

# Full wwPDB X-ray Structure Validation Report (i)

#### Jan 28, 2024 - 02:18 PM EST

PDB ID	:	1EGU
Title	:	CRYSTAL STRUCTURE OF STREPTOCOCCUS PNEUMONIAE
		HYALURONATE LYASE AT 1.56 A RESOLUTION
Authors	:	Li, S.; Kelly, S.J.; Lamani, E.; Ferraroni, M.; Jedrzejas, M.J.
Deposited on	:	2000-02-16
Resolution	:	1.56  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
R <sub>free</sub>	130704	1483 (1.56-1.56)		
Clashscore	141614	1529 (1.56-1.56)		
Ramachandran outliers	138981	1498 (1.56-1.56)		
Sidechain outliers	138945	1495 (1.56-1.56)		
RSRZ outliers	127900	1465 (1.56-1.56)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			6%		
1	А	731	77%	19%	• •



#### $1\mathrm{EGU}$

## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 6454 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called HYALURONATE LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	721	Total 5794	C 3643	N 971	0 1158	S 22	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	731	VAL	GLY	SEE REMARK 999	UNP Q54873
А	893	HIS	-	expression tag	UNP Q54873
А	894	HIS	-	expression tag	UNP Q54873
А	895	HIS	-	expression tag	UNP Q54873
А	896	HIS	-	expression tag	UNP Q54873
А	897	HIS	-	expression tag	UNP Q54873
А	898	HIS	-	expression tag	UNP Q54873

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	650	Total O 650 650	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: HYALURONATE LYASE



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	84.04Å 104.28Å 101.76Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	20.00 - 1.56	Depositor
Resolution (A)	38.98 - 1.56	EDS
% Data completeness	98.6 (20.00-1.56)	Depositor
(in resolution range)	98.0(38.98-1.56)	EDS
R <sub>merge</sub>	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.85 (at 1.56Å)	Xtriage
Refinement program	XTALVIEW, X-PLOR 3.843	Depositor
D D.	0.200 , $0.236$	Depositor
$\Pi, \Pi_{free}$	0.218 , $0.245$	DCC
$R_{free}$ test set	1232 reflections $(0.99%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	15.3	Xtriage
Anisotropy	0.584	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.41, $54.2$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.011 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6454	wwPDB-VP
Average B, all atoms $(Å^2)$	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.33	0/5913	0.56	1/7984~(0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^{o})$	$Ideal(^{o})$
1	А	220	ALA	N-CA-C	-5.23	96.89	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5794	0	5605	115	1
2	А	10	0	0	0	0
3	А	650	0	0	18	0
All	All	6454	0	5605	115	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



1EGU	

Atom 1 Atom 2		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:456:MET:SD	3:A:1424:HOH:O	2.15	1.02
1:A:458:MET:SD	3:A:1423:HOH:O	2.18	1.02
1:A:267:ARG:HD2	3:A:1267:HOH:O	1.67	0.94
1:A:219:GLN:HB3	1:A:222:ARG:HB3	1.53	0.91
1:A:613:ASN:H	1:A:698:GLN:HE22	1.13	0.91
1:A:581:LYS:HB3	1:A:768:GLU:HB2	1.55	0.87
1:A:864:GLU:HB2	1:A:869:LYS:NZ	1.91	0.85
1:A:219:GLN:CB	1:A:222:ARG:HB3	2.10	0.82
1:A:222:ARG:HG3	1:A:223:ILE:N	1.96	0.78
1:A:864:GLU:HB2	1:A:869:LYS:HZ1	1.49	0.75
1:A:491:GLY:O	1:A:495:GLN:HG3	1.87	0.75
1:A:845:SER:HB2	1:A:853:VAL:HG23	1.67	0.75
1:A:391:TYR:CD2	1:A:549:LYS:HG3	2.27	0.69
1:A:217:SER:HB2	1:A:219:GLN:HG2	1.73	0.68
1:A:855:LYS:HD3	1:A:886:VAL:HG12	1.79	0.65
1:A:219:GLN:HB2	1:A:222:ARG:H	1.61	0.64
1:A:217:SER:HB3	1:A:222:ARG:HE	1.62	0.63
1:A:862:ARG:HD3	1:A:864:GLU:OE1	1.99	0.63
1:A:222:ARG:HG3	1:A:224:TYR:H	1.62	0.63
1:A:373:THR:O	1:A:377:ILE:HG12	1.99	0.62
1:A:217:SER:O	1:A:219:GLN:HG3	2.01	0.61
1:A:327:LYS:HE2	3:A:1331:HOH:O	2.01	0.61
1:A:660:TRP:HA	3:A:1108:HOH:O	2.02	0.60
1:A:446:PHE:HD1	3:A:1468:HOH:O	1.85	0.59
1:A:458:MET:HE2	1:A:567:SER:HB2	1.83	0.59
1:A:229:PHE:HE2	1:A:244:LYS:HE3	1.66	0.59
1:A:291:TRP:O	1:A:295:GLU:HG3	2.03	0.59
1:A:650:ASN:HD21	1:A:832:GLN:HE22	1.50	0.59
1:A:422:VAL:HA	3:A:1224:HOH:O	2.02	0.58
1:A:705:ASN:HB3	3:A:1173:HOH:O	2.03	0.58
1:A:222:ARG:HG2	1:A:224:TYR:O	2.03	0.58
1:A:217:SER:CB	1:A:222:ARG:HE	2.16	0.57
1:A:283:SER:O	1:A:327:LYS:HE3	2.04	0.57
1:A:355:ARG:HH11	1:A:418:GLN:NE2	2.03	0.56
1:A:864:GLU:HB2	1:A:869:LYS:HZ3	1.66	0.56
1:A:426:THR:O	1:A:429:PRO:HD3	2.06	0.56
1:A:610:PRO:HG3	1:A:763:TRP:CE2	2.42	0.55
1:A:495:GLN:HG2	3:A:1452:HOH:O	2.07	0.55
1:A:579:MET:HG3	3:A:1317:HOH:O	2.08	0.54
1:A:549:LYS:CE	3:A:1136:HOH:O	2.57	0.53
1:A:820:ASN:HD22	1:A:825:GLN:HG2	1.73	0.53
1:A:254:ASN:C	1:A:254:ASN:HD22	2.12	0.52



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:314:SER:HB2	3:A:1278:HOH:O	2.09	0.52
1:A:282:ASN:HD22	1:A:284:GLU:N	2.07	0.52
1:A:874:ASN:C	1:A:874:ASN:HD22	2.14	0.51
1:A:219:GLN:HB2	1:A:222:ARG:N	2.24	0.51
1:A:888:LYS:O	1:A:889:LYS:HB2	2.09	0.51
1:A:456:MET:CE	1:A:565:GLY:HA3	2.41	0.51
1:A:584:LYS:HE3	1:A:629:ASP:HB3	1.93	0.50
1:A:579:MET:HG2	3:A:1316:HOH:O	2.12	0.49
1:A:584:LYS:HE2	3:A:1505:HOH:O	2.13	0.49
1:A:219:GLN:NE2	1:A:222:ARG:H	2.10	0.49
1:A:282:ASN:HD22	1:A:284:GLU:H	1.59	0.49
1:A:521:LYS:HE3	1:A:525:LEU:HG	1.94	0.49
1:A:681:GLY:O	1:A:792:SER:HB2	2.12	0.49
1:A:173:THR:O	1:A:176:ASP:HB2	2.12	0.49
1:A:235:SER:HB2	1:A:293:ASP:HB2	1.94	0.49
1:A:273:MET:O	1:A:277:HIS:HB3	2.13	0.48
1:A:727:GLU:OE2	1:A:749:LYS:HD2	2.13	0.48
1:A:316:GLU:H	1:A:316:GLU:CD	2.16	0.48
1:A:868:TYR:H	1:A:893:HIS:CD2	2.32	0.48
1:A:222:ARG:HG3	1:A:224:TYR:N	2.28	0.47
1:A:233:LYS:HB2	1:A:233:LYS:NZ	2.29	0.47
1:A:404:TYR:CE1	1:A:461:GLY:HA3	2.49	0.47
1:A:848:SER:O	1:A:850:GLN:HG3	2.14	0.47
1:A:314:SER:HB3	1:A:317:GLU:OE1	2.15	0.47
1:A:456:MET:HE3	1:A:565:GLY:HA3	1.97	0.47
1:A:456:MET:CE	1:A:597:TYR:HE2	2.28	0.47
1:A:607:GLY:C	1:A:610:PRO:HD2	2.34	0.47
1:A:882:PRO:HB2	1:A:884:GLN:NE2	2.30	0.46
1:A:354:GLY:O	1:A:358:VAL:HB	2.15	0.46
1:A:640:PHE:CD1	1:A:875:PRO:HG2	2.52	0.45
1:A:708:LYS:HE3	1:A:710:TYR:OH	2.17	0.45
1:A:213:LEU:HD11	1:A:265:VAL:HG22	1.98	0.45
1:A:868:TYR:O	1:A:893:HIS:HB3	2.17	0.45
1:A:219:GLN:HG3	1:A:219:GLN:H	1.41	0.45
1:A:314:SER:CB	3:A:1278:HOH:O	2.64	0.45
1:A:196:ASP:HB2	3:A:1445:HOH:O	2.17	0.44
1:A:254:ASN:ND2	1:A:256:SER:H	2.16	0.44
1:A:663:THR:HB	1:A:688:SER:HB3	1.99	0.44
1:A:295:GLU:HB3	1:A:329:VAL:HG22	1.99	0.44
1:A:447:ALA:HB3	1:A:448:PRO:HD3	1.99	0.44
1:A:623:THR:HA	1:A:691:THR:O	2.18	0.44



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	$distance ( m \AA)$	overlap (Å)
1:A:222:ARG:CG	1:A:223:ILE:N	2.75	0.44
1:A:255:PRO:HA	1:A:260:TYR:CG	2.53	0.44
1:A:269:VAL:O	1:A:273:MET:HG2	2.18	0.43
1:A:357:LYS:HB3	1:A:357:LYS:HE3	1.84	0.43
1:A:529:LEU:HD21	1:A:535:VAL:HG11	2.00	0.43
1:A:222:ARG:NH1	1:A:224:TYR:CZ	2.86	0.43
1:A:458:MET:HE2	1:A:458:MET:HB3	1.87	0.43
1:A:217:SER:HB2	1:A:219:GLN:CG	2.43	0.43
1:A:623:THR:HB	1:A:690:ASP:HB2	2.01	0.42
1:A:223:ILE:N	1:A:223:ILE:HD13	2.35	0.42
1:A:867:GLU:HA	1:A:893:HIS:NE2	2.34	0.42
1:A:647:ASP:OD2	1:A:862:ARG:NH1	2.53	0.42
1:A:661:ASN:H	1:A:661:ASN:HD22	1.68	0.42
1:A:674:LYS:HE3	3:A:1021:HOH:O	2.20	0.42
1:A:213:LEU:O	1:A:216:ILE:HG22	2.20	0.42
1:A:476:VAL:HG11	1:A:516:ASN:HB3	2.02	0.41
1:A:836:GLY:C	1:A:837:ILE:HG13	2.40	0.41
1:A:420:LEU:HD13	1:A:435:MET:CE	2.50	0.41
1:A:667:HIS:HD2	3:A:1052:HOH:O	2.03	0.41
1:A:738:SER:O	1:A:800:ASN:HA	2.20	0.41
1:A:219:GLN:C	1:A:221:ASP:H	2.24	0.41
1:A:580:ASN:O	1:A:581:LYS:HB2	2.20	0.41
1:A:621:THR:C	1:A:622:GLU:HG2	2.40	0.41
1:A:745:PHE:HB2	1:A:810:ILE:HG22	2.03	0.41
1:A:374:ILE:O	1:A:378:GLU:HG3	2.21	0.41
1:A:633:GLY:C	1:A:634:LYS:HG2	2.40	0.41
1:A:549:LYS:HD3	1:A:549:LYS:HA	1.62	0.40
1:A:254:ASN:C	1:A:254:ASN:ND2	2.74	0.40
1:A:888:LYS:O	1:A:889:LYS:CB	2.70	0.40
1:A:661:ASN:HD22	1:A:661:ASN:C	2.25	0.40
1:A:760:LYS:HE2	1:A:774:GLU:OE2	2.22	0.40
1:A:664:LEU:HA	1:A:685:GLN:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:316:GLU:OE2	1:A:867:GLU:OE1[3_555]	1.96	0.24



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	717/731~(98%)	685~(96%)	30 (4%)	2 (0%)	41	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	176	ASP
1	А	231	ASN

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	640/649~(99%)	595~(93%)	45~(7%)	15 1

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	171	LYS
1	А	196	ASP
1	А	214	SER
1	А	219	GLN
1	А	222	ARG
1	А	223	ILE
1	А	228	LYS
1	А	233	LYS
1	А	254	ASN



Mol	Chain	Res	Type
1	А	263	GLU
1	А	270	ARG
1	А	278	LYS
1	А	279	HIS
1	А	282	ASN
1	А	314	SER
1	А	319	LYS
1	А	333	GLU
1	А	337	LYS
1	А	368	GLN
1	А	375	ARG
1	А	380	VAL
1	А	385	ASP
1	А	404	TYR
1	А	425	LYS
1	А	432	LYS
1	А	433	ASP
1	А	444	LYS
1	А	495	GLN
1	А	496	ARG
1	А	529	LEU
1	А	549	LYS
1	А	579	MET
1	А	581	LYS
1	А	626	LYS
1	А	661	ASN
1	А	715	GLU
1	A	717	SER
1	А	748	LYS
1	А	773	LYS
1	А	854	LEU
1	А	867	GLU
1	A	869	LYS
1	А	874	ASN
1	А	888	LYS
1	А	889	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	202	ASN
1	А	219	GLN



Mol	Chain	Res	Type
1	А	237	ASN
1	А	254	ASN
1	А	261	GLN
1	А	282	ASN
1	А	368	GLN
1	А	386	GLN
1	А	418	GLN
1	А	495	GLN
1	А	661	ASN
1	А	667	HIS
1	А	698	GLN
1	А	759	GLN
1	А	788	GLN
1	А	808	GLN
1	А	820	ASN
1	А	825	GLN
1	А	832	GLN
1	А	874	ASN
1	А	893	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	SO4	А	2003	-	$4,\!4,\!4$	1.10	0	$6,\!6,\!6$	0.30	0
2	SO4	А	2002	-	4,4,4	1.13	0	$6,\!6,\!6$	0.27	0

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	721/731~(98%)	0.51	44 (6%) 21 25	14, 23, 45, 76	0

All (44) RSRZ outliers are listed below:

Mol	Mol Chain		Type	RSRZ
1	А	219	GLN	7.6
1	А	218	SER	7.1
1	А	893	HIS	6.8
1	А	223	ILE	6.3
1	А	220	ALA	5.9
1	А	867	GLU	4.8
1	А	279	HIS	4.8
1	А	224	TYR	4.7
1	А	175	THR	4.6
1	А	221	ASP	4.5
1	А	275	TRP	4.2
1	А	375	ARG	4.1
1	А	866	ASP	4.0
1	А	288	VAL	3.7
1	А	233	LYS	3.6
1	А	892	GLN	3.5
1	А	231	ASN	3.3
1	А	222	ARG	3.3
1	А	174	TYR	3.2
1	А	171	LYS	3.0
1	А	340	ASP	3.0
1	А	427	LYS	2.9
1	А	234	THR	2.8
1	А	230	SER	2.8
1	А	428	ASN	2.7
1	А	368	GLN	2.7
1	А	281	TYR	2.6



Mol	Chain	Res	Type	RSRZ
1	А	229	PHE	2.5
1	А	328	PHE	2.5
1	А	232	TYR	2.4
1	А	341	ASN	2.4
1	А	325	ILE	2.4
1	А	338	THR	2.3
1	А	358	VAL	2.3
1	А	865	GLY	2.3
1	А	797	LEU	2.2
1	А	217	SER	2.2
1	А	228	LYS	2.2
1	А	889	LYS	2.2
1	А	433	ASP	2.2
1	А	196	ASP	2.2
1	А	216	ILE	2.1
1	А	356	VAL	2.1
1	А	568	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	SO4	А	2003	5/5	0.91	0.12	40,40,40,40	0
2	SO4	А	2002	5/5	0.94	0.10	40,40,40,40	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

