

# wwPDB X-ray Structure Validation Summary Report (i)

#### Mar 10, 2024 – 01:22 AM EST

PDB ID	:	4EIW
Title	:	Whole cytosolic region of atp-dependent metalloprotease FtsH (G399L)
Authors	:	Suno, R.; Niwa, H.; Tsuchiya, D.; Yoshida, M.; Morikawa, K.
Deposited on	:	2012-04-06
Resolution	:	3.90  Å(reported)
	-	

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\#$ Entries)	Similar resolution $(\#Entries, resolution range(Å))$
R <sub>free</sub>	130704	1002 (4.14-3.66)
Clashscore	141614	1004 (4.12-3.68)
Ramachandran outliers	138981	1021 (4.14-3.66)
Sidechain outliers	138945	1014 (4.14-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain												
1	А	508	16%	49%	23%	• 10%									
1	В	508	9%	53%	22%	• 12%									
1	С	508	16%	49%	21%	• 10%									
1	D	508	9%	52%	23%	• 12%									
1	Е	508	17%	49%	22%	• 10%									
1	F	508	10%	52%	23%	• 12%									

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ADP	В	2001	-	-	Х	-
2	ADP	С	1001	-	-	Х	-
2	ADP	D	2001	-	-	Х	-
2	ADP	Е	1001	-	-	Х	-
2	ADP	F	2001	_	_	Х	-

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



#### 4EIW

# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 21429 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	458	Total	С	Ν	0	S	0	0	0
	A	400	3578	2245	658	662	13	0	0	0
1	р	446	Total	С	Ν	0	S	0	0	0
	D	440	3511	2206	641	651	13	0	0	0
1	C	159	Total	С	Ν	0	S	0	0	0
	U	400	3578	2245	658	662	13	0	0	0
1	П	446	Total	С	Ν	0	S	0	0	0
	D	440	3511	2206	641	651	13	0	0	0
1	Б	159	Total	С	Ν	Ο	S	0	0	0
	E	400	3578	2245	658	662	13	0	0	0
1	Б	446	Total	С	Ν	Ο	S	0	0	0
	.   F'	440	3511	2206	641	651	13	U	U	

• Molecule 1 is a protein called ATP-dependent zinc metalloprotease FtsH.

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	117	GLY	-	expression tag	UNP Q5SI82
А	118	PRO	-	expression tag	UNP Q5SI82
А	119	LEU	-	expression tag	UNP Q5SI82
А	120	GLY	-	expression tag	UNP Q5SI82
А	121	SER	-	expression tag	UNP Q5SI82
А	122	HIS	-	expression tag	UNP Q5SI82
А	123	MET	-	expression tag	UNP Q5SI82
А	124	GLY	-	expression tag	UNP Q5SI82
А	125	ALA	-	expression tag	UNP Q5SI82
А	399	LEU	GLY	engineered mutation	UNP Q5SI82
В	117	GLY	-	expression tag	UNP Q5SI82
В	118	PRO	-	expression tag	UNP Q5SI82
В	119	LEU	-	expression tag	UNP Q5SI82
В	120	GLY	-	expression tag	UNP Q5SI82
В	121	SER	-	expression tag	UNP Q5SI82
В	122	HIS	-	expression tag	UNP Q5SI82
В	123	MET	-	expression tag	UNP Q5SI82

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference						
В	124	GLY	-	expression tag	UNP Q5SI82						
В	125	ALA	-	expression tag	UNP Q5SI82						
В	399	LEU	GLY	engineered mutation	UNP Q5SI82						
С	117	GLY	-	expression tag	UNP Q5SI82						
С	118	PRO	-	expression tag	UNP Q5SI82						
С	119	LEU	-	expression tag	UNP Q5SI82						
С	120	GLY	-	expression tag	UNP Q5SI82						
С	121	SER	-	expression tag	UNP Q5SI82						
С	122	HIS	-	expression tag	UNP Q5SI82						
С	123	MET	-	expression tag	UNP Q5SI82						
С	124	GLY	-	expression tag	UNP Q5SI82						
C	125	ALA	-	expression tag	UNP Q5SI82						
C	399	LEU	GLY	engineered mutation	UNP Q5SI82						
D	117	GLY	-	expression tag	UNP Q5SI82						
D	118	PRO	-	expression tag	UNP Q5SI82						
D	119	LEU	-	expression tag	UNP Q5SI82						
D	120	GLY	-	expression tag	UNP Q5SI82						
D	121	SER	-	expression tag	UNP Q5SI82						
D	122	HIS	-	expression tag	UNP Q5SI82						
D	123	MET	-	expression tag	UNP Q5SI82						
D	124	GLY	-	expression tag	UNP Q5SI82						
D	125	ALA	-	expression tag	UNP Q5SI82						
D	399	LEU	GLY	engineered mutation	UNP Q5SI82						
E	117	GLY	-	expression tag	UNP Q5SI82						
E	118	PRO	-	expression tag	UNP Q5SI82						
E	119	LEU	-	expression tag	UNP Q5SI82						
E	120	GLY	-	expression tag	UNP Q5SI82						
E	121	SER	-	expression tag	UNP Q5SI82						
E	122	HIS	-	expression tag	UNP Q5SI82						
E	123	MET	-	expression tag	UNP Q5SI82						
E	124	GLY	-	expression tag	UNP Q5SI82						
E	125	ALA	-	expression tag	UNP Q5SI82						
E	399	LEU	GLY	engineered mutation	UNP Q5SI82						
F	117	GLY	-	expression tag	UNP Q5SI82						
F	118	PRO	-	expression tag	UNP Q5SI82						
F	119	LEU	-	expression tag	UNP Q5SI82						
F	120	GLY	-	expression tag	UNP Q5SI82						
F	121	SER	-	expression tag	UNP Q5SI82						
F	122	HIS	-	expression tag	UNP Q5SI82						
F	123	MET	-	expression tag	UNP Q5SI82						
F	124	GLY	-	expression tag	UNP Q5SI82						
F	125	ALA	-	expression tag	UNP Q5SI82						

Continued from previous page...

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference					
F	399	LEU	GLY	engineered mutation	UNP Q5SI82					

• Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues		Ate	oms	ZeroOcc	AltConf		
0	Δ	1	Total	С	Ν	Ο	Р	0	0
	A	L	27	10	5	10	2	0	0
0	В	1	Total	С	Ν	Ο	Р	0	0
	D	I	27	10	5	10	2	0	0
9	С	1	Total	С	Ν	Ο	Р	0	0
	U	I	27	10	5	10	2	0	0
9	Л	1	Total	С	Ν	Ο	Р	0	0
	D	I	27	10	5	10	2	0	0
9	F	1	Total	С	Ν	Ο	Р	0	0
	Ľ	I	27	10	5	10	2	0	0
9	F	1	Total	С	Ν	Ο	Р	0	0
	2 F		27	10	5	10	2	0	



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: ATP-dependent zinc metalloprotease FtsH

 $\bullet$  Molecule 1: ATP-dependent zinc metalloprotease FtsH









#### GLY ALA

 $\bullet$  Molecule 1: ATP-dependent zinc metalloprotease FtsH



• Molecule 1: ATP-dependent zinc metalloprotease FtsH



# A497 D433 A501 V431 A501 V433 A502 V433 A503 P4436 A504 P4436 A503 P4436 A504 P4436 A505 P4436 A506 P4436 A509 P4436 A516 P4444 A516 P4446 A516 P4446 A516 P4446 A516 P4446 A516 P445 A528 A546 A533 A546

 $\bullet$  Molecule 1: ATP-dependent zinc metalloprotease FtsH

Chai	in	F:		10	%										5	529	6												2	23%	6				•		1	2%	6				
GLY PRO LEU	GLY SER	HIS MET	GLY	ALA ARG	ASN	GLY	ALA	GLY	PRO	SER	SER	ALA	PHE	SER	PHE THR	LYS	SER	ARG	ALA	VAL	L147	T148	E149 A150	P151	K152	V153 T154	F155	K156	D157 V158	A159	G160	A161	E162	A164	K165	E166 E167	E167	K169	E170	1171	V172	F174	L175 K176
N177 P178 S179	R180 F181	H182 E183	M184	6185 A186	R187	I 188	K190	G191	V192	L193	L194 V195	G196	P197	P198	G199 V200	G201	K202	T203	H204	A206	R207	A208 11000	4210	G211	E212	A213 R214	V215	P216	F217 T218	T219	A220	S221	8222 8223	D224	F225	V226	MODR	F229	V230	G231	V232	A234	<mark>A235</mark> R236
V237 R238 D239	L240 F241	E242 T243	A244	K245 R246	H247	A248 D240	C250	1251	V252	F253	1254 D255	E256	1257	D258	A259 V260	G261	R262	K263	AKG GI V	SER	GLY	VAL	GLY	GLY	N272	D273 F274	R275	E276	ц277 тотв	1279 L279	N280	Q281	L282	V284	E285	M286	028/ C288	F289	E290	K291	D292	A294	1295 V296
97 98 99	<mark>0 1</mark>	02	4	a g	07	08		11	12	13	15	16	17	8	<b>6</b>	2	23	24		27	58	5 <mark>0</mark>	00 10 10	32	33	35	36	37	20 C	10	<u>1</u> 1	<u>12</u>	14	15	<u>16</u>	47 40		00	51	52	0 3 0 3	55	57
V29 M29 A29	T3(	N30 R30	P30		L3(	D3(	A3:	L3:	L3	R3	0.00	R3	F3	D3:	R. D.	) F	I3;	D3:	A3.	D3	V3:	K3	сэ ВЗ	E3	03		R3	I3	H3.	R3	C3	K3	1.30	A34	E3	D3		L3	A3	L3	L3	K3	T3
P358 G359 F360	V361 G362	A363 D364	L365	E366 N367	L368	L369	E371	A372	A373	L374	A376	A377	R378	E379	G380 B381	R382	K383	1384 2005	1385 M386	K387	D388	L389	E391	A392	A393	D394 R395	V396	M397	M398 1300	P400	A401	K402	8403 8404	L405	V406	L407	R410	D411	R412	R413	I414	Y417	H418 E419
A420 G421	L424 A425	A426 H427	F428	E430	H431	A432	G434	V435	H436	K437	V430 T439	1440	V441	P442	R443 C444	R445	A446	L447	6448 FAAO	M450	M451	P452	R453 R454	E455	D456	M457 L458	H459	W460	S461 BA67	K463	R464	L465	L466 D467	Q468	1469	A470	V4/1 4479	L473	A474	G475	R476	A478	E479 E480
1481 V482 F483	D484 D485	V486 T487	T488	6489 A490	E491	N492	F494	R495	Q496	A497	1490 E499	L500	A501	R502	R503 M504	I 505	T506	E507	008 2500	M510	H511	P512	E513		V517	A518 Y519	A520	V521	R522 FE73	D524	T525	Y526	L527 G528	G529	Y530	D531	P533	0534	Y535	<b>S536</b>	E537 E536	T539	A540 K541
D544	V547 R548	R549 1.550	I551	E553	Q554	Y555	R557	V558	K559	A560	L562	L563	E564	K565	R566 F567	V568	L569	E570	K5/1 VE70		T575	L576	E578	R579	E580	1581 1.582	T583	A584	E585 F586	F587	<mark>Q588</mark>	R589	V590	E592	G593	L594	1.596	E597	A5 98	P599	E600	ALA	ARG GLU
GLU ARG GLU	PR0 PR0	ARG VAL	VAL	LYS	VAL	LYS	GLY	GLY	ALA	LEU	GL.Y GL.Y	ALA																															



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants	146.15Å 146.15Å 349.06Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	71.53 - 3.90	Depositor
Resolution (A)	71.53 - 3.90	EDS
% Data completeness	97.1 (71.53-3.90)	Depositor
(in resolution range)	97.1 (71.53-3.90)	EDS
R <sub>merge</sub>	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	7.46 (at 3.89Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
D D.	0.299 , $0.312$	Depositor
$\Pi, \Pi_{free}$	0.298 , $0.309$	DCC
$R_{free}$ test set	1967 reflections $(5.01\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	99.1	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent $k_{sol}(e/A^3)$ , $B_{sol}(A^2)$	0.17, 19.0	EDS
L-test for twinning <sup>2</sup>	$< L >=0.27, < L^2>=0.11$	Xtriage
Estimated twinning fraction	0.237 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.78	EDS
Total number of atoms	21429	wwPDB-VP
Average B, all atoms $(Å^2)$	112.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 1.77% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles		
Moi Chain		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.60	3/3636~(0.1%)	0.95	16/4906~(0.3%)	
1	В	0.63	3/3568~(0.1%)	1.01	18/4815~(0.4%)	
1	С	0.59	2/3636~(0.1%)	1.03	24/4906~(0.5%)	
1	D	0.60	2/3568~(0.1%)	0.98	17/4815~(0.4%)	
1	Е	0.61	6/3636~(0.2%)	0.96	13/4906~(0.3%)	
1	F	0.58	2/3568~(0.1%)	0.96	12/4815~(0.2%)	
All	All	0.60	18/21612~(0.1%)	0.98	100/29163~(0.3%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2
1	В	0	5
1	С	0	1
1	D	0	6
1	Ε	0	1
1	F	0	5
All	All	0	20

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	В	586	GLU	CD-OE1	-12.04	1.12	1.25
1	С	214	ARG	CZ-NH2	8.16	1.43	1.33
1	Е	214	ARG	CZ-NH2	-8.12	1.22	1.33
1	Е	319	ARG	CZ-NH1	-7.39	1.23	1.33
1	D	586	GLU	CD-OE1	-6.72	1.18	1.25



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	Ε	319	ARG	NE-CZ-NH2	18.11	129.36	120.30
1	С	207	ARG	NE-CZ-NH1	-15.37	112.61	120.30
1	А	316	ARG	NE-CZ-NH1	-15.14	112.73	120.30
1	В	236	ARG	NE-CZ-NH1	-14.64	112.98	120.30
1	С	207	ARG	NE-CZ-NH2	13.93	127.26	120.30

The worst 5 of 100 bond angle outliers are listed below:

There are no chirality outliers.

5 of 20 planarity outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	Group
1	А	213	ALA	Peptide
1	А	532	VAL	Peptide
1	В	244	ALA	Peptide
1	В	288	GLY	Peptide
1	В	382	ARG	Peptide

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3578	0	3623	862	5
1	В	3511	0	3556	851	3
1	С	3578	0	3623	815	0
1	D	3511	0	3556	851	3
1	Е	3578	0	3623	805	3
1	F	3511	0	3556	857	2
2	А	27	0	12	8	0
2	В	27	0	12	9	0
2	С	27	0	12	10	0
2	D	27	0	12	9	0
2	Ε	27	0	12	11	0
2	F	27	0	12	10	0
All	All	21429	0	21609	4895	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 114.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:416:ALA:CB	1:E:577:LEU:HD23	1.18	1.63
1:A:416:ALA:HB3	1:A:577:LEU:CD2	1.33	1.58
1:A:416:ALA:CB	1:A:577:LEU:HD23	1.15	1.55
1:F:376:ALA:CA	1:F:381:ARG:HD2	1.31	1.55
1:E:416:ALA:HB3	1:E:577:LEU:CD2	1.35	1.55

The worst 5 of 4895 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

The worst 5 of 8 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic}\\ {\rm distance}~({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:570:GLU:CG	1:D:382:ARG:NH2[3_564]	1.61	0.59
1:A:417:TYR:OH	1:D:382:ARG:NE[3_564]	1.74	0.46
1:A:177:ASN:OD1	1:E:214:ARG:NH2[6_665]	1.85	0.35
1:B:238:ARG:NE	1:F:378:ARG:NH2[6_665]	1.91	0.29
1:A:570:GLU:CB	1:D:382:ARG:NH2[3_564]	2.08	0.12

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Pe	erc	entiles
1	А	456/508~(90%)	323 (71%)	111 (24%)	22 (5%)		2	24
1	В	442/508~(87%)	289~(65%)	125 (28%)	28 (6%)		1	19
1	С	456/508~(90%)	323 (71%)	112 (25%)	21 (5%)		2	25
1	D	442/508~(87%)	288~(65%)	121 (27%)	33 (8%)		1	16
1	Ε	456/508~(90%)	324 (71%)	110 (24%)	22 (5%)		2	24
1	F	442/508 (87%)	288~(65%)	120 (27%)	34 (8%)		1	16
All	All	2694/3048 (88%)	1835 (68%)	699 (26%)	160 (6%)		1	20



5 of 160 Ramachandran outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	153	VAL
1	А	511	HIS
1	В	153	VAL
1	В	274	GLU
1	В	379	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	365/402~(91%)	240 (66%)	125~(34%)	0 1
1	В	361/402~(90%)	242 (67%)	119 (33%)	0 1
1	С	365/402~(91%)	240 (66%)	125 (34%)	0 1
1	D	361/402~(90%)	241 (67%)	120 (33%)	0 1
1	Ε	365/402~(91%)	244 (67%)	121 (33%)	0 1
1	F	361/402~(90%)	240 (66%)	121 (34%)	0 1
All	All	2178/2412 (90%)	1447 (66%)	731 (34%)	0 1

5 of 731 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	D	460	TRP
1	Е	439	THR
1	D	533	ARG
1	D	459	HIS
1	Е	264	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 37 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	468	GLN
1	F	338	HIS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	Е	496	GLN
1	F	281	GLN
1	В	431	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Tuno Chain Dec		Link	Bond lengths			Bond angles		
INIOI	туре	Unain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ADP	А	1001	-	24,29,29	1.10	2 (8%)	29,45,45	1.50	4 (13%)
2	ADP	С	1001	-	24,29,29	1.10	3 (12%)	29,45,45	1.56	4 (13%)
2	ADP	D	2001	-	24,29,29	1.03	3 (12%)	29,45,45	1.45	5 (17%)
2	ADP	В	2001	-	24,29,29	1.05	2 (8%)	29,45,45	1.49	4 (13%)
2	ADP	E	1001	-	24,29,29	1.06	2 (8%)	29,45,45	1.45	4 (13%)
2	ADP	F	2001	-	24,29,29	1.01	1 (4%)	29,45,45	1.52	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	А	1001	-	-	2/12/32/32	0/3/3/3
2	ADP	С	1001	-	-	2/12/32/32	0/3/3/3
2	ADP	D	2001	-	-	2/12/32/32	0/3/3/3
2	ADP	В	2001	-	-	3/12/32/32	0/3/3/3
2	ADP	Е	1001	-	-	2/12/32/32	0/3/3/3
2	ADP	F	2001	-	-	2/12/32/32	0/3/3/3

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
2	А	1001	ADP	C5-C4	2.78	1.48	1.40
2	F	2001	ADP	C5-C4	2.76	1.48	1.40
2	Е	1001	ADP	C5-C4	2.73	1.48	1.40
2	С	1001	ADP	C5-C4	2.62	1.47	1.40
2	В	2001	ADP	C5-C4	2.59	1.47	1.40

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	1001	ADP	PA-O3A-PB	-4.26	118.21	132.83
2	С	1001	ADP	PA-O3A-PB	-4.20	118.40	132.83
2	Е	1001	ADP	PA-O3A-PB	-4.05	118.92	132.83
2	В	2001	ADP	PA-O3A-PB	-3.99	119.13	132.83
2	С	1001	ADP	N3-C2-N1	-3.72	122.86	128.68

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	1001	ADP	C5'-O5'-PA-O1A
2	В	2001	ADP	C5'-O5'-PA-O3A
2	С	1001	ADP	C5'-O5'-PA-O1A
2	С	1001	ADP	C5'-O5'-PA-O3A
2	D	2001	ADP	C5'-O5'-PA-O3A

There are no ring outliers.

6 monomers are involved in 57 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	1001	ADP	8	0
2	С	1001	ADP	10	0
2	D	2001	ADP	9	0
2	В	2001	ADP	9	0
2	Е	1001	ADP	11	0
2	F	2001	ADP	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

## 6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













## 6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

