

Full wwPDB X-ray Structure Validation Report (i)

Jun 28, 2021 – 02:14 PM BST

PDB ID : 6EIL

Title: DYRK1A in complex with XMD8-49

Authors : Rothweiler, U. Deposited on : 2017-09-19

Resolution : 2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp

with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as 541 be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.20

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.20

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\mathring{\rm A})}) \end{array}$
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	368	88%	5%	% 7%
1	В	368	81%	11%	8%
1	С	368	7%	10%	10%
1	D	368	82%	8%	10%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 22259 atoms, of which 10817 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DYRK1A.


Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace	
1	Λ	342	Total	С	Н	N	О	Р	S	0	0	0
1	A	042	5581	1798	2787	477	501	1	17	0	0	U
1	В	340	Total	С	Н	N	О	Р	S	0	0	0
1	Ъ	340	5492	1774	2735	470	495	1	17	0		U
1	С	333	Total	С	Н	N	О	Р	S	0	0	0
1			5280	1724	2602	452	484	1	17	U		U
1	D	333	Total	С	Н	N	О	Р	S	0	0	0
1	ע	ეეე	5404	1752	2693	455	486	1	17	U	U	U

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	GLY	-	expression tag	UNP Q13627
A	124	ALA	_	expression tag	UNP Q13627
A	125	SER	_	expression tag	UNP Q13627
В	123	GLY	_	expression tag	UNP Q13627
В	124	ALA	-	expression tag	UNP Q13627
В	125	SER	_	expression tag	UNP Q13627
С	123	GLY	_	expression tag	UNP Q13627
С	124	ALA	_	expression tag	UNP Q13627
С	125	SER	_	expression tag	UNP Q13627
D	123	GLY	-	expression tag	UNP Q13627
D	124	ALA	_	expression tag	UNP Q13627
D	125	SER	-	expression tag	UNP Q13627

• Molecule 2 is [3-azanyl-6-(5-azanyl-2-methoxy-phenyl)pyrazin-2-yl]-pyridin-3-yl-methanone (three-letter code: B6B) (formula: $C_{17}H_{15}N_5O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
2	А	1	Total C N O	0	0	
	11	1	24 17 5 2	O		
9	D	1	Total C N O	0	0	
	2 B	1	24 17 5 2	0	0	
9	D	1	Total C N O	0	0	
	ש	1	24 17 5 2	U	U	


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	146	Total O 146 146	0	0
3	В	92	Total O 92 92	0	0
3	С	74	Total O 74 74	0	0
3	D	118	Total O 118 118	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	87.39Å 87.70Å 229.96Å	Danasitan
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.16 - 2.46	Depositor
resolution (A)	48.16 - 2.46	EDS
% Data completeness	96.2 (48.16-2.46)	Depositor
(in resolution range)	95.7 (48.16-2.46)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.94 (at 2.45Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
D D.	0.209 , 0.254	Depositor
R, R_{free}	0.213 , 0.254	DCC
R_{free} test set	3077 reflections $(4.94%)$	wwPDB-VP
Wilson B-factor (Å ²)	43.8	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38 , 29.1	EDS
L-test for twinning ²	$< L >=0.43, < L^2>=0.26$	Xtriage
Estimated twinning fraction	0.089 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	22259	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.84% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $^{^{1}}$ Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: B6B, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Moi Chain		$\mid \text{RMSZ} \mid \# Z > 5$		# Z > 5
1	A	0.27	0/2842	0.44	0/3833
1	В	0.26	0/2804	0.42	0/3785
1	С	0.26	0/2724	0.42	0/3686
1	D	0.27	0/2758	0.43	0/3723
All	All	0.27	0/11128	0.43	0/15027

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2794	2787	2787	12	1
1	В	2757	2735	2734	29	0
1	С	2678	2602	2601	21	0
1	D	2711	2693	2693	19	2
2	A	24	0	0	3	0
2	В	24	0	0	2	0
2	D	24	0	0	2	0
3	A	146	0	0	8	0
3	В	92	0	0	13	1



Continued from previous page...

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
3	С	74	0	0	9	0
3	D	118	0	0	10	0
All	All	11442	10817	10815	85	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		${\rm distance}\;({\rm \AA})$	overlap (Å)
1:B:286:CYS:SG	3:B:686:HOH:O	2.28	0.90
1:C:203:GLU:OE1	3:C:501:HOH:O	1.94	0.86
1:A:146:ASP:OD2	3:A:601:HOH:O	1.98	0.81
1:B:180:VAL:N	3:B:604:HOH:O	2.14	0.81
1:B:433:GLY:O	3:B:602:HOH:O	1.99	0.80
1:C:182:GLN:OE1	3:C:502:HOH:O	2.02	0.78
1:B:430:GLU:OE2	3:B:603:HOH:O	2.02	0.78
1:B:178:ASP:OD1	3:B:604:HOH:O	2.03	0.75
1:A:239:GLU:OE1	3:A:602:HOH:O	2.04	0.74
1:D:366:GLU:OE2	3:D:601:HOH:O	2.06	0.73
1:B:457:LEU:O	3:B:605:HOH:O	2.07	0.72
1:C:199:GLN:NE2	3:C:501:HOH:O	2.22	0.72
1:B:471:TYR:O	3:B:606:HOH:O	2.07	0.71
1:B:349:CYS:O	3:B:607:HOH:O	2.07	0.71
2:A:501:B6B:NAN	3:A:606:HOH:O	2.24	0.69
1:A:396:GLU:OE1	3:A:603:HOH:O	2.08	0.69
1:B:247:ASP:OD1	1:B:250:ARG:NH2	2.25	0.69
1:B:459:MET:SD	3:B:691:HOH:O	2.51	0.68
1:C:385:LEU:O	3:C:503:HOH:O	2.12	0.68
1:B:183:GLU:OE2	1:B:225:LYS:NZ	2.27	0.67
1:C:163:SER:OG	3:C:504:HOH:O	2.13	0.66
2:A:501:B6B:NAC	2:A:501:B6B:OAD	2.26	0.64
1:D:214:ASP:O	3:D:602:HOH:O	2.14	0.64
1:A:241:LEU:N	3:A:606:HOH:O	2.27	0.63
1:C:458:ARG:NH2	1:C:466:THR:O	2.33	0.62
1:C:404:ASN:OD1	1:C:405:LEU:N	2.35	0.60
1:D:429:VAL:O	3:D:603:HOH:O	2.16	0.60
1:A:408:THR:OG1	3:A:605:HOH:O	2.17	0.59
1:D:190:ILE:O	1:D:233:HIS:ND1	2.34	0.59
1:B:328:ARG:NH1	1:B:366:GLU:OE2	2.36	0.58
1:B:134:LYS:NZ	3:B:613:HOH:O	2.31	0.57



Continued from previous page...

Continuea from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ ({\rm \AA})$	$\text{overlap } (\mathring{\mathrm{A}})$
1:B:239:GLU:OE1	1:B:304:LYS:NZ	2.38	0.57
2:B:501:B6B:OAD	2:B:501:B6B:NAC	2.35	0.57
1:C:250:ARG:NH1	3:C:514:HOH:O	2.37	0.56
1:C:390:LYS:NZ	3:C:513:HOH:O	2.35	0.56
1:B:280:GLU:N	1:B:280:GLU:OE2	2.36	0.56
1:A:392:ARG:NH1	3:A:616:HOH:O	2.39	0.55
1:D:231:ARG:O	3:D:604:HOH:O	2.19	0.53
1:C:325:ARG:NH1	1:C:363:GLY:O	2.40	0.53
1:D:257:VAL:HG11	1:D:261:LEU:HD23	1.90	0.53
1:B:396:GLU:N	1:B:404:ASN:O	2.42	0.52
1:C:158:ARG:O	1:C:179:ARG:HG3	2.10	0.52
1:D:150:LYS:NZ	3:D:607:HOH:O	2.28	0.51
1:A:198:ASN:ND2	3:A:621:HOH:O	2.43	0.50
1:C:257:VAL:HG11	1:C:261:LEU:HD23	1.94	0.50
1:C:430:GLU:OE2	3:C:506:HOH:O	2.20	0.50
1:C:154:LYS:HA	1:C:159:TYR:O	2.11	0.50
1:C:324:SER:OG	3:C:505:HOH:O	2.19	0.49
1:D:199:GLN:NE2	3:D:622:HOH:O	2.45	0.49
2:D:501:B6B:OAD	2:D:501:B6B:NAC	2.42	0.49
1:C:215:THR:HG22	1:C:217:MET:H	1.77	0.48
1:B:229:MET:O	3:B:608:HOH:O	2.19	0.47
1:C:420:THR:O	1:C:422:LYS:N	2.47	0.47
1:D:421:ARG:NH1	3:D:616:HOH:O	2.41	0.47
1:D:216:GLU:HA	1:D:219:TYR:CE2	2.50	0.47
1:A:319:TYR:OH	1:B:198:ASN:ND2	2.44	0.46
1:D:392:ARG:NH2	3:D:626:HOH:O	2.48	0.46
1:B:336:MET:SD	1:B:387:GLN:OE1	2.74	0.46
1:D:479:PHE:O	3:D:605:HOH:O	2.21	0.45
1:B:179:ARG:N	3:B:604:HOH:O	2.50	0.45
1:C:249:LEU:HD21	1:C:354:MET:HA	1.98	0.45
1:B:387:GLN:OE1	1:B:387:GLN:C	2.55	0.45
1:C:463:ASP:HB3	1:C:466:THR:HG22	1.98	0.44
1:D:222:VAL:HG21	1:D:306:VAL:HG12	2.00	0.44
2:A:501:B6B:CAG	2:A:501:B6B:NAO	2.78	0.44
1:B:436:GLY:N	3:B:602:HOH:O	2.41	0.44
1:C:374:VAL:HG11	1:C:405:LEU:HD11	2.00	0.44
1:D:445:THR:HG23	1:D:448:ASP:H	1.83	0.44
1:A:388:ALA:HB3	1:A:391:ALA:HB2	2.01	0.43
1:B:248:LEU:HD12	1:B:295:LEU:HD11	1.99	0.43
1:A:214:ASP:HB2	1:A:275:PHE:CZ	2.53	0.43
2:B:501:B6B:NAO	2:B:501:B6B:CAG	2.80	0.43



Continued from previous page...

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance } (ext{Å}) \end{array}$	Clash overlap (Å)
2:D:501:B6B:NAO	2:D:501:B6B:CAG	2.78	0.43
1:B:248:LEU:HD21	1:B:257:VAL:HG11	2.00	0.43
1:D:201:GLN:NE2	3:D:630:HOH:O	2.52	0.42
1:B:400:ASP:N	1:B:400:ASP:OD1	2.52	0.42
1:B:245:LEU:O	1:B:249:LEU:HG	2.18	0.42
1:D:383:HIS:CE1	1:D:384:ILE:HG13	2.55	0.42
1:B:249:LEU:HD21	1:B:354:MET:HA	2.01	0.42
1:D:398:LEU:O	1:D:400:ASP:N	2.53	0.42
1:A:472:TYR:HH	1:D:383:HIS:HD1	1.67	0.41
1:B:430:GLU:HA	1:B:446:VAL:HG12	2.03	0.41
1:D:400:ASP:OD1	1:D:400:ASP:O	2.39	0.41
1:B:383:HIS:HB2	1:C:472:TYR:OH	2.21	0.40
1:A:334:LEU:HB3	1:A:388:ALA:HB1	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{array}{c} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{array}{c} ext{Clash} \ ext{overlap } (ext{Å}) \end{array}$
1:D:181:GLU:OE2	3:B:601:HOH:O[3_554]	2.17	0.03
1:A:154:LYS:HZ3	1:D:251:ASN:O[1_545]	1.58	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	$337/368 \ (92\%)$	321 (95%)	16 (5%)	0	100	100
1	В	$335/368 \; (91\%)$	311 (93%)	23 (7%)	1 (0%)	41	49
1	С	326/368~(89%)	310 (95%)	15 (5%)	1 (0%)	41	49
1	D	326/368~(89%)	310 (95%)	15 (5%)	1 (0%)	41	49
All	All	1324/1472 (90%)	1252 (95%)	69 (5%)	3 (0%)	47	57



All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	323	GLN
1	С	323	GLN
1	D	323	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	301/324 (93%)	300 (100%)	1 (0%)	92 95
1	В	295/324~(91%)	290 (98%)	5 (2%)	60 73
1	С	282/324~(87%)	277 (98%)	5 (2%)	59 71
1	D	292/324~(90%)	290 (99%)	2 (1%)	84 90
All	All	$1170/1296 \ (90\%)$	1157 (99%)	13 (1%)	73 82

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	414	GLU
1	В	212	LYS
1	В	296	CYS
1	В	400	ASP
1	В	462	TYR
1	В	477	SER
1	С	136	TYR
1	С	199	GLN
1	С	365	ASN
1	С	431	THR
1	С	437	ARG
1	D	286	CYS
1	D	390	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	В	213	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Во	nd leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	В	321	1	15,16,17	1.34	1 (6%)	19,22,24	0.50	0
1	PTR	С	321	1	15,16,17	1.20	1 (6%)	19,22,24	0.52	0
1	PTR	A	321	1	15,16,17	1.28	1 (6%)	19,22,24	0.49	0
1	PTR	D	321	1	15,16,17	1.21	1 (6%)	19,22,24	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	В	321	1	-	0/10/11/13	0/1/1/1
1	PTR	С	321	1	-	0/10/11/13	0/1/1/1
1	PTR	A	321	1	-	2/10/11/13	0/1/1/1
1	PTR	D	321	1	-	0/10/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	Ideal(A)
1	В	321	PTR	OH-CZ	-4.50	1.30	1.40
1	A	321	PTR	OH-CZ	-4.24	1.31	1.40



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
1	D	321	PTR	OH-CZ	-4.22	1.31	1.40
1	С	321	PTR	OH-CZ	-4.17	1.31	1.40

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

	Mol	Chain	Res	Type	Atoms
	1	Α	321	PTR	CZ-OH-P-O2P
Ī	1	A	321	PTR	CZ-OH-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bond lengths			Bond angles		
	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2 $ 5 (15%)$ $ 5 (15%)$
2	B6B	A	501	-	26,26,26	1.97	7 (26%)	32,36,36	1.79	5 (15%)
2	B6B	D	501	-	26,26,26	1.86	6 (23%)	32,36,36	1.71	5 (15%)
2	B6B	В	501	-	26,26,26	1.88	7 (26%)	32,36,36	1.79	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



'-' means no outliers of that kind were identified.

Mol	\mathbf{Type}	Chain	${ m Res}$	Link	Chirals	${f Torsions}$	Rings
2	B6B	A	501	-	-	0/14/14/14	0/3/3/3
2	B6B	D	501	-	-	0/14/14/14	0/3/3/3
2	B6B	В	501	-	-	2/14/14/14	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
2	A	501	B6B	CAT-CAQ	-5.48	1.40	1.49
2	A	501	B6B	CAX-CAQ	-5.26	1.40	1.50
2	В	501	B6B	CAT-CAQ	-5.13	1.40	1.49
2	D	501	B6B	CAX-CAQ	-4.92	1.40	1.50
2	D	501	B6B	CAT-CAQ	-4.87	1.41	1.49
2	В	501	B6B	CAX-CAQ	-4.83	1.40	1.50
2	A	501	B6B	CAW-CAU	-3.49	1.39	1.48
2	В	501	B6B	CAW-CAU	-3.41	1.39	1.48
2	D	501	B6B	CAW-CAU	-3.38	1.40	1.48
2	D	501	B6B	CAJ-NAM	3.19	1.41	1.34
2	В	501	B6B	CAJ-NAM	2.99	1.40	1.34
2	В	501	B6B	CAK-NAN	2.89	1.40	1.34
2	A	501	B6B	CAK-NAN	2.86	1.40	1.34
2	D	501	B6B	CAK-NAN	2.84	1.40	1.34
2	A	501	B6B	CAJ-NAM	2.83	1.40	1.34
2	A	501	B6B	CAX-CAS	-2.48	1.39	1.43
2	В	501	B6B	CAF-NAM	2.22	1.40	1.33
2	A	501	B6B	CAF-NAM	2.18	1.40	1.33
2	D	501	B6B	CAF-NAM	2.17	1.40	1.33
2	В	501	B6B	CAX-CAS	-2.05	1.39	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	В	501	B6B	CAA-OAP-CAV	-6.12	108.29	117.53
2	A	501	B6B	CAX-CAS-NAC	-5.88	115.94	122.84
2	A	501	B6B	CAA-OAP-CAV	-5.46	109.28	117.53
2	В	501	B6B	CAX-CAS-NAC	-5.23	116.71	122.84
2	D	501	B6B	CAX-CAS-NAC	-5.10	116.86	122.84
2	D	501	B6B	CAA-OAP-CAV	-5.05	109.91	117.53
2	В	501	B6B	CAX-NAO-CAU	2.81	122.38	118.19
2	D	501	B6B	CAX-NAO-CAU	2.78	122.34	118.19
2	A	501	B6B	CAX-NAO-CAU	2.48	121.88	118.19
2	В	501	B6B	CAU-CAK-NAN	-2.17	118.72	122.27



$\alpha \cdots \tau$	e	•	
Continued	trom	meaningile	maaa
-	110116	predidus	puyc

Mol	Chain	Res	Type	${f Atoms}$	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	501	B6B	CAT-CAJ-NAM	-2.15	120.32	123.49
2	D	501	B6B	OAP-CAV-CAI	-2.12	120.73	124.37
2	A	501	B6B	CAF-NAM-CAJ	2.01	120.32	116.85
2	D	501	B6B	CAU-CAK-NAN	-2.00	119.00	122.27

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	501	B6B	CAI-CAV-OAP-CAA
2	В	501	B6B	CAW-CAV-OAP-CAA


There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	B6B	3	0
2	D	501	B6B	2	0
2	В	501	B6B	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	${f Analysed}$	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	341/368~(92%)	0.20	5 (1%) 73 71	23, 37, 55, 86	0
1	В	$339/368 \ (92\%)$	0.63	20 (5%) 22 19	29, 51, 79, 99	0
1	С	$332/368 \ (90\%)$	0.57	27 (8%) 12 9	29, 54, 84, 91	0
1	D	332/368~(90%)	0.37	10 (3%) 50 46	28, 43, 67, 82	0
All	All	$1344/1472 \ (91\%)$	0.44	62 (4%) 32 30	23, 46, 75, 99	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	399	PRO	6.9
1	В	213	HIS	6.6
1	A	216	GLU	6.2
1	В	215	THR	5.6
1	D	429	VAL	4.8
1	С	378	GLY	4.6
1	С	429	VAL	4.5
1	С	395	PHE	4.3
1	В	216	GLU	4.1
1	С	472	TYR	4.0
1	В	401	GLY	4.0
1	В	398	LEU	3.8
1	A	215	THR	3.7
1	В	374	VAL	3.7
1	С	450	LEU	3.6
1	С	376	VAL	3.5
1	D	164	LEU	3.4
1	В	422	LYS	3.4
1	В	217	MET	3.3
1	В	394	PHE	3.2
1	С	421	ARG	3.2



Continued from previous page...

Mol	Chain	Res	$\overline{\text{Type}}$	RSRZ
1	С	423	LEU	3.2
1	С	425	ASN	3.1
1	В	420	THR	3.1
1	A	219	TYR	3.0
1	D	243	TYR	3.0
1	С	254	PHE	3.0
1	С	440	GLY	3.0
1	С	442	SER	3.0
1	С	404	ASN	2.9
1	В	245	LEU	2.8
1	С	420	THR	2.8
1	С	362	SER	2.7
1	D	219	TYR	2.7
1	D	398	LEU	2.7
1	В	277	ALA	2.6
1	С	477	SER	2.6
1	A	217	MET	2.6
1	В	418	PRO	2.5
1	В	333	LEU	2.5
1	D	214	ASP	2.5
1	С	295	LEU	2.5
1	С	303	ILE	2.4
1	В	219	TYR	2.3
1	С	334	LEU	2.3
1	С	448	ASP	2.3
1	С	459	MET	2.2
1	С	443	GLY	2.2
1	D	443	GLY	2.2
1	С	447	ALA	2.2
1	A	251	ASN	2.2
1	В	318	ILE	2.2
1	С	384	ILE	2.2
1	В	335	GLY	2.1
1	С	394	PHE	2.1
1	В	274	LEU	2.1
1	D	180	VAL	2.1
1	В	478	PHE	2.1
1	В	480	LYS	2.1
1	D	423	LEU	2.0
1	С	135	VAL	2.0
1	С	145	TYR	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q < 0.9
1	PTR	В	321	16/17	0.93	0.16	42,47,57,58	0
1	PTR	С	321	16/17	0.94	0.16	40,48,58,61	0
1	PTR	A	321	16/17	0.96	0.15	22,29,37,38	0
1	PTR	D	321	16/17	0.97	0.16	25,35,41,47	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f A}^2)$	Q<0.9
2	B6B	D	501	24/24	0.82	0.30	46,55,65,69	0
2	B6B	A	501	24/24	0.86	0.20	37,43,58,61	0
2	B6B	В	501	24/24	0.87	0.20	41,51,58,61	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














6.5 Other polymers (i)

There are no such residues in this entry.

