

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 17, 2023 – 10:15 AM EDT

PDB ID : 8EL3

Title : Light harvesting phycobiliprotein HaPE555 from the cryptophyte Hemiselmis

andersenii CCMP644 in a loose interface filament

Authors: Rathbone, H.W.; Michie, K.A.; Laos, A.L.; Curmi, P.M.G.

Deposited on : 2022-09-23

Resolution : 1.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$ 

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
$R_{free}$	130704	5534 (1.60-1.56)
Clashscore	141614	5861 (1.60-1.56)
Ramachandran outliers	138981	5708 (1.60-1.56)
Sidechain outliers	138945	5703 (1.60-1.56)
RSRZ outliers	127900	5431 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	67	88%	7% •
1	G	67	3%	
1	G		94% 3%	• •
1	J	67	94%	
1	K	67	94%	
2	В	177	96%	•



Mol	Chain	Length	Quality of chain	
2	D	177	94%	
2	F	177	97%	•••
2	Н	177	95%	
3	С	62	3% 97%	
3	E	62	3%	<u>.</u>
3	Ī	62	2%	6%
3	L	62	3%	070



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 20240 atoms, of which 9709 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Phycoerythrin alpha-1 subunit.

Mol	Chain	Residues		Α	tom	S			ZeroOcc	AltConf	Trace	
1	Λ	64	Total	С	Н	N	О	S	0	64	0	
1	A	04	926	287	461	82	91	5	0	04	U	
1	G	64	Total	С	Н	N	О	S	0	64	0	
1	G	04	926	287	461	82	91	5	0	04	0	
1	J	64	Total	С	Н	N	О	S	0	64	0	
1	J	04	924	287	459	82	91	5	0	04	0	
1	K	64	Total	С	Н	N	О	S	0	64	0	
1	17	04	925	287	460	82	91	5	U	04	U	

• Molecule 2 is a protein called Phycoerythrin550 beta subunit.

Mol	Chain	Residues			Atom	ıs		ZeroOcc	AltConf	Trace		
2	В	177	Total	С	Н	N	О	S	0	9	0	
2	Б	111	2687	825	1345	230	277	10	0	9	U	
2	D	173	Total	С	Н	N	О	S	0	8	0	
2	D	175	2626	807	1316	226	268	9	0	8	U	
2	F	173	Total	С	Н	N	О	S	0	10	0	
2	I'	175	2654	815	1329	227	274	9	0	10	U	
2	Н	173	Total	С	Н	N	О	S	0	Q	0	
	11	173	2615	804	1308	225	269	9	U	8	U	

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
В	172	VAL	GLU	$\operatorname{conflict}$	UNP U5T8W0
D	172	VAL	GLU	conflict	UNP U5T8W0
F	172	VAL	GLU	conflict	UNP U5T8W0
Н	172	VAL	GLU	conflict	UNP U5T8W0

• Molecule 3 is a protein called Phycoerythrin alpha-2 subunit.



Mol	Chain	Residues		Atoms						AltConf	Trace
3	С	62	Total	С	Н	N	О	S	0	62	0
)		02	917	280	459	79	93	6	0	02	
3	Е	62	Total	С	Н	N	О	S	0	62	0
)	15	02	917	280	459	79	93	6	0	02	
3	т	62	Total	С	Н	N	О	S	0	62	0
)	1	02	916	280	458	79	93	6	0	02	
2	Т	62	Total	С	Н	N	О	S	0	62	0
3	3   L	L 62		280	458	79	93	6	U	02	

 $\bullet$  Molecule 4 is PHYCOERYTHROBILIN (three-letter code: PEB) (formula:  $\mathrm{C}_{33}\mathrm{H}_{40}\mathrm{N}_4\mathrm{O}_6).$ 




Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
4	A	1	Total	С	Н	N	О	0	1
4	A	1	80	33	37	4	6	U	1
4	В	1	Total	С	Н	N	О	0	0
4	Б	1	80	33	37	4	6	0	
4	В	1	Total	С	Н	N	О	0	0
4	Б	1	80	33	37	4	6	U	0
4	С	1	Total	С	Η	N	О	0	1
4		1	80	33	37	4	6	U	1
4	D	1	Total	С	Η	N	О	0	0
4	D	1	80	33	37	4	6	0	
4	D	1	Total	С	Η	N	О	0	0
4	D	1	80	33	37	4	6	0	
4	E	1	Total	С	Н	N	О	0	1
4	ند	1	80	33	37	4	6	U	1
4	F	1	Total	С	Н	N	О	0	0
4	1'	1	80	33	37	4	6	U	0



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
4	F	1	Total	С	Н	N	О	0	0
4	I'	1	80	33	37	4	6	0	0
4	G	1	Total	С	Н	N	О	0	1
4	G	1	80	33	37	4	6	0	1
4	Н	1	Total	С	Н	N	О	0	0
4	11	1	80	33	37	4	6	0	0
4	Н	1	Total	С	Н	N	О	0	0
4	11	1	80	33	37	4	6	0	0
4	I	1	Total	С	Н	N	О	0	1
4	1	1	80	33	37	4	6	0	1
4	J	1	Total	С	Н	N	О	0	1
4	J	1	80	33	37	4	6	0	1
4	K	1	Total	С	Н	N	О	0	1
4	IX.	1	80	33	37	4	6	U	1
4	L	1	Total	С	Н	N	О	0	1
4	ь	1	80	33	37	4	6	U	1

 $\bullet$  Molecule 5 is DiCys-(15,16)-Dihydrobiliverdin (three-letter code: AX9) (formula:  $C_{33}H_{40}N_4O_6).$ 



Mol	Chain	Residues		Ato	oms		ZeroOcc	AltConf	
5	В	1	Total	С	Н	N	О	0	0
	Ъ	1	79	33	36	4	6	0	
5	D	1	Total	С	Н	N	О	0	0
9	ט	1	79	33	36	4	6	0	
5	Г	1	Total	С	Н	N	О	0	0
	Г	1	79	33	36	4	6	U	



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
5	П	1	Total	С	Н	N	О	0	0
5	п	1	79	33	36	4	6	U	0

#### • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	72	Total O 72 72	0	4
6	В	125	Total O 125 125	0	4
6	С	46	Total O 46 46	0	1
6	D	94	Total O 94 94	0	2
6	Е	39	Total O 39 39	0	2
6	F	133	Total O 133 133	0	3
6	G	58	Total O 58 58	0	3
6	Н	102	Total O 102 102	0	5
6	I	6	Total O 6 6	0	3
6	J	5	Total O 5 5	0	2
6	K	9	Total O 9 9	0	3
6	L	6	Total O 6 6	0	1



## 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Phycoerythrin alpha-1 subunit



• Molecule 2: Phycoerythrin 550 beta subunit







## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	64.80Å 76.84Å 103.36Å	Donositon
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $110.73^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	48.33 - 1.57	Depositor
Resolution (A)	48.33 - 1.57	EDS
% Data completeness	98.0 (48.33-1.57)	Depositor
(in resolution range)	98.0 (48.33-1.57)	EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.27 (at 1.57Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
D.D.	0.170 , 0.201	Depositor
$R, R_{free}$	0.167 , $0.196$	DCC
$R_{free}$ test set	1989 reflections (1.53%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.6	Xtriage
Anisotropy	0.216	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.45, 44.4	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	20240	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.46% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



 $<sup>^1 {\</sup>rm Intensities}$  estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LYZ, PEB, AX9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.56	0/459	0.80	0/609
1	G	0.49	0/459	0.71	0/609
1	J	0.51	0/459	0.67	0/609
1	K	0.51	0/459	0.69	0/609
2	В	0.49	0/1363	0.69	0/1838
2	D	0.47	0/1331	0.65	0/1794
2	F	0.50	0/1346	0.67	0/1815
2	Н	0.47	0/1328	0.64	0/1791
3	С	0.51	0/450	0.65	0/597
3	Е	0.50	0/450	0.68	0/597
3	I	0.54	0/450	0.78	0/597
3	L	0.49	0/450	0.70	0/597
All	All	0.50	0/9004	0.68	0/12062

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	465	461	449	3	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	465	461	450	1	0
1	J	465	459	451	1	0
1	K	465	460	448	1	0
2	В	1342	1345	1331	4	0
2	D	1310	1316	1303	3	0
2	F	1325	1329	1314	3	0
2	Н	1307	1308	1296	4	0
3	С	458	459	449	2	0
3	Е	458	459	449	1	0
3	I	458	458	448	3	0
3	L	458	458	450	1	0
4	A	43	37	37	2	0
4	В	86	74	74	4	0
4	С	43	37	37	1	0
4	D	86	74	74	8	0
4	Е	43	37	37	1	0
4	F	86	74	74	5	0
4	G	43	37	37	0	0
4	Н	86	74	74	1	0
4	I	43	37	37	2	0
4	J	43	37	37	1	0
4	K	43	37	37	1	0
4	L	43	37	37	0	0
5	В	43	36	0	0	0
5	D	43	36	0	0	0
5	F	43	36	0	0	0
5	Н	43	36	0	0	0
6	A	72	0	0	0	0
6	В	125	0	0	0	0
6	С	46	0	0	0	0
6	D	94	0	0	1	0
6	Е	39	0	0	0	0
6	F	133	0	0	1	1
6	G	58	0	0	0	1
6	Н	102	0	0	0	0
6	I	6	0	0	1	0
6	J	5	0	0	0	0
6	K	9	0	0	0	0
6	L	6	0	0	0	0
All	All	10531	9709	9430	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap (Å)} \end{aligned}$
4:B:202:PEB:HNA	4:B:202:PEB:HMB2	1.49	0.77
4:D:202:PEB:HMB2	4:D:202:PEB:HNA	1.50	0.76
4:F:202:PEB:HNA	4:F:202:PEB:HMB2	1.50	0.75
4:H:202:PEB:HMB2	4:H:202:PEB:HNA	1.55	0.72
4:B:203:PEB:HMB2	4:B:203:PEB:NA	2.20	0.56

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
6:F:394:HOH:O	6:G:242:HOH:O[2_557]	2.12	0.08

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percent	iles
1	A	61/67~(91%)	60 (98%)	1 (2%)	0	100 1	.00
1	G	61/67~(91%)	61 (100%)	0	0	100 1	.00
1	J	61/67~(91%)	61 (100%)	0	0	100 1	.00
1	K	61/67~(91%)	61 (100%)	0	0	100 1	.00
2	В	185/177~(104%)	183 (99%)	2 (1%)	0	100 1	.00
2	D	180/177~(102%)	179 (99%)	1 (1%)	0	100 1	.00
2	F	183/177 (103%)	181 (99%)	2 (1%)	0	100 1	.00
2	Н	180/177~(102%)	177 (98%)	3 (2%)	0	100 1	.00
3	C	59/62~(95%)	59 (100%)	0	0	100 1	.00
3	E	59/62~(95%)	59 (100%)	0	0	100 1	.00
3	I	59/62~(95%)	58 (98%)	1 (2%)	0	100 1	.00



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
3	L	59/62~(95%)	59 (100%)	0	0	100	100
All	All	1208/1224 (99%)	1198 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	$45/48 \; (94\%)$	45 (100%)	0	100	100
1	G	45/48~(94%)	45 (100%)	0	100	100
1	J	45/48 (94%)	45 (100%)	0	100	100
1	K	45/48 (94%)	45 (100%)	0	100	100
2	В	150/140 (107%)	149 (99%)	1 (1%)	84	72
2	D	146/140 (104%)	145 (99%)	1 (1%)	84	72
2	F	149/140 (106%)	149 (100%)	0	100	100
2	Н	146/140 (104%)	146 (100%)	0	100	100
3	С	48/48 (100%)	48 (100%)	0	100	100
3	${ m E}$	48/48 (100%)	48 (100%)	0	100	100
3	I	48/48 (100%)	48 (100%)	0	100	100
3	L	48/48 (100%)	48 (100%)	0	100	100
All	All	963/944 (102%)	961 (100%)	2 (0%)	93	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	В	15	LYS
2	D	59	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:



Mol	Chain	Res	Type
1	G	26[A]	GLN
1	G	46[A]	ASN
1	K	46[B]	ASN
3	I	26[B]	GLN
3	Е	26[A]	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain Res		Link	В	ond leng	gths	В	ond ang	gles
IVIOI	Type	Chain	rtes	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
3	LYZ	I	4[B]	3	7,9,10	1.10	0	4,10,12	1.15	0
1	LYZ	G	4[A]	1	7,9,10	0.89	0	4,10,12	0.59	0
3	LYZ	Е	4[A]	3	7,9,10	0.72	0	4,10,12	1.07	0
1	LYZ	A	4[A]	1	7,9,10	1.10	0	4,10,12	1.15	0
1	LYZ	J	4[B]	1	7,9,10	0.72	0	4,10,12	1.26	0
3	LYZ	L	4[B]	3	7,9,10	0.89	0	4,10,12	0.59	0
3	LYZ	С	4[A]	3	7,9,10	0.72	0	4,10,12	1.26	0
1	LYZ	K	4[B]	1	7,9,10	0.72	0	4,10,12	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LYZ	I	4[B]	3	-	0/8/9/11	-
1	LYZ	G	4[A]	1	-	0/8/9/11	-
3	LYZ	Е	4[A]	3	-	0/8/9/11	-
1	LYZ	A	4[A]	1	-	0/8/9/11	-



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LYZ	J	4[B]	1	-	0/8/9/11	-
3	LYZ	L	4[B]	3	-	0/8/9/11	-
3	LYZ	С	4[A]	3	-	0/8/9/11	-
1	LYZ	K	4[B]	1	-	0/8/9/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	AX9	D	201	2	41,46,46	1.54	8 (19%)	41,67,67	1.21	4 (9%)
4	PEB	В	202	2	43,46,46	3.79	26 (60%)	45,67,67	2.23	14 (31%)
5	AX9	В	201	2	41,46,46	1.62	9 (21%)	41,67,67	1.40	6 (14%)
4	PEB	С	101[A]	3	43,46,46	4.29	29 (67%)	45,67,67	2.26	12 (26%)
5	AX9	F	201	2	41,46,46	1.41	8 (19%)	41,67,67	1.39	4 (9%)
4	PEB	F	203	2	43,46,46	3.98	25 (58%)	45,67,67	1.69	10 (22%)
4	PEB	A	101[A]	1	43,46,46	3.71	24 (55%)	45,67,67	2.09	14 (31%)
5	AX9	Н	201	2	41,46,46	1.51	8 (19%)	41,67,67	1.19	4 (9%)



Mol	Type	Chain	Res	Link	В	ond leng	$\operatorname{gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PEB	D	202	2	43,46,46	3.84	28 (65%)	45,67,67	2.12	10 (22%)
4	PEB	F	202	2	43,46,46	3.94	27 (62%)	45,67,67	2.19	13 (28%)
4	PEB	Н	203	2	43,46,46	3.98	26 (60%)	45,67,67	1.75	9 (20%)
4	PEB	K	101[B]	1	43,46,46	4.16	31 (72%)	45,67,67	2.19	9 (20%)
4	PEB	G	101[A]	1	43,46,46	3.76	25 (58%)	45,67,67	2.04	10 (22%)
4	PEB	D	203	2	43,46,46	4.29	33 (76%)	45,67,67	1.73	8 (17%)
4	PEB	В	203	2	43,46,46	3.96	26 (60%)	45,67,67	1.60	9 (20%)
4	PEB	L	101[B]	3	43,46,46	3.76	25 (58%)	45,67,67	2.04	10 (22%)
4	PEB	Н	202	2	43,46,46	4.14	31 (72%)	45,67,67	2.20	11 (24%)
4	PEB	J	101[B]	1	43,46,46	4.29	29 (67%)	45,67,67	2.26	12 (26%)
4	PEB	Е	101[A]	3	43,46,46	4.16	31 (72%)	45,67,67	2.19	9 (20%)
4	PEB	I	101[B]	3	43,46,46	3.71	24 (55%)	45,67,67	2.09	14 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AX9	D	201	2	-	6/26/74/74	0/4/4/4
4	PEB	В	202	2	-	3/24/74/74	0/4/4/4
5	AX9	В	201	2	-	5/26/74/74	0/4/4/4
4	PEB	С	101[A]	3	-	3/24/74/74	0/4/4/4
5	AX9	F	201	2	-	6/26/74/74	0/4/4/4
4	PEB	F	203	2	-	8/24/74/74	0/4/4/4
4	PEB	A	101[A]	1	-	3/24/74/74	0/4/4/4
5	AX9	Н	201	2	-	6/26/74/74	0/4/4/4
4	PEB	D	202	2	-	4/24/74/74	0/4/4/4
4	PEB	F	202	2	-	4/24/74/74	0/4/4/4
4	PEB	Н	203	2	-	6/24/74/74	0/4/4/4
4	PEB	K	101[B]	1	-	4/24/74/74	0/4/4/4
4	PEB	G	101[A]	1	-	2/24/74/74	0/4/4/4
4	PEB	D	203	2	-	8/24/74/74	0/4/4/4
4	PEB	В	203	2	-	6/24/74/74	0/4/4/4
4	PEB	L	101[B]	3		2/24/74/74	0/4/4/4
4	PEB	Н	202	2	-	4/24/74/74	0/4/4/4



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEB	J	101[B]	1	1	3/24/74/74	0/4/4/4
4	PEB	Е	101[A]	3	-	4/24/74/74	0/4/4/4
4	PEB	I	101[B]	3	-	3/24/74/74	0/4/4/4

The worst 5 of 473 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	Observed(A)	$\operatorname{Ideal}( ext{\AA})$
4	Н	202	PEB	CHB-C4B	10.72	1.44	1.35
4	G	101[A]	PEB	CHB-C4B	10.66	1.44	1.35
4	L	101[B]	PEB	CHB-C4B	10.66	1.44	1.35
4	С	101[A]	PEB	CHB-C4B	10.51	1.43	1.35
4	J	101[B]	PEB	CHB-C4B	10.51	1.43	1.35

The worst 5 of 192 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
4	Н	202	PEB	C1C-CHB-C4B	-9.46	117.51	128.81
4	D	202	PEB	C1C-CHB-C4B	-8.89	118.19	128.81
4	Е	101[A]	PEB	C1C-CHB-C4B	-8.21	119.01	128.81
4	K	101[B]	PEB	C1C-CHB-C4B	-8.21	119.01	128.81
4	В	202	PEB	C1C-CHB-C4B	-8.06	119.18	128.81

There are no chirality outliers.

5 of 90 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	101[A]	PEB	NB-C1B-CHA-C4A
4	A	101[A]	PEB	C2B-C1B-CHA-C4A
4	В	202	PEB	NB-C1B-CHA-C4A
4	В	202	PEB	C2B-C1B-CHA-C4A
4	В	203	PEB	NA-C4A-CHA-C1B

There are no ring outliers.

13 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	202	PEB	1	0
4	С	101[A]	PEB	1	0
4	F	203	PEB	3	0
4	A	101[A]	PEB	2	0




Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	202	PEB	3	0
4	F	202	PEB	2	0
4	K	101[B]	PEB	1	0
4	D	203	PEB	5	0
4	В	203	PEB	3	0
4	Н	202	PEB	1	0
4	J	101[B]	PEB	1	0
4	Е	101[A]	PEB	1	0
4	I	101[B]	PEB	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



































































































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2		$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	63/67~(94%)	0.02	2 (3%) 47	49	12, 18, 32, 41	63 (100%)
1	G	63/67 (94%)	-0.02	2 (3%) 47	49	14, 22, 34, 39	63 (100%)
1	J	63/67 (94%)	0.13	2 (3%) 47	49	16, 23, 36, 43	63 (100%)
1	K	63/67 (94%)	0.11	2 (3%) 47	49	15, 22, 36, 40	63 (100%)
2	В	177/177 (100%)	-0.06	1 (0%) 89	90	13, 20, 33, 46	0
2	D	173/177 (97%)	0.23	8 (4%) 32	32	13, 24, 47, 62	0
2	F	173/177 (97%)	0.02	2 (1%) 79	80	12, 19, 38, 75	0
2	Н	173/177 (97%)	0.05	4 (2%) 60	62	13, 21, 36, 49	0
3	С	61/62 (98%)	0.06	2 (3%) 46	47	16, 23, 36, 39	61 (100%)
3	E	$61/62 \; (98\%)$	0.01	2 (3%) 46	47	15, 22, 36, 40	61 (100%)
3	I	61/62 (98%)	-0.00	1 (1%) 72	74	12, 18, 31, 39	61 (100%)
3	L	61/62 (98%)	-0.09	2 (3%) 46	47	14, 21, 35, 39	61 (100%)
All	All	1192/1224 (97%)	0.05	30 (2%) 57	58	12, 21, 38, 75	496 (41%)

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ	
1	J	62[B]	PHE	6.0	
2	D	30	PHE	5.5	
3	I	62[B]	LEU	5.1	
1	K	62[B]	PHE	4.8	
2	F	14	GLY	4.6	

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
1	LYZ	G	4[A]	10/11	0.94	0.09	24,31,44,48	21
3	LYZ	L	4[B]	10/11	0.94	0.09	24,31,44,48	21
3	LYZ	I	4[B]	10/11	0.95	0.08	18,22,34,38	21
1	LYZ	A	4[A]	10/11	0.95	0.08	18,22,34,38	21
3	LYZ	Ε	4[A]	10/11	0.96	0.07	18,25,39,47	21
1	LYZ	J	4[B]	10/11	0.96	0.09	22,27,43,47	21
1	LYZ	K	4[B]	10/11	0.96	0.07	18,25,39,47	21
3	LYZ	С	4[A]	10/11	0.96	0.09	22,27,43,47	21

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q<0.9
4	PEB	Е	101[A]	43/43	0.94	0.10	16,25,42,51	80
4	PEB	K	101[B]	43/43	0.94	0.10	16,25,42,51	80
5	AX9	D	201	43/43	0.94	0.09	17,24,31,36	0
5	AX9	Н	201	43/43	0.94	0.09	14,23,30,35	0
4	PEB	J	101[B]	43/43	0.95	0.10	15,23,33,57	80
4	PEB	D	203	43/43	0.95	0.10	18,25,32,38	0
4	PEB	С	101[A]	43/43	0.95	0.10	15,23,33,57	80
5	AX9	F	201	43/43	0.95	0.09	12,17,25,29	0
4	PEB	Н	202	43/43	0.95	0.10	16,24,31,40	0
4	PEB	Н	203	43/43	0.96	0.09	13,19,24,31	0
4	PEB	I	101[B]	43/43	0.96	0.11	10,16,31,64	80
4	PEB	В	202	43/43	0.96	0.10	13,18,27,32	0
4	PEB	В	203	43/43	0.96	0.10	13,20,27,28	0
4	PEB	L	101[B]	43/43	0.96	0.11	13,18,33,48	80
5	AX9	В	201	43/43	0.96	0.08	15,22,33,40	0
4	PEB	F	203	43/43	0.96	0.09	12,19,33,50	0
4	PEB	G	101[A]	43/43	0.96	0.11	13,18,33,48	80
4	PEB	A	101[A]	43/43	0.96	0.11	10,16,31,64	80


Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B-factors}({f \AA}^2)$	Q<0.9
4	PEB	D	202	43/43	0.97	0.10	13,18,24,28	0
4	PEB	F	202	43/43	0.97	0.10	10,18,27,30	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











































## Electron density around PEB H 203:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










### Electron density around PEB I 101 (B):

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











































## 6.5 Other polymers (i)

There are no such residues in this entry.

