



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 21, 2023 – 10:50 AM EDT

PDB ID : 8EN5
Title : Structure of GII.4 norovirus in complex with Nanobody 56
Authors : Kher, G.; Sabin, C.; Pancera, M.; Koromyslova, A.; Hansman, G.
Deposited on : 2022-09-28
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	91% 6% .
1	B	317	2% 91% 6% .
1	C	317	2% 95% . .
1	D	317	4% 87% 9% . .
2	E	131	2% 89% 6% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	131	<p>2% 86% 8% 5%</p>
2	G	131	<p>5% 84% 15%</p>
2	H	131	<p>2% 82% 12% 5%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	606	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15074 atoms, of which 36 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GII.4 P domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	308	2418	1532	415	461	10	0	2	0
1	B	308	2427	1540	415	462	10	0	3	0
1	C	308	2406	1524	412	460	10	0	0	0
1	D	306	2395	1518	410	457	10	0	0	0

- Molecule 2 is a protein called Nanobody 56.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	126	973	611	170	188	4	0	0	0
2	F	125	963	605	167	187	4	0	0	0
2	G	131	1026	643	185	194	4	0	1	0
2	H	124	957	602	166	185	4	0	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C H O 10 2 6 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C H O 10 2 6 2	0	0
3	B	1	Total C H O 10 2 6 2	0	0
3	B	1	Total C H O 10 2 6 2	0	0
3	E	1	Total C H O 10 2 6 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C H O 10 2 6 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	G	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	H	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	322	Total	O	0	0
			322	322		
5	B	278	Total	O	0	0
			278	278		
5	E	83	Total	O	0	0
			83	83		
5	F	95	Total	O	0	0
			95	95		
5	C	238	Total	O	0	0
			238	238		
5	D	166	Total	O	0	0
			166	166		
5	G	69	Total	O	0	0
			69	69		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	66	Total 66	O 66	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GII.4 P domain




- Molecule 1: GII.4 P domain




- Molecule 1: GII.4 P domain




- Molecule 1: GII.4 P domain





- Molecule 2: Nanobody 56




- Molecule 2: Nanobody 56



- Molecule 2: Nanobody 56



- Molecule 2: Nanobody 56



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.24Å 235.38Å 70.57Å 90.00° 104.30° 90.00°	Depositor
Resolution (Å)	68.38 – 1.60 68.38 – 1.60	Depositor EDS
% Data completeness (in resolution range)	95.4 (68.38-1.60) 92.8 (68.38-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.186 , 0.219 0.185 , 0.218	Depositor DCC
R_{free} test set	12091 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	20.1	Xtrriage
Anisotropy	0.251	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15074	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/2494	0.82	3/3411 (0.1%)
1	B	0.76	0/2507	0.81	1/3429 (0.0%)
1	C	0.63	0/2476	0.74	0/3387
1	D	0.55	0/2465	0.69	0/3372
2	E	0.62	0/994	0.81	0/1348
2	F	0.65	0/983	0.76	0/1333
2	G	0.54	0/1055	0.71	0/1431
2	H	0.60	1/977 (0.1%)	0.71	0/1325
All	All	0.66	1/13951 (0.0%)	0.76	4/19036 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	50	CYS	CB-SG	-6.85	1.70	1.82

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	278	LEU	C-N-CA	-5.71	107.42	121.70
1	A	447	MET	CG-SD-CE	5.64	109.22	100.20
1	A	448	ASP	CB-CG-OD2	-5.58	113.28	118.30
1	B	310	ASP	CB-CG-OD1	5.00	122.81	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	410	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2418	0	2322	14	0
1	B	2427	0	2331	12	0
1	C	2406	0	2302	9	0
1	D	2395	0	2292	28	0
2	E	973	0	944	9	0
2	F	963	0	937	21	0
2	G	1026	0	984	24	0
2	H	957	0	932	22	0
3	A	32	6	48	6	0
3	B	52	18	78	8	0
3	C	20	6	30	3	0
3	D	16	0	24	3	0
3	E	4	6	6	0	0
3	F	12	0	18	2	0
3	G	4	0	6	1	0
3	H	4	0	6	1	0
4	A	12	0	12	0	0
5	A	322	0	0	3	0
5	B	278	0	0	3	0
5	C	238	0	0	4	0
5	D	166	0	0	1	0
5	E	83	0	0	0	0
5	F	95	0	0	2	0
5	G	69	0	0	0	0
5	H	66	0	0	0	0
All	All	15038	36	13272	133	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 133 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:31:ARG:HH11	2:F:72:ARG:HD2	1.11	1.15
2:H:72:ARG:HD3	2:H:74:ASN:HD21	1.35	0.91
2:G:125:SER:O	2:G:129:HIS:NE2	2.03	0.90
1:C:501:HIS:CE1	1:C:531:GLY:HA3	2.09	0.86
2:F:31:ARG:NH1	2:F:72:ARG:HD2	1.92	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/317 (97%)	303 (98%)	5 (2%)	0	100	100
1	B	309/317 (98%)	301 (97%)	8 (3%)	0	100	100
1	C	306/317 (96%)	300 (98%)	6 (2%)	0	100	100
1	D	304/317 (96%)	299 (98%)	4 (1%)	1 (0%)	41	21
2	E	124/131 (95%)	116 (94%)	7 (6%)	1 (1%)	19	6
2	F	123/131 (94%)	119 (97%)	3 (2%)	1 (1%)	19	6
2	G	130/131 (99%)	124 (95%)	5 (4%)	1 (1%)	19	6
2	H	122/131 (93%)	118 (97%)	4 (3%)	0	100	100
All	All	1726/1792 (96%)	1680 (97%)	42 (2%)	4 (0%)	47	26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	29	LEU
2	F	29	LEU
2	G	29	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	442	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/273 (98%)	269 (100%)	0	100	100
1	B	270/273 (99%)	268 (99%)	2 (1%)	84	73
1	C	267/273 (98%)	267 (100%)	0	100	100
1	D	266/273 (97%)	263 (99%)	3 (1%)	73	57
2	E	105/110 (96%)	104 (99%)	1 (1%)	76	61
2	F	104/110 (94%)	102 (98%)	2 (2%)	57	34
2	G	111/110 (101%)	110 (99%)	1 (1%)	78	65
2	H	103/110 (94%)	102 (99%)	1 (1%)	76	61
All	All	1495/1532 (98%)	1485 (99%)	10 (1%)	84	73

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	373	ARG
2	G	1	GLN
2	H	76	LYS
2	F	19	ARG
2	F	74	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	3	GLN
2	H	5	GLN
2	H	74	ASN
1	C	469	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	469	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

37 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	612	-	3,3,3	0.41	0	2,2,2	0.75	0
3	EDO	B	608	-	3,3,3	0.41	0	2,2,2	0.57	0
3	EDO	E	201	-	3,3,3	0.40	0	2,2,2	0.46	0
3	EDO	A	602	-	3,3,3	0.66	0	2,2,2	0.31	0
3	EDO	B	609	-	3,3,3	0.60	0	2,2,2	0.26	0
3	EDO	C	602	-	3,3,3	0.52	0	2,2,2	0.58	0
3	EDO	B	611	-	3,3,3	0.90	0	2,2,2	0.58	0
3	EDO	F	203	-	3,3,3	0.59	0	2,2,2	0.06	0
3	EDO	D	602	-	3,3,3	0.58	0	2,2,2	0.18	0
3	EDO	H	201	-	3,3,3	0.45	0	2,2,2	0.36	0
3	EDO	B	606	-	3,3,3	0.58	0	2,2,2	0.94	0
3	EDO	F	202	-	3,3,3	0.47	0	2,2,2	0.69	0
3	EDO	B	607	-	3,3,3	0.71	0	2,2,2	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	D	604	-	3,3,3	0.57	0	2,2,2	0.32	0
3	EDO	C	604	-	3,3,3	0.41	0	2,2,2	0.29	0
3	EDO	C	605	-	3,3,3	0.46	0	2,2,2	0.47	0
3	EDO	G	201	-	3,3,3	0.46	0	2,2,2	0.37	0
3	EDO	A	606	-	3,3,3	0.44	0	2,2,2	0.51	0
3	EDO	C	601	-	3,3,3	0.58	0	2,2,2	0.26	0
3	EDO	D	601	-	3,3,3	0.59	0	2,2,2	0.21	0
3	EDO	B	610	-	3,3,3	0.47	0	2,2,2	0.25	0
3	EDO	D	603	-	3,3,3	0.57	0	2,2,2	0.31	0
3	EDO	B	603	-	3,3,3	0.55	0	2,2,2	0.55	0
3	EDO	C	603	-	3,3,3	0.68	0	2,2,2	0.07	0
3	EDO	F	201	-	3,3,3	0.44	0	2,2,2	0.70	0
4	MES	A	608	-	12,12,12	1.89	1 (8%)	14,16,16	2.27	6 (42%)
3	EDO	B	602	-	3,3,3	0.50	0	2,2,2	0.47	0
3	EDO	B	604	-	3,3,3	0.46	0	2,2,2	0.51	0
3	EDO	A	609	-	3,3,3	0.48	0	2,2,2	1.22	0
3	EDO	A	604	-	3,3,3	0.55	0	2,2,2	0.31	0
3	EDO	B	605	-	3,3,3	0.63	0	2,2,2	0.91	0
3	EDO	A	607	-	3,3,3	0.54	0	2,2,2	0.34	0
3	EDO	B	601	-	3,3,3	0.63	0	2,2,2	0.50	0
3	EDO	B	613	-	3,3,3	0.43	0	2,2,2	1.51	0
3	EDO	A	605	-	3,3,3	0.74	0	2,2,2	0.30	0
3	EDO	A	601	-	3,3,3	0.50	0	2,2,2	0.45	0
3	EDO	A	603	-	3,3,3	0.37	0	2,2,2	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	612	-	-	1/1/1/1	-
3	EDO	B	608	-	-	1/1/1/1	-
3	EDO	E	201	-	-	0/1/1/1	-
3	EDO	A	602	-	-	0/1/1/1	-
3	EDO	B	609	-	-	0/1/1/1	-
3	EDO	C	602	-	-	0/1/1/1	-
3	EDO	B	611	-	-	0/1/1/1	-
3	EDO	F	203	-	-	0/1/1/1	-
3	EDO	D	602	-	-	1/1/1/1	-
3	EDO	H	201	-	-	1/1/1/1	-
3	EDO	B	606	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	F	202	-	-	1/1/1/1	-
3	EDO	B	607	-	-	1/1/1/1	-
3	EDO	D	604	-	-	1/1/1/1	-
3	EDO	C	604	-	-	0/1/1/1	-
3	EDO	C	605	-	-	0/1/1/1	-
3	EDO	G	201	-	-	1/1/1/1	-
3	EDO	A	606	-	-	0/1/1/1	-
3	EDO	C	601	-	-	0/1/1/1	-
3	EDO	D	601	-	-	0/1/1/1	-
3	EDO	B	610	-	-	1/1/1/1	-
3	EDO	D	603	-	-	0/1/1/1	-
3	EDO	B	603	-	-	0/1/1/1	-
3	EDO	C	603	-	-	0/1/1/1	-
3	EDO	F	201	-	-	0/1/1/1	-
4	MES	A	608	-	-	5/6/14/14	0/1/1/1
3	EDO	B	602	-	-	0/1/1/1	-
3	EDO	B	604	-	-	0/1/1/1	-
3	EDO	A	609	-	-	1/1/1/1	-
3	EDO	A	604	-	-	1/1/1/1	-
3	EDO	B	605	-	-	0/1/1/1	-
3	EDO	A	607	-	-	0/1/1/1	-
3	EDO	B	601	-	-	1/1/1/1	-
3	EDO	B	613	-	-	1/1/1/1	-
3	EDO	A	605	-	-	1/1/1/1	-
3	EDO	A	601	-	-	0/1/1/1	-
3	EDO	A	603	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	608	MES	C8-S	-6.01	1.69	1.77

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	608	MES	C5-N4-C3	5.10	120.31	108.83
4	A	608	MES	O3S-S-C8	3.74	111.82	105.77
4	A	608	MES	C7-N4-C5	2.73	118.22	111.23
4	A	608	MES	C7-N4-C3	2.72	118.20	111.23
4	A	608	MES	O2S-S-C8	-2.19	104.28	106.92

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	608	MES	C8-C7-N4-C5
4	A	608	MES	C7-C8-S-O1S
4	A	608	MES	C7-C8-S-O2S
4	A	608	MES	C7-C8-S-O3S
3	B	606	EDO	O1-C1-C2-O2

There are no ring outliers.

14 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	609	EDO	1	0
3	D	602	EDO	3	0
3	H	201	EDO	1	0
3	B	606	EDO	4	0
3	F	202	EDO	1	0
3	C	604	EDO	2	0
3	G	201	EDO	1	0
3	C	601	EDO	1	0
3	B	610	EDO	2	0
3	F	201	EDO	1	0
3	A	609	EDO	2	0
3	A	604	EDO	2	0
3	B	613	EDO	1	0
3	A	603	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/317 (97%)	-0.25	0 100 100	13, 19, 31, 41	0
1	B	308/317 (97%)	-0.16	7 (2%) 60 59	13, 20, 34, 67	0
1	C	308/317 (97%)	-0.15	7 (2%) 60 59	18, 26, 43, 58	0
1	D	306/317 (96%)	0.20	14 (4%) 32 29	23, 34, 50, 66	0
2	E	126/131 (96%)	0.05	3 (2%) 59 56	17, 27, 43, 66	0
2	F	125/131 (95%)	-0.13	3 (2%) 59 56	17, 27, 41, 51	0
2	G	131/131 (100%)	0.18	6 (4%) 32 29	22, 34, 56, 69	0
2	H	124/131 (94%)	0.12	3 (2%) 59 56	24, 34, 47, 57	0
All	All	1736/1792 (96%)	-0.05	43 (2%) 57 55	13, 27, 46, 69	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	42	GLY	6.6
1	D	412	ASN	6.1
2	G	126	HIS	5.8
2	E	125	SER	5.2
1	B	411	ASN	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	B	606	4/4	0.73	0.21	30,30,36,36	0
3	EDO	C	602	4/4	0.77	0.17	36,38,41,45	0
3	EDO	F	203	4/4	0.79	0.27	35,38,48,56	0
3	EDO	F	202	4/4	0.81	0.32	37,42,43,44	0
3	EDO	B	613	4/4	0.84	0.25	35,42,43,43	0
3	EDO	F	201	4/4	0.86	0.15	33,35,40,42	0
3	EDO	H	201	4/4	0.86	0.15	41,42,51,54	0
3	EDO	B	607	4/4	0.87	0.13	27,31,32,34	0
3	EDO	B	609	4/4	0.87	0.21	33,36,37,42	0
3	EDO	B	601	4/4	0.89	0.16	30,31,36,42	0
3	EDO	A	603	4/4	0.90	0.12	31,32,35,44	0
3	EDO	D	603	4/4	0.90	0.13	31,32,39,40	0
3	EDO	A	604	4/4	0.90	0.10	28,31,32,33	0
3	EDO	B	611	4/4	0.91	0.13	22,32,39,41	0
3	EDO	C	604	4/4	0.91	0.13	40,43,47,52	0
3	EDO	D	602	4/4	0.91	0.13	35,36,41,41	0
3	EDO	B	612	4/4	0.91	0.16	30,36,41,43	0
3	EDO	B	608	4/4	0.91	0.22	21,28,30,38	0
3	EDO	A	609	4/4	0.92	0.18	20,31,38,38	0
3	EDO	A	607	4/4	0.92	0.15	27,31,31,45	0
3	EDO	D	604	4/4	0.92	0.15	28,30,35,41	0
3	EDO	C	605	4/4	0.92	0.10	33,39,47,50	0
3	EDO	A	606	4/4	0.93	0.07	27,31,32,36	0
3	EDO	D	601	4/4	0.93	0.10	27,28,31,36	0
3	EDO	A	605	4/4	0.93	0.12	20,24,29,36	0
3	EDO	E	201	4/4	0.94	0.20	32,39,46,47	0
3	EDO	C	603	4/4	0.94	0.12	31,32,33,35	0
3	EDO	G	201	4/4	0.94	0.28	35,35,39,40	0
3	EDO	B	610	4/4	0.94	0.18	27,32,35,35	0
4	MES	A	608	12/12	0.94	0.20	28,44,52,53	0
3	EDO	C	601	4/4	0.95	0.12	26,28,31,34	0
3	EDO	A	602	4/4	0.95	0.09	21,23,29,31	0
3	EDO	B	602	4/4	0.95	0.09	23,26,29,30	0
3	EDO	B	604	4/4	0.95	0.12	20,22,30,37	0
3	EDO	B	603	4/4	0.97	0.07	21,26,27,30	0
3	EDO	B	605	4/4	0.97	0.08	21,24,28,32	0
3	EDO	A	601	4/4	0.98	0.06	21,21,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.