



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2021 – 03:43 PM EDT

PDB ID : 3EQV
Title : Crystal structure of penicillin-binding protein 2 from *Neisseria gonorrhoeae* containing four mutations associated with penicillin resistance
Authors : Powell, A.J.; Deacon, A.M.; Nicholas, R.A.; Davies, C.
Deposited on : 2008-10-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	 4% 69% 11% 18%
1	B	542	 6% 68% 10% 21%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	B	12	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6805 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Penicillin-binding protein 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	443	3395	2126	613	647	1	8	0	1	0
1	B	426	3244	2034	583	618	1	8	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	GLY	-	expression tag	UNP P08149
A	41	SER	-	expression tag	UNP P08149
A	42	GLY	-	expression tag	UNP P08149
A	43	GLY	-	expression tag	UNP P08149
A	504	LEU	PHE	engineered mutation	UNP P08149
A	510	VAL	ALA	engineered mutation	UNP P08149
A	516	GLY	ALA	engineered mutation	UNP P08149
A	551	SER	PRO	engineered mutation	UNP P08149
B	40	GLY	-	expression tag	UNP P08149
B	41	SER	-	expression tag	UNP P08149
B	42	GLY	-	expression tag	UNP P08149
B	43	GLY	-	expression tag	UNP P08149
B	504	LEU	PHE	engineered mutation	UNP P08149
B	510	VAL	ALA	engineered mutation	UNP P08149
B	516	GLY	ALA	engineered mutation	UNP P08149
B	551	SER	PRO	engineered mutation	UNP P08149

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	40	Total	O	0	0
			40	40		
4	B	39	Total	O	0	0
			39	39		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Penicillin-binding protein 2



- Molecule 1: Penicillin-binding protein 2





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.60Å 137.20Å 229.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.10 – 2.40 48.21 – 2.39	Depositor EDS
% Data completeness (in resolution range)	97.7 (48.10-2.40) 97.6 (48.21-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.217 , 0.253 0.260 , 0.293	Depositor DCC
R_{free} test set	3558 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	43.1	Xtrriage
Anisotropy	0.797	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6805	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	1/3452 (0.0%)	0.71	0/4671
1	B	0.52	0/3293	0.65	1/4457 (0.0%)
All	All	0.55	1/6745 (0.0%)	0.68	1/9128 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	333	ASN	CG-ND2	9.17	1.55	1.32

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	332	LEU	CA-CB-CG	6.79	130.91	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3395	0	3392	35	0
1	B	3244	0	3252	26	0
2	A	30	0	0	0	0
2	B	45	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	40	0	0	0	0
4	B	39	0	0	0	0
All	All	6805	0	6660	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:THR:HB	1:B:93:GLU:HA	1.49	0.94
1:A:251:LEU:HG	1:A:265:VAL:HG23	1.49	0.94
1:A:86:LEU:CD2	1:A:171:MET:HE3	2.18	0.74
1:A:86:LEU:CD2	1:A:171:MET:CE	2.65	0.74
1:A:522:ALA:HB3	1:A:557:MET:HE2	1.69	0.73
1:A:522:ALA:HB3	1:A:557:MET:CE	2.19	0.73
1:B:513:LYS:HB3	1:B:537:GLU:HG2	1.73	0.68
1:A:259:GLN:HB3	1:A:539:THR:HB	1.80	0.64
1:A:171:MET:SD	1:A:241:GLN:NE2	2.72	0.63
1:B:468:ARG:HH11	1:B:468:ARG:HG3	1.64	0.62
1:B:318:ALA:HB2	1:B:470:VAL:HG21	1.83	0.59
1:A:380:TYR:CE2	1:A:404:LEU:HD13	2.37	0.59
1:A:470:VAL:HG12	1:A:474:MET:HE2	1.84	0.59
1:A:281:THR:HG22	1:A:282:PRO:HA	1.85	0.58
1:A:86:LEU:HD23	1:A:171:MET:HE3	1.85	0.58
1:B:183:ASP:HB3	1:B:189:GLN:HE21	1.69	0.57
1:A:251:LEU:HG	1:A:265:VAL:CG2	2.30	0.57
1:A:53:THR:HB	1:A:58:LYS:HE3	1.87	0.56
1:A:86:LEU:CD2	1:A:171:MET:HE1	2.37	0.55
1:B:250:GLU:OE2	1:B:555:LYS:HD3	2.09	0.53
1:B:471:ARG:HG2	1:B:521:PHE:CE2	2.44	0.53
1:A:305:MET:HG2	1:A:402:GLY:HA2	1.90	0.53
1:A:293:ASP:HB2	1:A:296:GLN:OE1	2.11	0.51
1:A:356:ARG:NH1	1:A:476:SER:OG	2.44	0.50
1:A:471:ARG:HG2	1:A:521:PHE:CE2	2.46	0.49
1:B:246:LEU:HD11	1:B:559:GLY:HA3	1.94	0.49
1:A:557:MET:HE3	1:A:561:LEU:HD13	1.93	0.49
1:B:67:ARG:NH1	1:B:67:ARG:HB2	2.28	0.49
1:A:173:ASN:C	1:A:173:ASN:HD22	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:LEU:HD21	1:A:171:MET:HE1	1.95	0.48
1:A:167:ARG:HD2	1:A:193:GLU:OE2	2.14	0.47
1:B:470:VAL:HG12	1:B:474:MET:HE2	1.96	0.47
1:A:557:MET:HE2	1:A:561:LEU:HD21	1.97	0.47
1:A:428:LEU:HD13	1:A:534:THR:HG23	1.97	0.46
1:B:171:MET:SD	1:B:241:GLN:HG3	2.55	0.46
1:B:380:TYR:HB2	1:B:407:TRP:HB3	1.97	0.46
1:B:471:ARG:HD3	1:B:527:PRO:CG	2.45	0.46
1:A:251:LEU:HD13	1:A:281:THR:OG1	2.16	0.46
1:A:546:GLY:O	1:A:550:GLY:HA3	2.16	0.45
1:B:498:THR:HB	1:B:518:PHE:CD2	2.51	0.45
1:B:362:SER:OG	1:B:497:LYS:NZ	2.46	0.45
1:A:335:GLN:HG3	1:A:336:PRO:HD2	1.98	0.45
1:A:62:ASP:O	1:A:213:ARG:HB3	2.17	0.45
1:B:428:LEU:HD13	1:B:534:THR:HG23	1.99	0.44
1:B:350:TYR:CE1	1:B:361:LYS:HD3	2.53	0.44
1:B:438:LEU:HD13	1:B:474:MET:CE	2.48	0.44
1:A:428:LEU:HD13	1:A:534:THR:CG2	2.48	0.44
1:B:487:GLY:HA3	1:B:518:PHE:CZ	2.53	0.43
1:A:557:MET:HE2	1:A:561:LEU:CD2	2.48	0.43
1:B:318:ALA:CB	1:B:470:VAL:HG21	2.48	0.43
1:B:513:LYS:HD3	1:B:537:GLU:OE1	2.19	0.43
1:B:374:PHE:HB3	1:B:378:GLU:HG3	2.01	0.42
1:A:89:SER:HA	1:A:166:LYS:O	2.20	0.41
1:B:211:ARG:HG2	1:B:217:ILE:HA	2.02	0.41
1:A:86:LEU:HD22	1:A:171:MET:HE3	1.99	0.41
1:B:173:ASN:HD22	1:B:174:LEU:N	2.18	0.41
1:A:173:ASN:HD22	1:A:174:LEU:N	2.18	0.41
1:A:281:THR:CG2	1:A:282:PRO:HA	2.51	0.41
1:B:92:THR:HB	1:B:93:GLU:CA	2.36	0.41
1:A:522:ALA:HB3	1:A:557:MET:HE1	1.98	0.40
1:B:547:VAL:O	1:B:547:VAL:HG12	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	437/542 (81%)	429 (98%)	7 (2%)	1 (0%)	47	62
1	B	417/542 (77%)	406 (97%)	11 (3%)	0	100	100
All	All	854/1084 (79%)	835 (98%)	18 (2%)	1 (0%)	51	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	480	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	356/438 (81%)	323 (91%)	33 (9%)	9	13
1	B	340/438 (78%)	317 (93%)	23 (7%)	16	25
All	All	696/876 (80%)	640 (92%)	56 (8%)	11	18

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	THR
1	A	80	ASP
1	A	92	THR
1	A	167	ARG
1	A	173	ASN
1	A	182	THR
1	A	185	ASP
1	A	194	LEU
1	A	203	GLU
1	A	218	VAL
1	A	219	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	251	LEU
1	A	256	GLU
1	A	259	GLN
1	A	281	THR
1	A	293	ASP
1	A	321	LEU
1	A	332	LEU
1	A	338	LYS
1	A	345	ARG
1	A	362	SER
1	A	373	ARG
1	A	404	LEU
1	A	424	LEU
1	A	426	LEU
1	A	431	LEU
1	A	460	ARG
1	A	472	ASN
1	A	473	LEU
1	A	512	ASN
1	A	547	VAL
1	A	551	SER
1	A	561	LEU
1	B	80	ASP
1	B	93	GLU
1	B	173	ASN
1	B	194	LEU
1	B	237	LEU
1	B	241	GLN
1	B	246	LEU
1	B	256	GLU
1	B	271	ARG
1	B	298	ARG
1	B	302	VAL
1	B	306	ILE
1	B	331	ARG
1	B	332	LEU
1	B	345	ARG
1	B	349	VAL
1	B	362	SER
1	B	371	SER
1	B	390	VAL
1	B	403	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	424	LEU
1	B	473	LEU
1	B	562	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	173	ASN
1	A	329	ASN
1	A	360	GLN
1	A	425	GLN
1	B	69	GLN
1	B	173	ASN
1	B	189	GLN
1	B	562	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	B	465	1	8,9,10	1.68	1 (12%)	8,12,14	2.80	3 (37%)
1	SEP	A	465	1	8,9,10	1.59	1 (12%)	8,12,14	1.85	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	465	1	-	3/5/8/10	-
1	SEP	A	465	1	-	2/5/8/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	465	SEP	P-O1P	3.34	1.61	1.50
1	B	465	SEP	P-O1P	3.28	1.61	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	465	SEP	OG-CB-CA	6.94	114.90	108.14
1	A	465	SEP	OG-CB-CA	4.15	112.18	108.14
1	B	465	SEP	O2P-P-OG	2.27	112.78	106.73
1	A	465	SEP	OG-P-O1P	2.25	112.78	106.47
1	B	465	SEP	O3P-P-OG	2.09	112.30	106.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	465	SEP	N-CA-CB-OG
1	B	465	SEP	N-CA-CB-OG
1	B	465	SEP	CB-OG-P-O1P
1	A	465	SEP	CA-CB-OG-P
1	B	465	SEP	CA-CB-OG-P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	15	-	4,4,4	0.15	0	6,6,6	0.18	0
2	SO4	B	2	-	4,4,4	0.11	0	6,6,6	0.24	0
2	SO4	B	9	-	4,4,4	0.12	0	6,6,6	0.87	0
2	SO4	B	10	-	4,4,4	0.15	0	6,6,6	0.08	0
2	SO4	B	12	-	4,4,4	0.15	0	6,6,6	0.27	0
2	SO4	A	13	-	4,4,4	0.14	0	6,6,6	0.14	0
2	SO4	B	6	-	4,4,4	0.12	0	6,6,6	0.16	0
3	GOL	A	2	-	5,5,5	0.41	0	5,5,5	0.39	0
2	SO4	A	11	-	4,4,4	0.13	0	6,6,6	0.32	0
2	SO4	A	8	-	4,4,4	0.15	0	6,6,6	0.07	0
2	SO4	A	4	-	4,4,4	0.12	0	6,6,6	0.41	0
2	SO4	B	7	-	4,4,4	0.16	0	6,6,6	0.11	0
2	SO4	B	5	-	4,4,4	0.16	0	6,6,6	0.35	0
3	GOL	B	582	-	5,5,5	0.42	0	5,5,5	0.33	0
2	SO4	B	1	-	4,4,4	0.17	0	6,6,6	0.18	0
2	SO4	A	3	-	4,4,4	0.14	0	6,6,6	0.15	0
2	SO4	B	14	-	4,4,4	0.17	0	6,6,6	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	2	-	-	2/4/4/4	-
3	GOL	B	582	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2	GOL	C1-C2-C3-O3
3	B	582	GOL	O1-C1-C2-O2
3	B	582	GOL	O1-C1-C2-C3
3	B	582	GOL	C1-C2-C3-O3
3	A	2	GOL	O2-C2-C3-O3
3	B	582	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/542 (81%)	0.38	21 (4%) 30 29	33, 46, 56, 65	0
1	B	425/542 (78%)	0.65	31 (7%) 15 13	33, 49, 61, 71	0
All	All	867/1084 (79%)	0.52	52 (5%) 21 20	33, 48, 60, 71	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	218	VAL	6.1
1	B	217	ILE	5.9
1	A	543	TYR	5.1
1	B	209	VAL	4.5
1	B	211	ARG	4.4
1	B	216	ASN	4.4
1	B	220	SER	4.3
1	A	572	LEU	4.2
1	B	454	VAL	4.1
1	B	66	VAL	4.0
1	A	573	THR	4.0
1	A	214	GLN	3.8
1	B	67	ARG	3.6
1	A	92	THR	3.5
1	A	328	LEU	3.5
1	A	62	ASP	3.4
1	B	184	ILE	3.3
1	A	348	HIS	3.3
1	B	212	ASP	3.2
1	B	219	ASP	3.1
1	A	571	PRO	3.1
1	B	185	ASP	3.1
1	B	572	LEU	3.0
1	B	571	PRO	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	164	GLU	2.7
1	A	512	ASN	2.7
1	A	213	ARG	2.7
1	B	91	PRO	2.7
1	B	453	ALA	2.7
1	A	547	VAL	2.6
1	A	165	LEU	2.5
1	B	490	ASP	2.5
1	A	540	ALA	2.5
1	B	68	THR	2.4
1	B	69	GLN	2.4
1	A	539	THR	2.4
1	B	92	THR	2.3
1	A	61	GLY	2.3
1	B	93	GLU	2.3
1	B	421	GLY	2.1
1	B	207	GLU	2.1
1	B	417	THR	2.1
1	B	424	LEU	2.1
1	A	329	ASN	2.1
1	A	351	PRO	2.1
1	B	271	ARG	2.1
1	B	309	GLY	2.1
1	B	206	ALA	2.1
1	A	513	LYS	2.1
1	B	516	GLY	2.0
1	B	165	LEU	2.0
1	A	525	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	B	465	10/11	0.76	0.22	52,53,57,58	5
1	SEP	A	465	10/11	0.86	0.18	46,47,50,50	5

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	12	5/5	0.67	0.43	59,59,60,61	5
2	SO4	B	6	5/5	0.70	0.22	97,97,98,98	5
2	SO4	B	10	5/5	0.76	0.15	73,73,73,73	5
2	SO4	A	15	5/5	0.83	0.20	59,60,60,61	5
2	SO4	B	7	5/5	0.86	0.26	62,62,63,63	5
2	SO4	A	11	5/5	0.88	0.19	61,64,64,65	5
2	SO4	A	3	5/5	0.88	0.18	64,65,66,66	5
2	SO4	B	14	5/5	0.88	0.27	42,42,42,43	5
2	SO4	A	4	5/5	0.89	0.16	74,75,75,76	5
2	SO4	A	13	5/5	0.90	0.12	65,65,65,66	5
3	GOL	A	2	6/6	0.91	0.18	53,54,55,56	0
3	GOL	B	582	6/6	0.92	0.17	62,63,64,64	0
2	SO4	A	8	5/5	0.94	0.27	64,64,65,65	5
2	SO4	B	1	5/5	0.96	0.13	72,73,73,74	0
2	SO4	B	2	5/5	0.97	0.17	60,61,61,61	5
2	SO4	B	5	5/5	0.98	0.14	63,65,66,66	0
2	SO4	B	9	5/5	0.99	0.12	42,45,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.