



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 16, 2023 – 08:08 PM EDT

PDB ID : 2ERG  
Title : Crystal Structure of Leu3 DNA-binding domain with a single H50C mutation complexed with a 15mer DNA duplex  
Authors : Fitzgerald, M.X.; Marmorstein, R.  
Deposited on : 2005-10-24  
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1804 (3.20-3.12)
Ramachandran outliers	138981	1770 (3.20-3.12)
Sidechain outliers	138945	1769 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	15	13% (green), 80% (yellow), 7% (orange)
1	D	15	13% (green), 87% (yellow)
2	A	72	32% (green), 44% (yellow), 12% (orange), 10% (grey)
2	B	72	33% (green), 44% (yellow), 11% (orange), 10% (grey)

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*TP\*TP\*GP\*CP\*CP\*GP\*GP\*TP\*AP\*CP\*CP\*GP\*GP\*CP\*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	15	Total	C	N	O	P	0	0	0
			304	145	56	89	14			
1	D	15	Total	C	N	O	P	0	0	0
			304	145	56	89	14			

- Molecule 2 is a protein called Regulatory protein LEU3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	65	Total	C	N	O	S	0	0	0
			459	278	90	85	6			
2	B	65	Total	C	N	O	S	0	0	0
			459	278	90	85	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	CYS	HIS	engineered mutation	UNP P08638
B	50	CYS	HIS	engineered mutation	UNP P08638

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	2	Total O 2 2	0	0
4	D	2	Total O 2 2	0	0
4	A	4	Total O 4 4	0	0
4	B	5	Total O 5 5	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: 5'-D>(\*TP\*TP\*GP\*CP\*CP\*GP\*GP\*TP\*AP\*CP\*CP\*GP\*GP\*CP\*A)-3'

Chain C: 

T1 T2 G3 C4 C5 G6 G7 T8 A9 C10 C11 C12 G13 G14 A15

- Molecule 1: 5'-D(\*TP\*TP\*GP\*CP\*CP\*GP\*GP\*TP\*AP\*CP\*CP\*GP\*GP\*CP\*A)-3'

Chain D: 

T16 T17 G18 C19 C20 G21 G22 T23 A24 C25 C26 G27 G28 C29 A30


- Molecule 2: Regulatory protein LEU3

Chain A: 

LYS ARG LYS F35 A36 C37 V38 E39 C40 R41 Q42 Q43 K44 C47 D48 A49 C50 E51 R52 A53 P56 C57 T58 K59 C60 A61 K62 K63 M64 I68 L69 K70 R71 D72 F73 R74 R75 K78 R79 A80 R81 M82 E83 A84 I85 K90 E91 L92 T93 R94 T97 M98 L99 THR

SER ASP ASP GLU

- Molecule 2: Regulatory protein LEU3

Chain B: 

LYS ARG LYS F35 A36 C37 V38 E39 C40 R41 Q42 Q43 K44 C47 D48 A49 C50 E51 R52 A53 P56 C57 T58 A61 K62 K63 M64 I68 L69 K70 R71 D72 F73 R74 R75 K78 R79 A80 R81 M82 E83 A84 I85 K90 E91 L92 T93 R94 T97 M98 L99 THR SER

ASP GLU

## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.77Å 66.77Å 122.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.09 – 3.15	Depositor
% Data completeness (in resolution range)	99.8 (42.09-3.15)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.273 , 0.277	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1543	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	80.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	C	0.66	0/340	0.78	0/523
1	D	0.65	0/340	0.77	0/523
2	A	0.48	0/464	0.66	0/627
2	B	0.45	0/464	0.66	0/627
All	All	0.55	0/1608	0.71	0/2300

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	7	DG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	304	0	170	25	0
1	D	304	0	170	25	0
2	A	459	0	387	47	0
2	B	459	0	387	46	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	4	0	0	0	0
4	B	5	0	0	3	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
All	All	1543	0	1114	132	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 50.

The worst 5 of 132 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:12:DG:H2''	1:C:13:DG:H5''	1.39	1.03
1:D:27:DG:H2''	1:D:28:DG:H5''	1.45	0.95
1:D:17:DT:H2''	1:D:18:DG:C8	2.10	0.86
1:D:27:DG:C2'	1:D:28:DG:H5''	2.08	0.83
1:D:18:DG:H2''	1:D:19:DC:H5'	1.61	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	63/72 (88%)	34 (54%)	19 (30%)	10 (16%)	<b>0</b> <b>0</b>
2	B	63/72 (88%)	34 (54%)	19 (30%)	10 (16%)	<b>0</b> <b>0</b>
All	All	126/144 (88%)	68 (54%)	38 (30%)	20 (16%)	<b>0</b> <b>0</b>



5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	51	GLU
2	A	52	ARG
2	A	53	ALA
2	A	71	ARG
2	A	79	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	39/66 (59%)	35 (90%)	4 (10%)	7 26
2	B	39/66 (59%)	35 (90%)	4 (10%)	7 26
All	All	78/132 (59%)	70 (90%)	8 (10%)	7 26

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	81	ARG
2	B	69	LEU
2	B	35	PHE
2	A	81	ARG
2	B	57	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	43	GLN
2	A	82	ASN
2	B	43	GLN
2	B	82	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.