



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 6, 2023 – 05:34 AM EDT

PDB ID : 4ERM  
Title : Crystal structure of the dATP inhibited E. coli class Ia ribonucleotide reductase complex at 4 Angstroms resolution  
Authors : Zimanyi, C.M.; Drennan, C.L.  
Deposited on : 2012-04-20  
Resolution : 3.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1025 (4.22-3.70)
Clashscore	141614	1085 (4.22-3.70)
Ramachandran outliers	138981	1047 (4.22-3.70)
Sidechain outliers	138945	1039 (4.22-3.70)
RSRZ outliers	127900	1013 (4.28-3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	761	 42% 52% . .
1	B	761	 44% 49% . .
1	C	761	 45% 47% . .
1	D	761	 46% 47% . .
2	E	375	 44% 47% . 6%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	375	<p>% 44% 49% • 5%</p>
2	G	375	<p>% 41% 49% • • 5%</p>
2	H	375	<p>2% 45% 48% • 5%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DAT	D	802	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 35345 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase 1 subunit alpha.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	733	Total 5841	C 3710	N 1003	O 1104	S 24	0	0	0
1	B	734	Total 5845	C 3712	N 1004	O 1105	S 24	0	0	0
1	C	731	Total 5821	C 3696	N 1000	O 1101	S 24	0	0	0
1	D	733	Total 5837	C 3708	N 1002	O 1103	S 24	0	0	0

- Molecule 2 is a protein called Ribonucleoside-diphosphate reductase 1 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	352	Total 2885	C 1841	N 478	O 553	S 13	0	0	0
2	F	356	Total 2917	C 1861	N 483	O 560	S 13	0	0	0
2	G	356	Total 2917	C 1861	N 483	O 560	S 13	0	0	0
2	H	356	Total 2914	C 1859	N 482	O 560	S 13	0	0	0

- Molecule 3 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).






Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	Total	C	N	O	P	0	0
			26	10	5	9	2		
4	B	1	Total	C	N	O	P	0	0
			26	10	5	9	2		
4	C	1	Total	C	N	O	P	0	0
			26	10	5	9	2		
4	D	1	Total	C	N	O	P	0	0
			26	10	5	9	2		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		
5	D	1	Total	Mg	0	0
			1	1		

- Molecule 6 is MU-OXO-DIIRON (three-letter code: FEO) (formula: Fe<sub>2</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	E	1	Total	Fe	O	0	0
			3	2	1		
6	F	1	Total	Fe	O	0	0
			3	2	1		
6	G	1	Total	Fe	O	0	0
			3	2	1		
6	H	1	Total	Fe	O	0	0
			3	2	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	2	Total	O	0	0
			2	2		
7	F	2	Total	O	0	0
			2	2		
7	G	2	Total	O	0	0
			2	2		
7	H	2	Total	O	0	0
			2	2		













• Molecule 1: Ribonucleoside-diphosphate reductase 1 subunit alpha







• Molecule 2: Ribonucleoside-diphosphate reductase 1 subunit beta





• Molecule 2: Ribonucleoside-diphosphate reductase 1 subunit beta







• Molecule 2: Ribonucleoside-diphosphate reductase 1 subunit beta



• Molecule 2: Ribonucleoside-diphosphate reductase 1 subunit beta





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	280.47Å 155.74Å 166.92Å 90.00° 119.07° 90.00°	Depositor
Resolution (Å)	30.00 – 3.95 29.99 – 3.93	Depositor EDS
% Data completeness (in resolution range)	96.1 (30.00-3.95) 95.2 (29.99-3.93)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.16	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.62 (at 3.98Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.259 , 0.284 0.241 , 0.264	Depositor DCC
$R_{free}$ test set	2527 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	95.3	Xtrriage
Anisotropy	0.470	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 61.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	35345	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DAT, FEO, DTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/5969	0.46	1/8085 (0.0%)
1	B	0.25	0/5973	0.45	1/8090 (0.0%)
1	C	0.25	0/5949	0.44	0/8057
1	D	0.25	0/5965	0.44	1/8079 (0.0%)
2	E	0.30	1/2949 (0.0%)	0.44	1/3998 (0.0%)
2	F	0.25	0/2981	0.41	0/4042
2	G	0.25	0/2981	0.42	0/4042
2	H	0.25	0/2978	0.43	0/4038
All	All	0.26	1/35745 (0.0%)	0.44	4/48431 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	203	LEU	C-N	5.68	1.47	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	48	GLN	O-C-N	5.88	132.10	122.70
1	A	48	GLN	O-C-N	5.86	132.08	122.70
1	B	48	GLN	O-C-N	5.85	132.07	122.70
2	E	203	LEU	C-N-CA	-5.43	108.12	121.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5841	0	5767	494	0
1	B	5845	0	5770	497	0
1	C	5821	0	5742	483	0
1	D	5837	0	5764	459	0
2	E	2885	0	2813	211	0
2	F	2917	0	2845	240	0
2	G	2917	0	2845	285	0
2	H	2914	0	2842	232	0
3	A	60	0	22	1	0
3	B	60	0	22	8	0
3	C	60	0	22	10	0
3	D	60	0	22	9	0
4	A	26	0	12	3	0
4	B	26	0	12	1	0
4	C	26	0	12	5	0
4	D	26	0	12	9	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	E	3	0	0	0	0
6	F	3	0	0	0	0
6	G	3	0	0	0	0
6	H	3	0	0	0	0
7	E	2	0	0	1	0
7	F	2	0	0	1	0
7	G	2	0	0	0	0
7	H	2	0	0	1	0
All	All	35345	0	34524	2786	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 40.

The worst 5 of 2786 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:354:THR:HG21	1:B:356:PHE:CZ	1.44	1.52
1:A:646:ALA:HB2	1:A:651:ILE:CG2	1.40	1.52
1:A:646:ALA:CB	1:A:651:ILE:HG22	1.41	1.46
1:A:297:VAL:CG1	1:A:298:ARG:HG2	1.50	1.41
1:A:53:ILE:HD11	1:A:58:ILE:CD1	1.50	1.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	731/761 (96%)	687 (94%)	40 (6%)	4 (0%)	29	66
1	B	732/761 (96%)	693 (95%)	37 (5%)	2 (0%)	41	74
1	C	729/761 (96%)	690 (95%)	36 (5%)	3 (0%)	34	70
1	D	731/761 (96%)	694 (95%)	33 (4%)	4 (0%)	29	66
2	E	348/375 (93%)	336 (97%)	12 (3%)	0	100	100
2	F	352/375 (94%)	338 (96%)	13 (4%)	1 (0%)	41	74
2	G	352/375 (94%)	336 (96%)	13 (4%)	3 (1%)	17	54
2	H	352/375 (94%)	339 (96%)	13 (4%)	0	100	100
All	All	4327/4544 (95%)	4113 (95%)	197 (5%)	17 (0%)	34	70

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	GLY
1	A	300	GLY
1	B	300	GLY
1	C	299	GLY
1	C	300	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	629/651 (97%)	593 (94%)	36 (6%)	20	49
1	B	629/651 (97%)	594 (94%)	35 (6%)	21	49
1	C	626/651 (96%)	591 (94%)	35 (6%)	21	49
1	D	628/651 (96%)	592 (94%)	36 (6%)	20	49
2	E	319/340 (94%)	305 (96%)	14 (4%)	28	54
2	F	323/340 (95%)	311 (96%)	12 (4%)	34	59
2	G	323/340 (95%)	299 (93%)	24 (7%)	13	41
2	H	323/340 (95%)	312 (97%)	11 (3%)	37	61
All	All	3800/3964 (96%)	3597 (95%)	203 (5%)	22	51

5 of 203 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	80	LEU
2	E	51	GLU
2	H	295	SER
1	D	154	LYS
1	D	474	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 135 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	143	ASN
2	G	247	HIS
2	H	281	GLN
1	C	158	GLN
1	C	130	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	DTP	B	801	-	26,32,32	1.63	6 (23%)	30,50,50	1.45	4 (13%)
4	DAT	C	802	-	24,28,28	1.67	8 (33%)	28,43,43	1.50	3 (10%)
3	DTP	A	801	-	26,32,32	1.62	6 (23%)	30,50,50	1.36	4 (13%)
3	DTP	C	803	5	26,32,32	1.59	5 (19%)	30,50,50	1.38	4 (13%)
4	DAT	A	802	-	24,28,28	1.57	7 (29%)	28,43,43	1.56	4 (14%)
6	FEO	H	501	7,2	0,2,2	-	-	-		
6	FEO	G	501	7,2	0,2,2	-	-	-		
3	DTP	D	801	-	26,32,32	1.62	6 (23%)	30,50,50	1.36	4 (13%)
4	DAT	B	802	-	24,28,28	1.58	7 (29%)	28,43,43	1.52	4 (14%)
3	DTP	A	803	5	26,32,32	1.58	5 (19%)	30,50,50	1.41	4 (13%)
6	FEO	F	501	7,2	0,2,2	-	-	-		
3	DTP	C	801	-	26,32,32	1.62	6 (23%)	30,50,50	1.36	4 (13%)
3	DTP	D	803	5	26,32,32	1.60	5 (19%)	30,50,50	1.40	4 (13%)
3	DTP	B	803	5	26,32,32	1.59	5 (19%)	30,50,50	1.41	4 (13%)
6	FEO	E	501	7,2	0,2,2	-	-	-		
4	DAT	D	802	-	24,28,28	1.59	7 (29%)	28,43,43	1.46	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DTP	B	801	-	-	6/18/34/34	0/3/3/3
4	DAT	C	802	-	-	3/12/28/28	0/3/3/3
3	DTP	A	801	-	-	4/18/34/34	0/3/3/3
3	DTP	C	803	5	-	4/18/34/34	0/3/3/3
4	DAT	A	802	-	-	3/12/28/28	0/3/3/3
3	DTP	D	801	-	-	4/18/34/34	0/3/3/3
4	DAT	B	802	-	-	8/12/28/28	0/3/3/3
3	DTP	A	803	5	-	1/18/34/34	0/3/3/3
3	DTP	C	801	-	-	4/18/34/34	0/3/3/3
3	DTP	D	803	5	-	5/18/34/34	0/3/3/3
3	DTP	B	803	5	-	2/18/34/34	0/3/3/3
4	DAT	D	802	-	-	4/12/28/28	0/3/3/3

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	803	DTP	O3'-C3'	-4.05	1.34	1.43
3	B	801	DTP	O3'-C3'	-4.01	1.34	1.43
3	D	803	DTP	O3'-C3'	-4.00	1.34	1.43
3	B	803	DTP	O3'-C3'	-3.96	1.35	1.43
3	A	801	DTP	O3'-C3'	-3.91	1.35	1.43

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	802	DAT	N3-C2-N1	-4.55	121.57	128.68
4	C	802	DAT	N3-C2-N1	-4.52	121.61	128.68
4	B	802	DAT	N3-C2-N1	-4.48	121.68	128.68
4	D	802	DAT	N3-C2-N1	-4.40	121.81	128.68
3	B	801	DTP	PB-O3B-PG	-3.95	119.25	132.83

There are no chirality outliers.

5 of 48 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	DTP	C5'-O5'-PA-O1A
3	A	801	DTP	C5'-O5'-PA-O2A

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	A	803	DTP	PB-O3B-PG-O3G
3	B	801	DTP	C5'-O5'-PA-O3A
3	C	801	DTP	C5'-O5'-PA-O1A


There are no ring outliers.



11 monomers are involved in 46 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	801	DTP	5	0
4	C	802	DAT	5	0
3	C	803	DTP	3	0
4	A	802	DAT	3	0
3	D	801	DTP	5	0
4	B	802	DAT	1	0
3	A	803	DTP	1	0
3	C	801	DTP	7	0
3	D	803	DTP	4	0
3	B	803	DTP	3	0
4	D	802	DAT	9	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	733/761 (96%)	-0.10	9 (1%) 79 70	56, 95, 121, 142	0
1	B	734/761 (96%)	-0.04	10 (1%) 75 66	59, 101, 127, 151	0
1	C	731/761 (96%)	0.19	27 (3%) 41 32	101, 130, 150, 160	0
1	D	733/761 (96%)	0.22	27 (3%) 41 32	89, 128, 151, 163	0
2	E	352/375 (93%)	0.03	12 (3%) 45 36	72, 117, 154, 176	0
2	F	356/375 (94%)	-0.13	5 (1%) 75 66	55, 91, 131, 158	0
2	G	356/375 (94%)	-0.09	3 (0%) 86 79	59, 97, 145, 192	0
2	H	356/375 (94%)	-0.07	8 (2%) 62 52	65, 98, 145, 174	0
All	All	4351/4544 (95%)	0.02	101 (2%) 60 51	55, 110, 146, 192	0

The worst 5 of 101 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	737	GLY	6.3
2	H	363	SER	4.4
1	D	298	ARG	4.1
1	D	297	VAL	3.4
1	D	617	SER	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	DAT	D	802	26/26	0.73	0.34	156,156,156,156	0
3	DTP	C	801	30/30	0.76	0.35	175,175,175,175	0
4	DAT	B	802	26/26	0.78	0.36	149,149,149,149	0
4	DAT	C	802	26/26	0.82	0.28	156,156,156,156	0
4	DAT	A	802	26/26	0.83	0.29	126,126,126,126	0
3	DTP	D	801	30/30	0.83	0.36	140,140,140,140	0
3	DTP	B	801	30/30	0.90	0.32	111,111,111,111	0
3	DTP	A	801	30/30	0.91	0.22	124,124,124,124	0
5	MG	D	804	1/1	0.92	0.28	118,118,118,118	0
5	MG	B	804	1/1	0.93	0.18	94,94,94,94	0
3	DTP	B	803	30/30	0.94	0.16	94,94,94,94	0
3	DTP	D	803	30/30	0.94	0.15	118,118,118,118	0
3	DTP	A	803	30/30	0.94	0.16	94,94,94,94	0
3	DTP	C	803	30/30	0.94	0.16	123,123,123,123	0
5	MG	A	804	1/1	0.95	0.13	94,94,94,94	0
6	FEO	G	501	3/3	0.96	0.18	78,78,78,78	0
5	MG	C	804	1/1	0.97	0.13	123,123,123,123	0
6	FEO	H	501	3/3	0.97	0.14	69,69,69,69	0
6	FEO	F	501	3/3	0.98	0.15	61,61,61,61	0
6	FEO	E	501	3/3	0.99	0.15	89,89,89,89	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













**Electron density around DAT B 802:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

















**Electron density around DTP A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)














**Electron density around DTP D 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around DTP A 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.