

wwPDB X-ray Structure Validation Summary Report (i)

Oct 23, 2021 – 02:37 PM EDT

PDB ID : 1EX4

Title : HIV-1 INTEGRASE CATALYTIC CORE AND C-TERMINAL DOMAIN Authors : Chen, J.C.-H.; Krucinski, J.; Miercke, L.J.W.; Finer-Moore, J.S.; Tang, A.H.;

Leavitt, A.D.; Stroud, R.M.

Deposited on : 2000-04-28

Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.23.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain				
1	A	239	43%	41%	5% 11%		
1	В	239	38%	42%	• 15%		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CPS	A	302	X	-	-	-
2	CPS	A	304	X	-	-	-
2	CPS	В	301	X	-	-	-
2	CPS	В	303	X	-	-	-



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3450 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called INTEGRASE.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	212	Total 1653	C 1046	11	O 304	S 4	0	0	0
1	В	204	Total 1601	C 1016	N 290	O 291	S 4	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	GLY	-	cloning artifact	UNP P04585
A	51	SER	-	cloning artifact	UNP P04585
A	56	SER	CYS	engineered mutation	UNP P04585
A	131	ASP	TRP	engineered mutation	UNP P04585
A	139	ASP	PHE	engineered mutation	UNP P04585
A	185	LYS	PHE	engineered mutation	UNP P04585
A	280	SER	CYS	engineered mutation	UNP P04585
В	50	GLY	-	cloning artifact	UNP P04585
В	51	SER	_	cloning artifact	UNP P04585
В	56	SER	CYS	engineered mutation	UNP P04585
В	131	ASP	TRP	engineered mutation	UNP P04585
В	139	ASP	PHE	engineered mutation	UNP P04585
В	185	LYS	PHE	engineered mutation	UNP P04585
В	280	SER	CYS	engineered mutation	UNP P04585

• Molecule 2 is 3-[(3-CHOLAMIDOPROPYL)DIMETHYLAMMONIO]-1-PROPANESULFO NATE (three-letter code: CPS) (formula: C₃₂H₅₈N₂O₇S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 25 22 3	0	0
2	A	1	Total C O 25 22 3	0	0
2	В	1	Total C O 25 22 3	0	0
2	В	1	Total C N O S 42 32 2 7 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	${f Atoms}$	$\mathbf{ZeroOcc}$	AltConf	
3	A	49	Total O 49 49	0	0	
3	В	30	Total O 30 30	0	0	




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: INTEGRASE



• Molecule 1: INTEGRASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 3 1 2	Depositor
Cell constants	103.99Å 103.99Å 101.38Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.26 - 2.80	Depositor
% Data completeness	99.7 (28.26-2.80)	Depositor
(in resolution range)	33.1 (20.20 2.00)	Беровног
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.258 , 0.306	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3450	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.36	0/1681	0.64	0/2267	
1	В	0.35	0/1628	0.60	0/2195	
All	All	0.36	0/3309	0.62	0/4462	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1653	0	1688	113	1
1	В	1601	0	1647	118	0
2	A	50	0	68	7	2
2	В	67	0	91	18	0
3	A	49	0	0	4	0
3	В	30	0	0	10	1
All	All	3450	0	3494	235	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 235 close contacts within the same asymmetric unit are listed below, sorted by



their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	Clash overlap (Å)
1:A:269:ARG:H	1:A:269:ARG:HD2	1.32	0.95
1:B:221:GLN:HE21	1:B:221:GLN:HA	1.30	0.95
1:A:177:GLN:HE21	1:A:177:GLN:HA	1.36	0.90
1:A:95:GLN:HE21	1:A:95:GLN:HA	1.37	0.88
1:A:177:GLN:HA	1:A:177:GLN:NE2	1.93	0.82

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:216:GLN:OE1	2:A:304:CPS:O2[6_565]	2.15	0.05
2:A:302:CPS:C16	3:B:315:HOH:O[3_565]	2.16	0.04

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles			
1	A	208/239 (87%)	167 (80%)	27 (13%)	14 (7%)	1 3			
1	В	200/239 (84%)	165 (82%)	24 (12%)	11 (6%)	2 5			
All	All	408/478 (85%)	332 (81%)	51 (12%)	25 (6%)	1 4			

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	GLU
1	В	79	VAL
1	A	119	SER
1	A	123	GLY
1	A	212	GLU



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	174/196 (89%)	166 (95%)	8 (5%)	27	60	
1	В	169/196 (86%)	161 (95%)	8 (5%)	26	59	
All	All	343/392 (88%)	327 (95%)	16 (5%)	26	59	

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	242	LEU
1	В	221	GLN
1	В	102	LEU
1	В	220	ILE
1	A	269	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	155	ASN
1	В	214	GLN
1	В	183	HIS
1	В	216	GLN
1	A	177	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Во	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2	
2	CPS	A	302	-	28,28,45	0.91	1 (3%)	46,46,70	4.42	26 (56%)	
2	CPS	В	301	-	28,28,45	0.82	1 (3%)	46,46,70	3.73	23 (50%)	
2	CPS	В	303	-	45,45,45	1.28	4 (8%)	69,70,70	3.74	26 (37%)	
2	CPS	A	304	-	28,28,45	0.92	2 (7%)	46,46,70	3.98	21 (45%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CPS	A	302	-	3/3/11/15	4/4/69/90	0/4/4/4
2	CPS	В	301	-	3/3/11/15	0/4/69/90	0/4/4/4
2	CPS	В	303	-	4/4/14/15	10/25/90/90	0/4/4/4
2	CPS	A	304	-	3/3/11/15	0/4/69/90	0/4/4/4

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	В	303	CPS	O3S-S	5.55	1.61	1.45
2	В	303	CPS	C2-C19	-2.77	1.51	1.56
2	В	303	CPS	C24-N1	2.76	1.39	1.33
2	В	301	CPS	C2-C19	-2.74	1.51	1.56
2	A	304	CPS	C2-C19	-2.59	1.51	1.56



The worst	5	of 9	96	bond	angle	outliers	are	listed	below:
TIIC WOID	$\mathbf{\mathcal{I}}$	OI 6	, 0	Olia	ansi	Outilors	COL	iibuca	DOIOW.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	В	303	CPS	C2-C19-C18	16.56	129.60	111.82
2	A	304	CPS	C3-C19-C2	12.46	126.58	113.73
2	A	304	CPS	C2-C19-C18	11.59	124.26	111.82
2	A	302	CPS	C6-C5-C4	11.30	117.92	107.40
2	В	301	CPS	C3-C19-C2	11.12	125.20	113.73

5 of 13 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	302	CPS	C18
2	A	302	CPS	C6
2	A	302	CPS	C19
2	A	304	CPS	C18
2	A	304	CPS	C6

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	302	CPS	C21-C20-C9-C5
2	A	302	CPS	C22-C20-C9-C5
2	A	302	CPS	C21-C20-C9-C8
2	A	302	CPS	C22-C20-C9-C8
2	В	303	CPS	C21-C20-C9-C8

There are no ring outliers.


3 monomers are involved in 27 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	302	CPS	5	1
2	В	303	CPS	18	0
2	A	304	CPS	2	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and




any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






Rings

Torsions





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

