



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2021 – 12:12 PM JST

PDB ID : 7EXM
Title : The N-terminal crystal structure of SARS-CoV-2 NSP2
Authors : Ma, J.; Chen, Z.
Deposited on : 2021-05-27
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	 8% 92% 8% .
1	B	278	 % 91% 6% ..
1	C	278	 6% 95% ..
1	D	278	 % 94% ..

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-structural protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	278	2124	1334	361	407	22	0	3	0
1	B	273	2092	1315	355	399	23	0	2	0
1	C	274	2080	1303	355	400	22	0	0	0
1	D	274	2110	1322	362	404	22	0	2	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P0DTC1
A	1	MET	-	expression tag	UNP P0DTC1
B	0	SER	-	expression tag	UNP P0DTC1
B	1	MET	-	expression tag	UNP P0DTC1
C	0	SER	-	expression tag	UNP P0DTC1
C	1	MET	-	expression tag	UNP P0DTC1
D	0	SER	-	expression tag	UNP P0DTC1
D	1	MET	-	expression tag	UNP P0DTC1

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	Zn 3	0	0
2	B	3	Total 3	Zn 3	0	0
2	C	3	Total 3	Zn 3	0	0
2	D	3	Total 3	Zn 3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	140	Total	O	0	0
			140	140		
4	B	288	Total	O	0	0
			288	288		
4	C	246	Total	O	0	0
			246	246		
4	D	251	Total	O	0	0
			251	251		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.93Å 159.60Å 63.55Å 90.00° 91.19° 90.00°	Depositor
Resolution (Å)	42.40 – 1.96 42.36 – 1.74	Depositor EDS
% Data completeness (in resolution range)	95.9 (42.40-1.96) 74.1 (42.36-1.74)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.74Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.200 , 0.226 0.211 , 0.215	Depositor DCC
R_{free} test set	4226 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.045 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9349	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/2182	0.73	0/2961
1	B	0.67	0/2146	0.72	0/2909
1	C	0.68	0/2126	0.71	0/2878
1	D	0.68	0/2164	0.72	0/2933
All	All	0.68	0/8618	0.72	0/11681

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2124	0	1987	10	0
1	B	2092	0	1968	13	0
1	C	2080	0	1936	6	0
1	D	2110	0	1994	3	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
2	C	3	0	0	0	0
2	D	3	0	0	0	0
3	C	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	140	0	0	0	0
4	B	288	0	0	1	0
4	C	246	0	0	0	0
4	D	251	0	0	1	0
All	All	9349	0	7893	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:ASN:HD22	1:B:40:LEU:HD11	1.53	0.72
1:A:0:SER:CB	1:A:17:TYR:HB2	2.35	0.56
1:D:219:ARG:NH1	4:D:401:HOH:O	2.30	0.49
1:A:46:LYS:O	1:A:51:CYS:HA	2.12	0.49
1:B:180:TYR:HB2	1:D:33:SER:HB3	1.95	0.48
1:A:108:ARG:NE	1:A:122:ARG:O	2.38	0.47
1:A:76:PHE:CE1	1:A:247:ARG:HB3	2.49	0.47
1:B:106:GLN:HG3	1:C:86:THR:CG2	2.44	0.47
1:A:63[A]:THR:HG22	1:A:65:ARG:HG3	1.97	0.47
1:A:105:ILE:HG21	1:A:124:VAL:HG13	1.98	0.46
1:C:104:THR:HA	1:C:247:ARG:O	2.17	0.46
1:B:106:GLN:HG3	1:C:86:THR:HG22	1.98	0.45
1:A:118:MET:HA	1:A:118:MET:HE2	1.98	0.45
1:B:162:CYS:SG	1:B:164:PHE:HB2	2.57	0.45
1:B:46:LYS:O	1:B:51:CYS:HA	2.16	0.45
1:A:272:LEU:O	1:A:276:GLN:HG2	2.17	0.44
1:B:25:LEU:C	1:B:25:LEU:HD23	2.38	0.44
1:A:254:CYS:HA	1:C:71:GLU:OE1	2.19	0.42
1:B:196:ASN:HB3	1:B:199:VAL:HG13	2.02	0.42
1:B:97:PRO:HG2	1:B:100:SER:HB3	2.02	0.41
1:B:104:THR:HA	1:B:247:ARG:O	2.20	0.41
1:A:209:HIS:CG	1:A:223:ARG:HD3	2.56	0.41
1:C:17:TYR:CZ	1:C:219:ARG:HG3	2.56	0.41
1:B:109:VAL:N	4:B:416:HOH:O	2.53	0.41
1:B:196:ASN:C	1:B:196:ASN:HD22	2.24	0.41
1:D:110:GLU:OE2	1:D:122:ARG:NH2	2.54	0.41
1:B:46:LYS:O	1:B:47:ARG:HB2	2.20	0.40
1:C:142:MET:CE	1:C:160:ALA:HB3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/278 (100%)	269 (96%)	8 (3%)	2 (1%)	22	11
1	B	271/278 (98%)	261 (96%)	9 (3%)	1 (0%)	34	22
1	C	268/278 (96%)	259 (97%)	9 (3%)	0	100	100
1	D	274/278 (99%)	266 (97%)	8 (3%)	0	100	100
All	All	1092/1112 (98%)	1055 (97%)	34 (3%)	3 (0%)	41	30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	47	ARG
1	B	47	ARG
1	A	199	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/238 (94%)	218 (97%)	6 (3%)	44	34
1	B	224/238 (94%)	218 (97%)	6 (3%)	44	34
1	C	220/238 (92%)	217 (99%)	3 (1%)	67	62
1	D	228/238 (96%)	219 (96%)	9 (4%)	32	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	896/952 (94%)	872 (97%)	24 (3%)	47 34

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	63[A]	THR
1	A	63[B]	THR
1	A	217	ILE
1	A	224	THR
1	A	247	ARG
1	B	56	GLU
1	B	91	CYS
1	B	196	ASN
1	B	199	VAL
1	B	219	ARG
1	B	277	LYS
1	C	71	GLU
1	C	131	ASN
1	C	135	GLN
1	D	28[A]	ARG
1	D	28[B]	ARG
1	D	53	ARG
1	D	71	GLU
1	D	135	GLN
1	D	217	ILE
1	D	247	ARG
1	D	273	GLU
1	D	275	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
1	B	10	ASN
1	B	196	ASN
1	C	99	ASN
1	C	134	ASN
1	C	183	GLN
1	D	93	ASN
1	D	99	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	169	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 12 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	C	304	-	5,5,5	0.10	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	304	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	304	GOL	C1-C2-C3-O3
3	C	304	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	278/278 (100%)	0.32	21 (7%) 13 21	24, 37, 70, 81	0
1	B	273/278 (98%)	-0.11	2 (0%) 87 92	16, 26, 41, 73	0
1	C	274/278 (98%)	0.22	16 (5%) 23 31	21, 36, 63, 69	0
1	D	274/278 (98%)	-0.15	4 (1%) 73 81	19, 28, 47, 61	0
All	All	1099/1112 (98%)	0.07	43 (3%) 39 49	16, 31, 60, 81	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	199	VAL	6.6
1	B	115	ASP	4.4
1	A	1	MET	4.3
1	C	201	PRO	4.0
1	C	203	HIS	4.0
1	A	199	VAL	3.9
1	A	277	LYS	3.8
1	A	109	VAL	3.8
1	A	2	ALA	3.7
1	A	113	LYS	3.4
1	A	200	GLY	3.4
1	A	196	ASN	3.3
1	A	210	ASN	3.2
1	C	189	ILE	3.1
1	D	53	ARG	3.1
1	C	117	PHE	2.9
1	A	0	SER	2.9
1	A	197	SER	2.9
1	C	120	ARG	2.8
1	C	205	LEU	2.8
1	C	217	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	241	CYS	2.8
1	C	115	ASP	2.6
1	D	50	TYR	2.6
1	C	193	ALA	2.6
1	A	195	HIS	2.5
1	A	120	ARG	2.5
1	A	207	GLU	2.4
1	A	192	PRO	2.4
1	A	276	GLN	2.3
1	C	208	TYR	2.3
1	C	90	GLU	2.2
1	A	137	CYS	2.2
1	A	190	TYR	2.2
1	D	112	LYS	2.2
1	C	197	SER	2.1
1	B	3	TYR	2.1
1	C	211	GLU	2.0
1	A	154[A]	THR	2.0
1	C	196	ASN	2.0
1	A	164	PHE	2.0
1	A	211	GLU	2.0
1	D	262	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	C	304	6/6	0.81	0.14	52,54,55,55	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	A	303	1/1	0.89	0.09	44,44,44,44	0
2	ZN	C	302	1/1	0.92	0.06	48,48,48,48	0
2	ZN	C	303	1/1	0.94	0.05	48,48,48,48	0
2	ZN	D	301	1/1	0.98	0.07	35,35,35,35	0
2	ZN	B	303	1/1	0.98	0.08	23,23,23,23	0
2	ZN	C	301	1/1	0.99	0.09	23,23,23,23	0
2	ZN	A	301	1/1	0.99	0.07	26,26,26,26	0
2	ZN	B	301	1/1	0.99	0.07	25,25,25,25	0
2	ZN	B	302	1/1	0.99	0.07	23,23,23,23	0
2	ZN	D	302	1/1	0.99	0.07	25,25,25,25	0
2	ZN	D	303	1/1	0.99	0.07	24,24,24,24	0
2	ZN	A	302	1/1	0.99	0.06	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ZN A 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN D 302:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN D 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.