

Full wwPDB X-ray Structure Validation Report (i)

May 14, 2020 – 05:02 am BST

PDB ID : 1EYR

Title: Structure of a sialic acid activating synthetase, CMP acylneuraminate syn-

thetase in the presence and absence of CDP

Authors: Mosimann, S.C.; Gilbert, M.; Dombrowski, D.; Wakarchuk, W.; Strynadka,

N.C.

Deposited on : 2000-05-08

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED

EDS : NOT EXECUTED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	228	82%	14%	
1	В	228	75%	21%	•••



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called CMP-N-ACETYLNEURAMINIC ACID SYNTHETASE.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	225		C 1079				0	0	0
1	В	221	Total 1680	C 1055				0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
A	26	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
A	136	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
A	155	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
A	202	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
В	1	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
В	26	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
В	136	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
В	155	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8
В	202	MSE	MET	MODIFIED RESIDUE	UNP P0A0Z8

• Molecule 2 is CYTIDINE-5'-DIPHOSPHATE (three-letter code: CDP) (formula: C₉H₁₅N₃O₁₁P₂).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Р	0	1
	A	1	50	18	6	22	4	U	1
9	D	1	Total	С	N	О	Р	0	1
	D	1	50	18	6	22	4	U	1

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	242	Total O 242 242	0	2
3	В	208	Total O 208 208	0	2




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CMP-N-ACETYLNEURAMINIC ACID SYNTHETASE





4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	43.01Å 69.89Å 157.48Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.90 - 2.20	Depositor
% Data completeness	87.9 (19.90-2.20)	Depositor
(in resolution range)	,	Берозног
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.185 , 0.246	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3952	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.49	0/1746	0.63	$1/2360 \ (0.0\%)$	
1	В	0.46	0/1702	0.63	0/2302	
All	All	0.48	0/3448	0.63	1/4662~(0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	A	186	LEU	CA-CB-CG	6.21	129.57	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1722	0	1744	29	0
1	В	1680	0	1693	37	0
2	A	50	0	24	0	0
2	В	50	0	24	0	0
3	A	242	0	0	2	0
3	В	208	0	0	3	0
All	All	3952	0	3485	59	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:165:ARG:HD2	1:B:165:ARG:H	1.42	0.85
1:B:154:PRO:HG3	1:B:158:LEU:HD22	1.70	0.72
1:B:6:ILE:HD13	1:B:93:ILE:HD11	1.70	0.71
3:A:1219:HOH:O	1:B:158:LEU:HD21	1.91	0.70
1:B:208:ILE:HD13	1:B:219:ALA:HB2	1.72	0.70
1:B:205:GLN:HG2	1:B:222:ILE:HD13	1.72	0.70
1:A:191:CYS:HB2	1:B:162:GLU:OE2	1.95	0.67
1:A:96:ASN:C	1:A:96:ASN:HD22	1.97	0.66
1:B:156:ARG:HB3	1:B:160:ASP:OD2	1.97	0.64
1:A:26:MSE:HE3	1:A:223:LEU:HD12	1.80	0.64
1:A:26:MSE:HE3	1:A:223:LEU:CD1	2.28	0.64
1:B:26:MSE:HE3	1:B:223:LEU:HD11	1.80	0.63
1:B:113:ALA:O	1:B:117:GLU:HG3	1.99	0.62
1:A:26:MSE:HG3	1:A:220:GLU:HG2	1.84	0.60
1:A:155:MSE:HE3	1:B:140:PRO:HB2	1.84	0.59
1:B:175:ASN:C	1:B:175:ASN:HD22	2.07	0.57
1:A:136:MSE:HE2	1:B:143:THR:HG21	1.86	0.57
1:A:147:ILE:HD11	1:A:153:ALA:HB2	1.87	0.56
1:B:104:GLN:HG3	3:B:2032:HOH:O	2.05	0.56
1:A:146:GLN:HB3	1:B:170:GLN:HB2	1.88	0.55
1:B:175:ASN:HD22	1:B:176:GLY:N	2.04	0.55
1:A:162:GLU:OE1	1:B:191:CYS:HB2	2.07	0.53
1:A:223:LEU:C	1:A:225:HIS:H	2.11	0.53
1:A:175:ASN:ND2	1:A:177:ALA:H	2.07	0.52
1:A:146:GLN:CB	1:B:170:GLN:HB2	2.39	0.52
1:A:164:PRO:HG2	1:A:167:GLN:HG2	1.91	0.52
1:A:205:GLN:HG3	1:A:222:ILE:HD13	1.92	0.52
1:A:205:GLN:OE1	1:A:222:ILE:HG23	2.10	0.51
1:B:175:ASN:ND2	1:B:177:ALA:H	2.09	0.51
1:A:96:ASN:ND2	1:A:96:ASN:C	2.62	0.50
1:A:176:GLY:HA2	1:A:179:TYR:OH	2.12	0.49
1:B:166:GLN:NE2	3:B:2117:HOH:O	2.46	0.48
1:A:158:LEU:HD12	1:A:161:LEU:HD12	1.95	0.47
1:A:140:PRO:HB2	1:B:155:MSE:HE3	1.96	0.47
1:B:139:HIS:HB3	1:B:142:LYS:HG3	1.98	0.46
1:A:72:PRO:HD2	1:A:75:LEU:HD12	1.98	0.46
1:B:124:GLU:H	1:B:124:GLU:CD	2.19	0.45

Continued on next page...



Continued from previous page...

A 4 a mag 1	A 4 a ma 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \; ({\rm \AA})$	$overlap (\AA)$
1:B:108:PRO:CD	1:B:210:ILE:HD13	2.47	0.45
1:B:159:SER:O	1:B:163:GLN:HG2	2.17	0.45
1:B:118:ALA:O	1:B:130:VAL:HG11	2.16	0.45
1:B:24:ARG:HD3	3:B:2074:HOH:O	2.17	0.44
1:B:27:ASN:HD22	1:B:27:ASN:HA	1.59	0.44
1:A:26:MSE:HG3	1:A:220:GLU:CG	2.46	0.44
1:B:165:ARG:HD2	1:B:165:ARG:N	2.21	0.43
1:B:148:ASN:HA	1:B:151:GLU:HA	2.01	0.43
1:B:15:SER:HB3	1:B:18:LEU:O	2.18	0.43
1:A:223:LEU:O	1:A:225:HIS:N	2.52	0.43
1:B:108:PRO:CG	1:B:210:ILE:HD13	2.49	0.43
1:B:176:GLY:HA2	1:B:179:TYR:OH	2.19	0.42
1:A:15:SER:HB3	1:A:18:LEU:O	2.20	0.41
1:B:71:ARG:HA	1:B:72:PRO:HD3	1.94	0.41
1:A:36:ILE:HG21	1:A:64:PHE:HB3	2.02	0.41
1:A:48:ILE:C	1:A:49:ILE:HD12	2.41	0.41
1:B:23:LEU:HG	1:B:57:ILE:HD13	2.02	0.41
1:A:104:GLN:HG3	3:A:1101:HOH:O	2.20	0.40
1:A:175:ASN:HD22	1:A:175:ASN:C	2.25	0.40
1:A:175:ASN:HD22	1:A:176:GLY:N	2.20	0.40
1:B:148:ASN:HD22	1:B:151:GLU:HA	1.85	0.40
1:B:6:ILE:HD13	1:B:93:ILE:CD1	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	$223/228 \ (98\%)$	218 (98%)	3 (1%)	2 (1%)	17 16
1	В	$217/228 \ (95\%)$	207 (95%)	9 (4%)	1 (0%)	29 31
All	All	440/456 (96%)	425 (97%)	12 (3%)	3 (1%)	22 22



All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	149	ASN
1	A	224	ASN
1	В	17	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	192/190 (101%)	186 (97%)	6 (3%)	40 51
1	В	186/190 (98%)	180 (97%)	6 (3%)	39 50
All	All	378/380 (100%)	366 (97%)	12 (3%)	39 50

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	ASN
1	A	106	THR
1	A	119	PHE
1	A	175	ASN
1	A	186	LEU
1	A	225	HIS
1	В	27	ASN
1	В	119	PHE
1	В	145	LEU
1	В	175	ASN
1	В	204	HIS
1	В	205	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	27	ASN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	37	ASN
1	A	96	ASN
1	A	104	GLN
1	A	157	HIS
1	A	175	ASN
1	A	221	ASN
1	A	224	ASN
1	В	14	ASN
1	В	27	ASN
1	В	63	ASN
1	В	148	ASN
1	В	163	GLN
1	В	175	ASN
1	В	204	HIS
1	В	218	GLN
1	В	221	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Т	Chain	D. a.g. T. ! 1.		Во	ond leng	$_{ m ths}$	Bond angles		
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CDP	В	2001[B]	-	21,26,26	2.45	5 (23%)	28,40,40	2.10	5 (17%)
2	CDP	A	1001[A]	-	21,26,26	2.56	4 (19%)	28,40,40	2.17	5 (17%)
2	CDP	A	1001[B]	-	21,26,26	2.58	5 (23%)	28,40,40	2.10	5 (17%)
2	CDP	В	2001[A]	-	21,26,26	2.55	5 (23%)	28,40,40	2.12	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CDP	В	2001[B]	-	-	2/14/32/32	0/2/2/2
2	CDP	A	1001[A]	-	-	3/14/32/32	0/2/2/2
2	CDP	A	1001[B]	_	-	1/14/32/32	0/2/2/2
2	CDP	В	2001[A]	-	-	1/14/32/32	0/2/2/2

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	A	1001[B]	CDP	C6-N1	10.17	1.48	1.35
2	A	1001[A]	CDP	C6-N1	10.15	1.48	1.35
2	В	2001[A]	CDP	C6-N1	10.03	1.48	1.35
2	В	2001[B]	CDP	C6-N1	9.52	1.47	1.35
2	В	2001[A]	CDP	O4'-C4'	-2.66	1.39	1.45
2	В	2001[A]	CDP	PB-O1B	2.57	1.58	1.50
2	A	1001[B]	CDP	PB-O1B	2.56	1.58	1.50
2	A	1001[B]	CDP	O4'-C4'	-2.56	1.39	1.45
2	В	2001[B]	CDP	O4'-C4'	-2.54	1.39	1.45
2	A	1001[A]	CDP	O4'-C4'	-2.53	1.39	1.45
2	A	1001[A]	CDP	PB-O1B	2.45	1.58	1.50
2	В	2001[A]	CDP	C6-C5	-2.42	1.32	1.38
2	В	2001[B]	CDP	C6-C5	-2.41	1.32	1.38
2	A	1001[B]	CDP	C6-C5	-2.39	1.32	1.38
2	A	1001[A]	CDP	C6-C5	-2.34	1.32	1.38
2	В	2001[A]	CDP	PA-O1A	2.18	1.58	1.50
2	В	2001[B]	CDP	PB-O1B	2.17	1.57	1.50
2	A	1001[B]	CDP	PA-O1A	2.17	1.58	1.50
2	В	2001[B]	CDP	PA-O1A	2.09	1.58	1.50

All (20) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	1001[A]	CDP	C2-N3-C4	6.32	122.75	116.34
2	A	1001[B]	CDP	C2-N3-C4	6.31	122.73	116.34
2	В	2001[A]	CDP	C2-N3-C4	6.26	122.68	116.34
2	В	2001[B]	CDP	C2-N3-C4	5.99	122.41	116.34
2	A	1001[A]	CDP	PA-O3A-PB	-5.70	113.28	132.83
2	В	2001[A]	CDP	PA-O3A-PB	-5.46	114.10	132.83
2	В	2001[B]	CDP	PA-O3A-PB	-5.10	115.32	132.83
2	A	1001[B]	CDP	PA-O3A-PB	-5.06	115.47	132.83
2	A	1001[A]	CDP	C6-N1-C2	-4.63	113.84	121.20
2	В	2001[B]	CDP	C6-N1-C2	-4.59	113.90	121.20
2	A	1001[B]	CDP	C6-N1-C2	-4.58	113.92	121.20
2	В	2001[A]	CDP	C6-N1-C2	-4.43	114.16	121.20
2	В	2001[B]	CDP	C5-C4-N3	-4.18	116.90	121.72
2	A	1001[B]	CDP	C5-C4-N3	-4.07	117.02	121.72
2	A	1001[A]	CDP	C5-C4-N3	-4.06	117.04	121.72
2	В	2001[A]	CDP	C5-C4-N3	-4.03	117.07	121.72
2	В	2001[B]	CDP	N4-C4-N3	3.04	121.29	116.49
2	A	1001[B]	CDP	N4-C4-N3	2.91	121.08	116.49
2	A	1001[A]	CDP	N4-C4-N3	2.89	121.05	116.49
2	В	2001[A]	CDP	N4-C4-N3	2.73	120.80	116.49

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001[A]	CDP	C5'-O5'-PA-O1A
2	В	2001[B]	CDP	PB-O3A-PA-O1A
2	A	1001[A]	CDP	C5'-O5'-PA-O3A
2	A	1001[A]	CDP	C5'-O5'-PA-O2A
2	В	2001[B]	CDP	PB-O3A-PA-O2A
2	A	1001[B]	CDP	C5'-O5'-PA-O1A
2	В	2001[A]	CDP	C5'-O5'-PA-O1A


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring





in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

