



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 25, 2020 – 02:59 pm BST

PDB ID : 6EYS  
Title : Crystal structure of the periplasmic pyoverdine maturation protein PvdP  
Authors : Poppe, J.; Blankenfeldt, W.  
Deposited on : 2017-11-13  
Resolution : 2.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	536	<p>3% 85% 13%</p>
1	B	536	<p>2% 87% 11%</p>
1	C	536	<p>3% 86% 12%</p>
1	D	536	<p>3% 85% 12%</p>

## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 30248 atoms, of which 14481 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PvdP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	B	478	7548	2472	3679	708	682	7	0	4	0
1	A	469	7379	2426	3590	686	670	7	0	2	0
1	C	470	7404	2427	3613	692	664	8	0	2	0
1	D	472	7404	2432	3599	695	671	7	0	4	0

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	9	HIS	-	expression tag	UNP A0A0H2ZBG1
B	10	HIS	-	expression tag	UNP A0A0H2ZBG1
B	11	HIS	-	expression tag	UNP A0A0H2ZBG1
B	12	HIS	-	expression tag	UNP A0A0H2ZBG1
B	13	HIS	-	expression tag	UNP A0A0H2ZBG1
B	14	HIS	-	expression tag	UNP A0A0H2ZBG1
B	15	SER	-	expression tag	UNP A0A0H2ZBG1
B	16	SER	-	expression tag	UNP A0A0H2ZBG1
B	17	GLY	-	expression tag	UNP A0A0H2ZBG1
B	18	LEU	-	expression tag	UNP A0A0H2ZBG1
B	19	GLU	-	expression tag	UNP A0A0H2ZBG1
B	20	VAL	-	expression tag	UNP A0A0H2ZBG1
B	21	LEU	-	expression tag	UNP A0A0H2ZBG1
B	22	PHE	-	expression tag	UNP A0A0H2ZBG1
B	23	GLN	-	expression tag	UNP A0A0H2ZBG1
B	24	GLY	-	expression tag	UNP A0A0H2ZBG1
B	25	THR	-	expression tag	UNP A0A0H2ZBG1
A	9	HIS	-	expression tag	UNP A0A0H2ZBG1
A	10	HIS	-	expression tag	UNP A0A0H2ZBG1
A	11	HIS	-	expression tag	UNP A0A0H2ZBG1
A	12	HIS	-	expression tag	UNP A0A0H2ZBG1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	HIS	-	expression tag	UNP A0A0H2ZBG1
A	14	HIS	-	expression tag	UNP A0A0H2ZBG1
A	15	SER	-	expression tag	UNP A0A0H2ZBG1
A	16	SER	-	expression tag	UNP A0A0H2ZBG1
A	17	GLY	-	expression tag	UNP A0A0H2ZBG1
A	18	LEU	-	expression tag	UNP A0A0H2ZBG1
A	19	GLU	-	expression tag	UNP A0A0H2ZBG1
A	20	VAL	-	expression tag	UNP A0A0H2ZBG1
A	21	LEU	-	expression tag	UNP A0A0H2ZBG1
A	22	PHE	-	expression tag	UNP A0A0H2ZBG1
A	23	GLN	-	expression tag	UNP A0A0H2ZBG1
A	24	GLY	-	expression tag	UNP A0A0H2ZBG1
A	25	THR	-	expression tag	UNP A0A0H2ZBG1
C	9	HIS	-	expression tag	UNP A0A0H2ZBG1
C	10	HIS	-	expression tag	UNP A0A0H2ZBG1
C	11	HIS	-	expression tag	UNP A0A0H2ZBG1
C	12	HIS	-	expression tag	UNP A0A0H2ZBG1
C	13	HIS	-	expression tag	UNP A0A0H2ZBG1
C	14	HIS	-	expression tag	UNP A0A0H2ZBG1
C	15	SER	-	expression tag	UNP A0A0H2ZBG1
C	16	SER	-	expression tag	UNP A0A0H2ZBG1
C	17	GLY	-	expression tag	UNP A0A0H2ZBG1
C	18	LEU	-	expression tag	UNP A0A0H2ZBG1
C	19	GLU	-	expression tag	UNP A0A0H2ZBG1
C	20	VAL	-	expression tag	UNP A0A0H2ZBG1
C	21	LEU	-	expression tag	UNP A0A0H2ZBG1
C	22	PHE	-	expression tag	UNP A0A0H2ZBG1
C	23	GLN	-	expression tag	UNP A0A0H2ZBG1
C	24	GLY	-	expression tag	UNP A0A0H2ZBG1
C	25	THR	-	expression tag	UNP A0A0H2ZBG1
D	9	HIS	-	expression tag	UNP A0A0H2ZBG1
D	10	HIS	-	expression tag	UNP A0A0H2ZBG1
D	11	HIS	-	expression tag	UNP A0A0H2ZBG1
D	12	HIS	-	expression tag	UNP A0A0H2ZBG1
D	13	HIS	-	expression tag	UNP A0A0H2ZBG1
D	14	HIS	-	expression tag	UNP A0A0H2ZBG1
D	15	SER	-	expression tag	UNP A0A0H2ZBG1
D	16	SER	-	expression tag	UNP A0A0H2ZBG1
D	17	GLY	-	expression tag	UNP A0A0H2ZBG1
D	18	LEU	-	expression tag	UNP A0A0H2ZBG1
D	19	GLU	-	expression tag	UNP A0A0H2ZBG1
D	20	VAL	-	expression tag	UNP A0A0H2ZBG1

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
D	21	LEU	-	expression tag	UNP A0A0H2ZBG1
D	22	PHE	-	expression tag	UNP A0A0H2ZBG1
D	23	GLN	-	expression tag	UNP A0A0H2ZBG1
D	24	GLY	-	expression tag	UNP A0A0H2ZBG1
D	25	THR	-	expression tag	UNP A0A0H2ZBG1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	157	Total O 157 157	0	0
2	A	124	Total O 124 124	0	0
2	C	131	Total O 131 131	0	0
2	D	101	Total O 101 101	0	0



- Molecule 1: PvdP



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.35Å 107.79Å 107.94Å 90.00° 99.97° 90.00°	Depositor
Resolution (Å)	45.61 – 2.09 45.61 – 2.09	Depositor EDS
% Data completeness (in resolution range)	93.2 (45.61-2.09) 66.2 (45.61-2.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.18 (at 2.08Å)	Xtrriage
Refinement program	PHENIX (dev_2875)	Depositor
R, $R_{free}$	0.214 , 0.241 0.214 , 0.241	Depositor DCC
$R_{free}$ test set	1999 reflections (2.33%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.1	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 33.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	30248	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/3921	0.41	0/5336
1	B	0.25	0/4015	0.41	0/5462
1	C	0.25	0/3925	0.41	0/5339
1	D	0.25	0/3945	0.41	0/5372
All	All	0.25	0/15806	0.41	0/21509

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3789	3590	3577	11	0
1	B	3869	3679	3651	5	0
1	C	3791	3613	3602	5	0
1	D	3805	3599	3575	9	0
2	A	124	0	0	3	0
2	B	157	0	0	2	0
2	C	131	0	0	1	0
2	D	101	0	0	1	0
All	All	15767	14481	14405	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:HIS:NE2	2:B:601:HOH:O	2.24	0.70
1:C:271:HIS:NE2	2:C:603:HOH:O	2.30	0.64
1:A:111:ARG:NH1	1:A:132:ASP:OD1	2.32	0.63
1:A:111:ARG:HH12	1:A:132:ASP:CG	2.02	0.61
1:A:111:ARG:HG3	1:A:112:PHE:CD2	2.39	0.58

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/536 (86%)	451 (97%)	12 (3%)	0	100	100
1	B	474/536 (88%)	459 (97%)	15 (3%)	0	100	100
1	C	462/536 (86%)	447 (97%)	15 (3%)	0	100	100
1	D	468/536 (87%)	451 (96%)	17 (4%)	0	100	100
All	All	1867/2144 (87%)	1808 (97%)	59 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/454 (85%)	383 (100%)	2 (0%)	88	92
1	B	395/454 (87%)	393 (100%)	2 (0%)	88	92
1	C	385/454 (85%)	383 (100%)	2 (0%)	88	92
1	D	387/454 (85%)	384 (99%)	3 (1%)	81	86
All	All	1552/1816 (86%)	1543 (99%)	9 (1%)	86	90

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	129	ARG
1	D	212	HIS
1	D	129	ARG
1	A	129	ARG
1	C	181	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	469/536 (87%)	0.38	18 (3%) 40 46	27, 47, 73, 115	0
1	B	478/536 (89%)	0.19	11 (2%) 60 65	26, 41, 73, 91	1 (0%)
1	C	470/536 (87%)	0.28	14 (2%) 50 56	27, 45, 76, 98	1 (0%)
1	D	472/536 (88%)	0.39	17 (3%) 42 49	27, 48, 77, 108	3 (0%)
All	All	1889/2144 (88%)	0.31	60 (3%) 47 54	26, 45, 76, 115	5 (0%)

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	390	SER	5.1
1	B	36	PRO	4.8
1	C	36	PRO	4.5
1	B	396	MET	4.1
1	B	387	ARG	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.