



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 12:12 pm BST

PDB ID : 2F0Y  
Title : Crystal Structure Of Human Protein Farnesyltransferase Complexed With Farnesyl Diphosphate and hydantoin derivative  
Authors : Kim, K.H.; Lee, J.; Kim, J.  
Deposited on : 2005-11-14  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	379	
2	B	437	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	314	2676	1707	466	498	5	0	0	0


- Molecule 2 is a protein called Protein farnesyltransferase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	395	3107	1984	533	568	22	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	B	1	1	1	0	0

- Molecule 4 is FARNESYL DIPHOSPHATE (three-letter code: FPP) (formula: C<sub>15</sub>H<sub>28</sub>O<sub>7</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
4	B	1	24	15	7	2	0	0

- Molecule 5 is 3-({3-[3-(1H-IMIDAZOL-1-YL)PROPYL]-5-METHYL-5-(1-NAPHTHYL)-2,4-DIOXOIMIDAZOLIDIN-1-YL}METHYL)BENZONITRILE (three-letter code: 3MN) (formula: C<sub>28</sub>H<sub>25</sub>N<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	B	1	35	28	5	2	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	201	201	201	0	0
6	B	208	208	208	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit



- Molecule 2: Protein farnesyltransferase beta subunit



PHE	G349	H362	C366	G374	H375	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY		
GLU	L350	L351	L367	L369	L370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
LEU	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
LYS	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
ASP	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
GLU	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
THR	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
ALA	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
PRO	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
ALA	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
THR	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY
ASP	L351	L351	S368	S369	S370	S371	Q374	F376	G377	S378	G379	A380	M381	L382	H383	E384	V385	V386	L387	G388	V389	P390	E391	L394	Q395	H398	P399	V400	Y401	N402	I403	G404	K407	V408	A411	T412	T413	Y414	F415	L416	Q417	K418	P419	V420	P421	GLY

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	171.89Å 171.89Å 71.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.70	Depositor
% Data completeness (in resolution range)	(Not available) (30.00-2.70)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.212 , 0.258	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3MN, ZN, FPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/2743	0.61	0/3725
2	B	0.43	0/3188	0.63	0/4328
All	All	0.44	0/5931	0.62	0/8053

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2676	0	2594	150	0
2	B	3107	0	3038	185	0
3	B	1	0	0	0	0
4	B	24	0	25	2	0
5	B	35	0	25	3	0
6	A	201	0	0	20	0
6	B	208	0	0	21	0
All	All	6252	0	5682	329	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.



The worst 5 of 329 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:127:CYS:HB3	2:B:171:ILE:HD11	1.54	0.89
1:A:75:ILE:HD11	1:A:115:ARG:HH22	1.38	0.87
2:B:389:VAL:HG23	2:B:391:GLU:H	1.40	0.86
1:A:303:GLN:HA	1:A:303:GLN:HE21	1.42	0.84
1:A:294:ASN:HB3	1:A:298:GLN:HE21	1.43	0.81

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	312/379 (82%)	257 (82%)	43 (14%)	12 (4%)	<b>3</b>   <b>7</b>
2	B	391/437 (90%)	351 (90%)	30 (8%)	10 (3%)	<b>5</b>   <b>13</b>
All	All	703/816 (86%)	608 (86%)	73 (10%)	22 (3%)	<b>4</b>   <b>9</b>

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	307	SER
2	B	74	GLN
2	B	378	SER
1	A	71	GLU
1	A	365	SER

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/341 (86%)	281 (96%)	12 (4%)	30	59
2	B	332/370 (90%)	316 (95%)	16 (5%)	25	53
All	All	625/711 (88%)	597 (96%)	28 (4%)	27	55

5 of 28 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	29	GLU
2	B	88	GLN
2	B	387	LEU
2	B	53	GLU
2	B	62	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	335	ASN
1	A	364	GLN
2	B	327	HIS
1	A	326	GLN
2	B	30	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	FPP	B	1001	-	21,23,23	0.86	0	27,31,31	1.53	5 (18%)
5	3MN	B	963	3	37,39,39	2.03	9 (24%)	47,56,56	1.98	9 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FPP	B	1001	-	-	8/25/25/25	-
5	3MN	B	963	3	-	0/18/42/42	0/5/5/5

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	963	3MN	C26-C25	5.41	1.44	1.37
5	B	963	3MN	C21-N4	4.99	1.46	1.37
5	B	963	3MN	C14-C25	4.31	1.56	1.53
5	B	963	3MN	C14-C4	3.34	1.57	1.53
5	B	963	3MN	C25-C24	2.61	1.49	1.43

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	963	3MN	C14-C25-C24	8.34	125.34	120.46
5	B	963	3MN	C19-C11-N2	4.30	124.83	114.55
5	B	963	3MN	N2-C18-N1	3.58	109.84	107.34
4	B	1001	FPP	C10-C8-C9	-3.47	109.44	115.27
4	B	1001	FPP	C5-C6-C7	3.45	123.23	111.88

There are no chirality outliers.

5 of 8 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1001	FPP	C1-O1-PA-O1A
4	B	1001	FPP	C1-O1-PA-O2A
4	B	1001	FPP	C10-C8-C9-C11
4	B	1001	FPP	C7-C8-C9-C11
4	B	1001	FPP	C4-C3-C5-C6


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1001	FPP	2	0
5	B	963	3MN	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.