

wwPDB X-ray Structure Validation Summary Report (i)

Oct 24, 2023 – 08:20 PM EDT

PDB ID	:	3F5O
Title	:	Crystal Structure of hTHEM2(undecan-2-one-CoA) complex
Authors	:	Xu, H.; Gong, W.
Deposited on	:	2008-11-04
Resolution	:	1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Chain	Length	Quality of chain	
А	148	% • 84%	9% 7%
В		3%	13% • 8%
		%	19% • 7%
		3%	14% 7%
		2%	14% •• 7%
		A 148 B 148 C 148 D 148	A 148 % B 148 3% C 148 78% D 148 71% 2% 79%



Mol	Chain	Length	Quality of chain		
1	F	148	78%	14%	• 7%
1	G	148	81%	11%	8%
1	Н	148	3% 76%	16%	•• 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	COA	А	150	Х	-	-	-



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	1 1	190	Total	С	Ν	0	S	0	3	0
	А	138	1039	651	180	198	10	0	0	0
1	В	136	Total	С	Ν	0	S	0	4	0
	D	150	1025	643	182	192	8	0	4	0
1	С	137	Total	С	Ν	0	S	0	4	0
	U	157	1036	650	182	195	9	0	4	0
1	D	137	Total	С	Ν	0	S	0	5	0
1	D	157	1025	643	177	195	10			0
1	Е	138	Total	С	Ν	Ο	\mathbf{S}	0	1	0
1	Ľ	150	1021	639	178	196	8	0	T	0
1	F	138	Total	С	Ν	Ο	\mathbf{S}	0	1	0
	Ľ	100	1024	639	181	196	8	0	T	0
1	G	136	Total	С	Ν	0	\mathbf{S}	0	3	0
	G	130	1017	642	177	189	9	0	5	0
1	Н	138	Total	С	Ν	Ο	\mathbf{S}	0	0 5	0
	11	100	1036	651	181	194	10	0		0

• Molecule 1 is a protein called Thioesterase superfamily member 2.

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	141	LEU	-	expression tag	UNP Q9NPJ3
А	142	GLU	-	expression tag	UNP Q9NPJ3
А	143	HIS	-	expression tag	UNP Q9NPJ3
А	144	HIS	-	expression tag	UNP Q9NPJ3
А	145	HIS	-	expression tag	UNP Q9NPJ3
А	146	HIS	-	expression tag	UNP Q9NPJ3
А	147	HIS	-	expression tag	UNP Q9NPJ3
А	148	HIS	-	expression tag	UNP Q9NPJ3
В	141	LEU	-	expression tag	UNP Q9NPJ3
В	142	GLU	-	expression tag	UNP Q9NPJ3
В	143	HIS	-	expression tag	UNP Q9NPJ3
В	144	HIS	-	expression tag	UNP Q9NPJ3
В	145	HIS	-	expression tag	UNP Q9NPJ3



	Continued from previous page								
Chain	Residue	Modelled	Actual	Comment	Reference				
B	146	HIS	-	expression tag	UNP Q9NPJ3				
B	147	HIS	-	expression tag	UNP Q9NPJ3				
B	148	HIS	-	expression tag	UNP Q9NPJ3				
C	141	LEU	-	expression tag	UNP Q9NPJ3				
С	142	GLU	-	expression tag	UNP Q9NPJ3				
С	143	HIS	-	expression tag	UNP Q9NPJ3				
С	144	HIS	-	expression tag	UNP Q9NPJ3				
С	145	HIS	-	expression tag	UNP Q9NPJ3				
С	146	HIS	-	expression tag	UNP Q9NPJ3				
С	147	HIS	-	expression tag	UNP Q9NPJ3				
С	148	HIS	-	expression tag	UNP Q9NPJ3				
D	141	LEU	-	expression tag	UNP Q9NPJ3				
D	142	GLU	-	expression tag	UNP Q9NPJ3				
D	143	HIS	-	expression tag	UNP Q9NPJ3				
D	144	HIS	-	expression tag	UNP Q9NPJ3				
D	145	HIS	-	expression tag	UNP Q9NPJ3				
D	146	HIS	-	expression tag	UNP Q9NPJ3				
D	147	HIS	-	expression tag	UNP Q9NPJ3				
D	148	HIS	-	expression tag	UNP Q9NPJ3				
E	141	LEU	-	expression tag	UNP Q9NPJ3				
E	142	GLU	-	expression tag	UNP Q9NPJ3				
E	143	HIS	-	expression tag	UNP Q9NPJ3				
E	144	HIS	-	expression tag	UNP Q9NPJ3				
Е	145	HIS	-	expression tag	UNP Q9NPJ3				
E	146	HIS	-	expression tag	UNP Q9NPJ3				
Е	147	HIS	-	expression tag	UNP Q9NPJ3				
E	148	HIS	-	expression tag	UNP Q9NPJ3				
F	141	LEU	-	expression tag	UNP Q9NPJ3				
F	142	GLU	-	expression tag	UNP Q9NPJ3				
F	143	HIS	-	expression tag	UNP Q9NPJ3				
F	144	HIS	-	expression tag	UNP Q9NPJ3				
F	145	HIS	-	expression tag	UNP Q9NPJ3				
F	146	HIS	-	expression tag	UNP Q9NPJ3				
F	147	HIS	-	expression tag	UNP Q9NPJ3				
F	148	HIS	-	expression tag	UNP Q9NPJ3				
G	141	LEU	-	expression tag	UNP Q9NPJ3				
G	142	GLU	-	expression tag	UNP Q9NPJ3				
G	143	HIS	-	expression tag	UNP Q9NPJ3				
G	144	HIS	-	expression tag	UNP Q9NPJ3				
G	145	HIS	-	expression tag	UNP Q9NPJ3				
G	146	HIS	-	expression tag	UNP Q9NPJ3				
G	147	HIS	-	expression tag	UNP Q9NPJ3				
					-				




Continu	Continuea from previous page								
Chain	Residue	Modelled	Actual	Comment	Reference				
G	148	HIS	-	expression tag	UNP Q9NPJ3				
Н	141	LEU	-	expression tag	UNP Q9NPJ3				
H	142	GLU	-	expression tag	UNP Q9NPJ3				
Н	143	HIS	-	expression tag	UNP Q9NPJ3				
Н	144	HIS	-	expression tag	UNP Q9NPJ3				
Н	145	HIS	-	expression tag	UNP Q9NPJ3				
Н	146	HIS	-	expression tag	UNP Q9NPJ3				
Н	147	HIS	-	expression tag	UNP Q9NPJ3				
Н	148	HIS	-	expression tag	UNP Q9NPJ3				

• Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total Cl 1 1	0	0
2	С	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0
2	Ε	1	Total Cl 1 1	0	0
2	F	1	Total Cl 1 1	0	0
2	G	1	Total Cl 1 1	0	0
2	Н	1	Total Cl 1 1	0	0






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C O 12 11 1	0	0
3	В	1	Total C O 12 11 1	0	0
3	С	1	Total C O 12 11 1	0	0
3	D	1	Total C O 12 11 1	0	0
3	Е	1	Total C O 12 11 1	0	0
3	F	1	Total C O 12 11 1	0	0
3	G	1	Total C O 12 11 1	0	0
3	Н	1	Total C O 12 11 1	0	0

• Molecule 4 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).






Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
4	А	1	Total	С	Ν	Ο	Р	S	0	0	
4	A	1	38	16	2	16	3	1	0	0	
4	В	1	Total	С	Ν	Ο	Р	S	0	0	
4	D	1	48	21	7	16	3	1	0	0	
4	С	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
4		1	48	21	7	16	3	1	0	0	
4	D	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
4	D	1	48	21	7	16	3	1	0		
4	Е	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
4	Ľ	T	48	21	7	16	3	1	0	0	
4	F	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
4	T,	1	48	21	7	16	3	1	0	0	
4	G	1	Total	С	Ν	Ο	Р	\mathbf{S}	0	0	
-	0	1	48	21	7	16	3	1	0	U	
4	Н	1	Total	С	Ν	Ο	Р	S	0	0	
±	11	1	38	16	2	16	3	1	U	0	

• Molecule 5 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $C_{12}H_{26}O_7$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 15 10 5	0	0
5	С	1	Total C O 15 10 5	0	0
5	Е	1	Total C O 15 10 5	0	0
5	G	1	Total C O 15 10 5	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	111	Total O 111 111	0	0
6	В	95	Total O 95 95	0	0
6	С	127	Total O 127 127	0	0
6	D	92	Total O 92 92	0	0
6	Е	118	Total O 118 118	0	0
6	F	107	Total O 107 107	0	0
6	G	102	Total O 102 102	0	0
6	Н	98	Total O 98 98	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Thioesterase superfamily member 2





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	91.03Å 110.63Å 119.82Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.73 - 1.70	Depositor
Resolution (A)	20.73 - 1.70	EDS
% Data completeness	98.8 (20.73-1.70)	Depositor
(in resolution range)	98.9(20.73-1.70)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.82 (at 1.69 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
B B.	0.199 , 0.237	Depositor
R, R_{free}	0.197 , 0.236	DCC
R_{free} test set	6645 reflections $(5.01%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	26.4	Xtriage
Anisotropy	0.368	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36 , 54.5	EDS
L-test for twinning ²	$ < L >=0.52, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9600	wwPDB-VP
Average B, all atoms $(Å^2)$	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 45.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3689e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, P6G, UOC, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	ond angles
10101	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.63	0/1055	0.68	0/1422
1	В	0.63	0/1048	0.75	0/1415
1	С	0.61	0/1059	0.71	0/1427
1	D	0.63	0/1050	0.71	0/1419
1	Е	0.66	0/1034	0.75	2/1397~(0.1%)
1	F	0.64	0/1037	0.76	2/1401~(0.1%)
1	G	0.62	0/1037	0.69	0/1399
1	Н	0.58	0/1062	0.74	1/1434~(0.1%)
All	All	0.62	0/8382	0.72	5/11314~(0.0%)

There are no bond length outliers.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	Ε	116	THR	N-CA-CB	-5.93	99.04	110.30
1	Н	116	THR	N-CA-CB	-5.81	99.26	110.30
1	Е	116	THR	OG1-CB-CG2	5.72	123.17	110.00
1	F	121	THR	OG1-CB-CG2	5.54	122.75	110.00
1	F	121	THR	N-CA-CB	-5.23	100.37	110.30

All (5) bond angle outliers are listed below:

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1039	0	1090	19	0
1	В	1025	0	1074	19	0
1	С	1036	0	1092	32	0
1	D	1025	0	1068	23	0
1	Е	1021	0	1062	32	0
1	F	1024	0	1073	18	0
1	G	1017	0	1058	15	0
1	Н	1036	0	1087	37	0
2	А	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	Е	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	Н	1	0	0	0	0
3	А	12	0	20	5	0
3	В	12	0	21	0	0
3	С	12	0	21	1	0
3	D	12	0	21	1	0
3	Е	12	0	21	2	0
3	F	12	0	21	0	0
3	G	12	0	21	1	0
3	Н	12	0	21	2	0
4	А	38	0	26	6	0
4	В	48	0	31	1	0
4	С	48	0	31	0	0
4	D	48	0	31	0	0
4	Е	48	0	31	1	0
4	F	48	0	31	3	0
4	G	48	0	31	5	0
4	Н	38	0	26	3	0
5	В	15	0	16	0	0
5	С	15	0	16	0	0
5	Е	15	0	16	1	0
5	G	15	0	16	1	0
6	А	111	0	0	1	0
6	В	95	0	0	3	0
6	С	127	0	0	0	0
6	D	92	0	0	1	0
6	Е	118	0	0	5	0
6	F	107	0	0	1	0
6	G	102	0	0	1	0
6	Н	98	0	0	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9600	0	9073	184	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 184 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:56:HIS:HD2	1:C:58:GLY:H	1.08	0.99
1:D:56:HIS:HD2	1:D:58:GLY:H	1.08	0.97
1:C:27:LYS:HE3	1:C:45:GLU:HG3	1.45	0.97
1:A:133:ARG:HE	1:D:131:GLN:HE21	1.10	0.96
1:F:133:ARG:HE	1:G:131:GLN:HE21	1.14	0.96

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	139/148~(94%)	136~(98%)	3~(2%)	0	100	100
1	В	138/148~(93%)	134~(97%)	4 (3%)	0	100	100
1	С	139/148~(94%)	137~(99%)	2(1%)	0	100	100
1	D	140/148~(95%)	139~(99%)	1 (1%)	0	100	100
1	Ε	137/148~(93%)	133~(97%)	4(3%)	0	100	100
1	F	137/148~(93%)	135~(98%)	2(2%)	0	100	100
1	G	136/148~(92%)	134~(98%)	2(2%)	0	100	100
1	Н	141/148~(95%)	140 (99%)	1 (1%)	0	100	100
All	All	1107/1184~(94%)	1088 (98%)	19 (2%)	0	100	100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	114/123~(93%)	113~(99%)	1 (1%)	78 70
1	В	111/123~(90%)	107~(96%)	4 (4%)	35 16
1	С	115/123~(94%)	108~(94%)	7~(6%)	18 5
1	D	112/123~(91%)	111 (99%)	1 (1%)	78 70
1	Ε	111/123~(90%)	105~(95%)	6~(5%)	22 7
1	\mathbf{F}	113/123~(92%)	111 (98%)	2(2%)	59 43
1	G	108/123~(88%)	108 (100%)	0	100 100
1	Н	114/123~(93%)	110~(96%)	4 (4%)	36 17
All	All	898/984~(91%)	873~(97%)	25~(3%)	43 25

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	13	LYS
1	Ε	116	THR
1	Н	116	THR
1	Е	108	LYS
1	Е	121	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	105	HIS
1	F	131	GLN
1	F	56	HIS
1	G	56	HIS
1	С	56	HIS



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 27 ligands modelled in this entry, 7 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	gths	B	ond ang	les
MOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	UOC	Н	149	4	11,11,11	0.62	0	11,11,11	0.72	0
3	UOC	Е	149	4	11,11,11	0.39	0	11,11,11	0.62	0
3	UOC	D	149	4	11,11,11	0.60	0	11,11,11	0.94	0
5	P6G	С	1001	-	$14,\!14,\!18$	0.63	0	$13,\!13,\!17$	0.62	0
3	UOC	F	149	4	11,11,11	0.47	0	11,11,11	1.27	2 (18%)
4	COA	В	150	3	41,50,50	1.55	9 (21%)	52,75,75	1.31	5 (9%)
4	COA	G	150	3	$41,\!50,\!50$	1.62	6 (14%)	52,75,75	1.39	7 (13%)
3	UOC	G	149	4	11,11,11	0.54	0	11,11,11	1.08	1 (9%)
4	COA	Е	150	3	41,50,50	1.92	9 (21%)	52,75,75	1.29	5 (9%)
4	COA	А	150	3	32,38,50	1.71	10 (31%)	$45,\!56,\!75$	1.69	9 (20%)
3	UOC	В	149	4	11,11,11	0.50	0	11,11,11	0.85	0
3	UOC	А	149	4	11,11,11	0.73	0	11,11,11	1.06	1 (9%)
5	P6G	В	1001	-	14,14,18	0.71	0	13,13,17	0.40	0
5	P6G	G	1001	-	14,14,18	0.65	0	13,13,17	0.74	0
5	P6G	Е	1001	-	14,14,18	0.78	0	$13,\!13,\!17$	0.56	0
4	COA	F	150	3	$41,\!50,\!50$	1.53	8 (19%)	52,75,75	1.08	3 (5%)



Mol	Turne	Chain	Res Link		B	Bond lengths			Bond angles		
10101	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
4	COA	D	150	3	41,50,50	1.95	10 (24%)	52,75,75	1.51	8 (15%)	
4	COA	С	150	3	41,50,50	1.76	9 (21%)	52,75,75	1.40	9 (17%)	
4	COA	Н	150	3	32,38,50	1.60	7 (21%)	45,56,75	1.19	5 (11%)	
3	UOC	С	149	4	11,11,11	0.60	0	11,11,11	0.61	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UOC	Н	149	4	-	7/9/9/9	-
3	UOC	D	149	4	-	4/9/9/9	-
3	UOC	Е	149	4	-	4/9/9/9	-
5	P6G	С	1001	-	-	7/12/12/16	-
3	UOC	F	149	4	-	4/9/9/9	-
4	COA	В	150	3	-	4/44/64/64	0/3/3/3
4	COA	G	150	3	-	1/44/64/64	0/3/3/3
3	UOC	G	149	4	-	4/9/9/9	-
4	COA	Е	150	3	-	2/44/64/64	0/3/3/3
4	COA	А	150	3	1/1/10/13	18/44/57/64	0/1/1/3
3	UOC	В	149	4	-	5/9/9/9	-
3	UOC	А	149	4	-	7/9/9/9	-
5	P6G	В	1001	-	-	7/12/12/16	-
5	P6G	G	1001	-	-	8/12/12/16	-
5	P6G	Е	1001	-	-	7/12/12/16	-
4	COA	F	150	3	-	0/44/64/64	0/3/3/3
4	COA	D	150	3	-	4/44/64/64	0/3/3/3
4	COA	С	150	3	-	5/44/64/64	0/3/3/3
4	COA	Н	150	3	-	5/44/57/64	0/1/1/3
3	UOC	С	149	4	-	2/9/9/9	-

The worst 5 of 68 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
4	D	150	COA	P2A-O4A	-5.09	1.32	1.50
4	G	150	COA	P2A-O4A	-4.66	1.34	1.50



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
4	D	150	COA	P1A-O5B	-4.46	1.41	1.59
4	Е	150	COA	P2A-O4A	-4.31	1.35	1.50
4	С	150	COA	P1A-O2A	-4.27	1.35	1.55

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	D	150	COA	O6A-CCP-CBP	4.67	118.06	110.55
4	Е	150	COA	O4B-C1B-C2B	-4.43	100.45	106.93
4	В	150	COA	N3A-C2A-N1A	-4.41	121.79	128.68
4	G	150	COA	N3A-C2A-N1A	-4.38	121.83	128.68
4	А	150	COA	C6P-C7P-N8P	-4.20	103.42	111.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	А	150	COA	CAP

5 of 105 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	150	COA	C2B-C3B-O3B-P3B
4	А	150	COA	C3B-C4B-C5B-O5B
4	А	150	COA	O4B-C4B-C5B-O5B
4	А	150	COA	CCP-O6A-P2A-O4A
4	А	150	COA	CDP-CBP-CCP-O6A

There are no ring outliers.

14 monomers are involved in 32 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Н	149	UOC	2	0
3	Е	149	UOC	2	0
3	D	149	UOC	1	0
4	В	150	COA	1	0
4	G	150	COA	5	0
3	G	149	UOC	1	0
4	Е	150	COA	1	0
4	А	150	COA	6	0
3	А	149	UOC	5	0
5	G	1001	P6G	1	0



0 0	iraca ji cii	Procee	rao pago.	••	
Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	Е	1001	P6G	1	0
4	F	150	COA	3	0
4	Н	150	COA	3	0
3	С	149	UOC	1	0


Continued from previous page...

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(A^2)$	$\mathbf{Q}{<}0.9$
1	А	138/148~(93%)	-0.01	2 (1%) 75 79	18, 26, 36, 44	4 (2%)
1	В	136/148~(91%)	0.14	5 (3%) 41 46	18, 27, 37, 42	6 (4%)
1	С	137/148~(92%)	-0.11	2 (1%) 73 77	18, 24, 32, 43	6 (4%)
1	D	137/148~(92%)	0.07	4 (2%) 51 56	17, 25, 37, 45	3(2%)
1	Ε	138/148~(93%)	0.01	3 (2%) 62 66	18, 25, 34, 40	6 (4%)
1	F	138/148~(93%)	0.15	3 (2%) 62 66	17, 25, 37, 53	11 (7%)
1	G	136/148~(91%)	0.02	2 (1%) 73 77	18, 26, 33, 38	9~(6%)
1	Н	138/148~(93%)	0.08	5 (3%) 42 47	18, 27, 37, 45	1 (0%)
All	All	1098/1184~(92%)	0.04	26 (2%) 59 63	17, 25, 36, 53	46 (4%)

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	2	THR	10.0
1	Н	2	THR	7.5
1	F	139	GLY	6.4
1	Н	139	GLY	5.9
1	В	25	LEU	4.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
5	P6G	В	1001	15/19	0.32	0.32	72,77,79,79	0
3	UOC	А	149	12/12	0.61	0.31	35,38,45,46	12
3	UOC	G	149	12/12	0.68	0.25	37,45,50,51	0
3	UOC	Н	149	12/12	0.69	0.19	35,47,53,53	0
5	P6G	Е	1001	15/19	0.71	0.24	63,67,70,70	0
3	UOC	В	149	12/12	0.73	0.19	36,46,52,52	0
5	P6G	С	1001	15/19	0.77	0.22	48,52,53,53	0
3	UOC	F	149	12/12	0.77	0.18	37,46,53,53	0
3	UOC	С	149	12/12	0.78	0.17	31,38,42,42	0
4	COA	А	150	38/48	0.79	0.22	33,43,54,55	38
5	P6G	G	1001	15/19	0.79	0.17	63,67,69,69	0
4	COA	В	150	48/48	0.80	0.25	33,43,61,61	33
4	COA	Н	150	38/48	0.80	0.20	33,47,64,64	21
4	COA	F	150	48/48	0.84	0.19	32,43,56,56	27
3	UOC	D	149	12/12	0.86	0.17	29,38,49,51	0
4	COA	G	150	48/48	0.88	0.18	28,45,56,57	30
3	UOC	Е	149	12/12	0.90	0.12	26,37,48,49	0
4	COA	Е	150	48/48	0.91	0.16	20,36,52,53	17
4	COA	С	150	48/48	0.91	0.15	22,34,48,48	17
4	COA	D	150	48/48	0.92	0.14	21,40,58,60	14
2	CL	Е	151	1/1	0.98	0.05	33,33,33,33	0
2	CL	G	151	1/1	0.98	0.03	36,36,36,36	0
2	CL	D	151	1/1	0.99	0.04	27,27,27,27	0
2	CL	А	151	1/1	0.99	0.07	36,36,36,36	0
2	CL	F	151	1/1	0.99	0.04	25,25,25,25	0
2	CL	С	151	1/1	0.99	0.04	25,25,25,25	0
2	CL	Н	151	1/1	0.99	0.03	33,33,33,33	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















































6.5 Other polymers (i)

There are no such residues in this entry.

