



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 26, 2023 – 12:27 PM EDT

PDB ID : 3F6X
Title : c-Src kinase domain in complex with small molecule inhibitor
Authors : Seeliger, M.A.; Statsuk, A.V.; Maly, D.J.; Patrick, P.Z.; Kuriyan, J.; Shokat, K.M.
Deposited on : 2008-11-06
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	 13% 66% 24% • 8%
1	B	286	 9% 67% 25% • 7%
1	C	286	 12% 59% 30% • 7%
1	D	286	 10% 66% 26% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IHH	B	1	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	2124	1364	355	389	16	0	0	0
1	B	267	2137	1372	358	391	16	0	0	0
1	C	265	2133	1373	357	387	16	0	0	0
1	D	267	2142	1375	359	391	17	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523
C	248	GLY	-	expression tag	UNP P00523
C	249	HIS	-	expression tag	UNP P00523
C	250	MET	-	expression tag	UNP P00523
D	248	GLY	-	expression tag	UNP P00523
D	249	HIS	-	expression tag	UNP P00523
D	250	MET	-	expression tag	UNP P00523


- Molecule 2 is [4-({4-[(5-cyclopropyl-1H-pyrazol-3-yl)amino]quinazolin-2-yl}amino)phenyl]acetonitrile (three-letter code: IHH) (formula: C₂₂H₁₉N₇).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C N	0	0
			29	22 7		
2	B	1	Total	C N	0	0
			29	22 7		
2	C	1	Total	C N	0	0
			29	22 7		
2	D	1	Total	C N	0	0
			29	22 7		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	51	Total	O	0	0
			51	51		
3	B	57	Total	O	0	0
			57	57		
3	C	47	Total	O	0	0
			47	47		
3	D	65	Total	O	0	0
			65	65		



● Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.70Å 74.49Å 84.65Å 89.06° 89.97° 78.43°	Depositor
Resolution (Å)	34.99 – 2.35 34.98 – 2.35	Depositor EDS
% Data completeness (in resolution range)	79.8 (34.99-2.35) 80.2 (34.98-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.34Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.234 , 0.266 0.227 , 0.260	Depositor DCC
R_{free} test set	2984 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtrriage
Anisotropy	0.311	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8872	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 91.92 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1880e-08. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IHH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2174	0.58	0/2942
1	B	0.35	0/2187	0.61	0/2961
1	C	0.36	0/2184	0.61	0/2955
1	D	0.36	0/2194	0.60	0/2970
All	All	0.36	0/8739	0.60	0/11828

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2124	0	2104	68	0
1	B	2137	0	2123	57	0
1	C	2133	0	2125	75	0
1	D	2142	0	2124	54	0
2	A	29	0	19	6	0
2	B	29	0	19	9	0
2	C	29	0	19	5	0
2	D	29	0	19	8	0
3	A	51	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	57	0	0	1	0
3	C	47	0	0	1	0
3	D	65	0	0	1	0
All	All	8872	0	8552	275	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 275 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:334:ILE:HD13	1:C:334:ILE:H	1.29	0.96
2:B:1:IHH:H17	2:B:1:IHH:H23	1.52	0.91
2:D:1:IHH:H17	2:D:1:IHH:H23	1.54	0.90
1:C:262:ILE:HD13	1:C:286:TRP:HE1	1.44	0.83
2:B:1:IHH:H17	2:B:1:IHH:N9	1.93	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/286 (90%)	240 (93%)	15 (6%)	3 (1%)	13	11
1	B	261/286 (91%)	246 (94%)	14 (5%)	1 (0%)	34	38
1	C	259/286 (91%)	245 (95%)	11 (4%)	3 (1%)	13	11
1	D	263/286 (92%)	254 (97%)	9 (3%)	0	100	100
All	All	1041/1144 (91%)	985 (95%)	49 (5%)	7 (1%)	22	23

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	404	ASP
1	A	336	ILE
1	A	425	PRO
1	A	404	ASP
1	C	411	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/245 (93%)	217 (96%)	10 (4%)	28	34
1	B	228/245 (93%)	220 (96%)	8 (4%)	36	44
1	C	228/245 (93%)	217 (95%)	11 (5%)	25	30
1	D	229/245 (94%)	218 (95%)	11 (5%)	25	30
All	All	912/980 (93%)	872 (96%)	40 (4%)	28	34

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	526	GLN
1	D	438	ARG
1	D	277	CYS
1	D	351	LYS
1	D	491	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	GLN
1	C	309	GLN
1	C	381	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	IHH	B	1	-	32,33,33	2.03	6 (18%)	42,46,46	2.04	11 (26%)
2	IHH	D	1	-	32,33,33	2.03	6 (18%)	42,46,46	2.06	12 (28%)
2	IHH	C	1	-	32,33,33	2.00	6 (18%)	42,46,46	2.10	12 (28%)
2	IHH	A	1	-	32,33,33	2.00	6 (18%)	42,46,46	2.09	13 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IHH	B	1	-	-	0/12/17/17	0/5/5/5
2	IHH	D	1	-	-	0/12/17/17	0/5/5/5
2	IHH	C	1	-	-	0/12/17/17	0/5/5/5
2	IHH	A	1	-	-	0/12/17/17	0/5/5/5

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	IHH	N26-N25	-7.25	1.23	1.37
2	D	1	IHH	N26-N25	-7.24	1.23	1.37
2	A	1	IHH	N26-N25	-7.18	1.23	1.37
2	C	1	IHH	N26-N25	-7.12	1.23	1.37
2	B	1	IHH	C10-C5	-4.48	1.39	1.44

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	IHH	C10-C5-C6	5.79	119.53	115.88
2	B	1	IHH	C10-C5-C6	5.78	119.52	115.88
2	D	1	IHH	N21-C10-N9	5.65	123.96	118.66
2	C	1	IHH	C10-C5-C6	5.60	119.40	115.88
2	D	1	IHH	C10-C5-C6	5.47	119.32	115.88

There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

4 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	IHH	9	0
2	D	1	IHH	8	0
2	C	1	IHH	5	0
2	A	1	IHH	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/286 (92%)	0.67	38 (14%) 2 4	12, 30, 76, 98	0
1	B	267/286 (93%)	0.53	25 (9%) 8 13	11, 31, 75, 86	0
1	C	265/286 (92%)	0.70	34 (12%) 3 6	12, 31, 78, 90	0
1	D	267/286 (93%)	0.71	30 (11%) 5 8	12, 32, 75, 84	0
All	All	1063/1144 (92%)	0.65	127 (11%) 4 7	11, 31, 77, 98	0

The worst 5 of 127 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	277	CYS	11.6
1	C	299	PRO	10.9
1	D	412	GLU	10.8
1	D	278	PHE	9.2
1	B	300	GLY	8.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	IHH	A	1	29/29	0.90	0.21	30,33,54,55	0
2	IHH	C	1	29/29	0.91	0.18	32,34,54,56	0
2	IHH	B	1	29/29	0.92	0.20	31,34,59,61	0
2	IHH	D	1	29/29	0.93	0.18	28,30,51,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around IHH C 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around IHH B 1:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.