



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 7, 2023 – 09:34 AM EDT

PDB ID : 4F7T  
Title : Crystal Structure of HLA-A\*2402 Complexed with a Newly Identified Peptide from 2009 H1N1 PB1 (498-505)  
Authors : Liu, J.; Zhang, S.; Tan, S.; Yi, Y.; Wu, B.; Zhu, F.; Wang, H.; Qi, J.; Gao, G.F.  
Deposited on : 2012-05-16  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	 6% 87% 12%
1	D	275	 4% 85% 13%
2	B	100	 3% 82% 14%
2	E	100	 10% 81% 17%
3	C	8	 88% 12%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	8	 88% 12%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-24 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	Total	C	N	O	S	0	1	0
			2228	1385	404	429	10			
1	D	274	Total	C	N	O	S	0	0	0
			2222	1382	403	427	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P05534
D	0	MET	-	expression tag	UNP P05534

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	99	Total	C	N	O	S	0	1	0
			835	531	141	160	3			
2	E	99	Total	C	N	O	S	0	0	0
			829	528	140	158	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	expression tag	UNP P61769

- Molecule 3 is a protein called RNA-directed RNA polymerase catalytic subunit.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	8	Total	C	N	O	0	0	0
			70	47	12	11			
3	F	8	Total	C	N	O	0	0	0
			70	47	12	11			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	315	Total 315	O 315	0	0
4	B	116	Total 116	O 116	0	0
4	C	17	Total 17	O 17	0	0
4	D	333	Total 333	O 333	0	0
4	E	89	Total 89	O 89	0	0
4	F	11	Total 11	O 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-24 alpha chain




- Molecule 1: HLA class I histocompatibility antigen, A-24 alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin




- Molecule 3: RNA-directed RNA polymerase catalytic subunit

Chain C:  88% 12%



- Molecule 3: RNA-directed RNA polymerase catalytic subunit

Chain F:  88% 12%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.84Å 66.58Å 70.36Å 97.70° 101.03° 111.84°	Depositor
Resolution (Å)	24.42 – 1.70 24.42 – 1.70	Depositor EDS
% Data completeness (in resolution range)	91.9 (24.42-1.70) 95.9 (24.42-1.70)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.80 (at 1.70Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.5_2)	Depositor
R, $R_{free}$	0.196 , 0.227 0.193 , 0.224	Depositor DCC
$R_{free}$ test set	4527 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtrriage
Anisotropy	0.594	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 56.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7135	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2288	0.51	0/3100
1	D	0.35	0/2282	0.50	1/3092 (0.0%)
2	B	0.28	0/858	0.45	0/1160
2	E	0.26	0/852	0.43	0/1152
3	C	0.32	0/72	0.44	0/94
3	F	0.32	0/72	0.41	0/94
All	All	0.31	0/6424	0.49	1/8692 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	82	LEU	CB-CG-CD1	5.16	119.78	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2086	29	0
1	D	2222	0	2082	31	0
2	B	835	0	798	12	0
2	E	829	0	794	11	0
3	C	70	0	65	2	0
3	F	70	0	65	2	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	315	0	0	6	0
4	B	116	0	0	0	0
4	C	17	0	0	0	0
4	D	333	0	0	6	0
4	E	89	0	0	2	0
4	F	11	0	0	0	0
All	All	7135	0	5890	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:58:LYS:HD2	2:B:58:LYS:H	1.27	0.96
1:A:234:ARG:HE	1:A:242:GLN:HE21	1.33	0.73
1:D:93:HIS:HD2	1:D:119:ASP:OD2	1.73	0.71
2:B:58:LYS:HD2	2:B:58:LYS:N	2.05	0.70
1:D:70:HIS:HD2	3:F:2:TYR:OH	1.75	0.68
2:E:28:SER:OG	4:E:178:HOH:O	2.08	0.68
1:D:234:ARG:HE	1:D:242:GLN:HE21	1.43	0.67
1:D:154:GLU:HG2	4:D:581:HOH:O	1.94	0.67
2:E:15:ALA:HB3	2:E:97:ARG:HG3	1.76	0.66
1:A:70:HIS:HD2	3:C:2:TYR:OH	1.78	0.66
1:D:129:ASP:O	1:D:131:ARG:HG3	1.99	0.62
1:A:151:HIS:O	1:A:154:GLU:HG2	2.01	0.60
1:A:93:HIS:HD2	1:A:119:ASP:OD2	1.83	0.59
1:D:268:LYS:HG2	1:D:269:PRO:HD2	1.83	0.59
1:A:83:ARG:HG3	4:A:430:HOH:O	2.03	0.59
1:A:219:ARG:HG3	1:A:256:ARG:HB3	1.85	0.57
1:A:111:ARG:HG2	1:A:113:TYR:CZ	2.39	0.57
1:A:234:ARG:HE	1:A:242:GLN:NE2	2.03	0.56
1:D:128:GLU:HA	4:D:468:HOH:O	2.04	0.56
1:D:272:LEU:HD12	1:D:272:LEU:N	2.21	0.56
1:A:151:HIS:HB3	1:A:154:GLU:CG	2.37	0.55
1:A:261:VAL:HB	1:A:270:LEU:HB2	1.88	0.55
2:E:51:HIS:HD2	2:E:52:SER:O	1.90	0.54
1:D:199:ALA:HB3	4:D:518:HOH:O	2.08	0.54
2:B:31:HIS:CD2	2:B:32:PRO:HA	2.43	0.53
1:A:219:ARG:HA	1:A:256:ARG:O	2.09	0.52
1:D:192:HIS:HE1	2:E:98:ASP:OD1	1.93	0.51

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:TRP:CD1	1:A:247:VAL:HG13	2.46	0.51
1:D:70:HIS:HE1	1:D:74:ASP:OD2	1.93	0.51
1:A:255:GLN:C	1:A:257:TYR:H	2.13	0.51
2:E:25:CYS:HB2	2:E:39:LEU:HD21	1.91	0.51
1:A:218:GLN:N	4:A:549:HOH:O	2.33	0.51
1:A:70:HIS:CD2	3:C:2:TYR:OH	2.62	0.50
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.47	0.50
1:D:232:GLU:HG3	4:E:178:HOH:O	2.12	0.49
1:A:258:THR:OG1	4:A:549:HOH:O	2.20	0.49
1:D:271:THR:C	1:D:272:LEU:HD12	2.32	0.49
1:D:173:GLU:OE1	1:D:173:GLU:HA	2.12	0.49
2:B:58:LYS:H	2:B:58:LYS:CD	2.11	0.49
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.47	0.48
1:A:96:GLN:OE1	2:B:31:HIS:HE1	1.97	0.48
1:A:214:THR:HG22	1:A:216:THR:HG23	1.96	0.47
1:D:268:LYS:HG2	1:D:269:PRO:CD	2.44	0.47
1:A:188:HIS:HD2	4:A:335:HOH:O	1.98	0.46
1:D:268:LYS:HD3	1:D:269:PRO:O	2.15	0.46
2:B:47:GLU:C	2:B:48:LYS:HG2	2.37	0.45
1:A:218:GLN:HB2	4:A:549:HOH:O	2.16	0.45
1:D:154:GLU:HG3	1:D:157:ARG:HH21	1.82	0.45
1:D:270:LEU:HB3	1:D:272:LEU:HD11	2.00	0.44
1:D:264:GLU:HG2	4:D:305:HOH:O	2.16	0.44
1:D:49:ALA:O	1:D:52:ILE:HG22	2.18	0.44
1:D:121:LYS:CE	2:E:1:ILE:HB	2.48	0.43
2:B:83:ASN:HD22	2:B:83:ASN:HA	1.66	0.43
2:E:97:ARG:O	2:E:97:ARG:HG2	2.17	0.43
1:D:70:HIS:CD2	3:F:2:TYR:OH	2.64	0.43
1:A:230:LEU:HG	1:A:243:LYS:HE3	2.00	0.43
1:A:234:ARG:HH21	1:A:242:GLN:NE2	2.17	0.43
1:D:114:HIS:HD2	4:D:407:HOH:O	2.02	0.43
2:B:21:ASN:HB3	2:B:70:PHE:CE1	2.53	0.42
1:A:151:HIS:HB3	1:A:154:GLU:HG2	2.01	0.42
1:A:114:HIS:HD2	4:A:340:HOH:O	2.02	0.42
2:E:19:LYS:O	2:E:72:PRO:HD2	2.19	0.42
1:D:272:LEU:N	1:D:272:LEU:CD1	2.82	0.42
2:E:21:ASN:HB3	2:E:70:PHE:CE1	2.54	0.42
1:D:234:ARG:HE	1:D:242:GLN:NE2	2.13	0.42
2:E:51:HIS:HA	2:E:65:LEU:O	2.20	0.42
2:B:19:LYS:O	2:B:72:PRO:HD2	2.21	0.41
1:D:186:LYS:N	1:D:186:LYS:HD3	2.35	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:29:GLY:HA2	2:B:61:SER:HB2	2.03	0.41
1:A:70:HIS:HE1	1:A:74:ASP:OD2	2.03	0.41
1:A:111:ARG:HD2	1:A:128:GLU:HG3	2.01	0.41
1:A:255:GLN:C	1:A:257:TYR:N	2.74	0.41
1:D:70:HIS:CE1	1:D:74:ASP:OD2	2.73	0.41
1:D:127:LYS:HD2	1:D:132:SER:OG	2.21	0.41
1:D:146:LYS:HG3	4:D:363:HOH:O	2.21	0.41
1:A:121:LYS:HE3	2:B:1:ILE:HG12	2.03	0.40
1:D:195:SER:C	1:D:197:HIS:H	2.23	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	263 (96%)	9 (3%)	1 (0%)	34	18
1	D	272/275 (99%)	266 (98%)	6 (2%)	0	100	100
2	B	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	E	97/100 (97%)	95 (98%)	2 (2%)	0	100	100
3	C	6/8 (75%)	6 (100%)	0	0	100	100
3	F	6/8 (75%)	6 (100%)	0	0	100	100
All	All	752/766 (98%)	733 (98%)	18 (2%)	1 (0%)	51	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	254	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	224 (97%)	7 (3%)	41	22
1	D	230/231 (100%)	219 (95%)	11 (5%)	25	9
2	B	95/95 (100%)	89 (94%)	6 (6%)	18	5
2	E	94/95 (99%)	90 (96%)	4 (4%)	29	11
3	C	6/6 (100%)	6 (100%)	0	100	100
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	662/664 (100%)	634 (96%)	28 (4%)	30	12

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	HIS
1	A	97	MET
1	A	138	MET
1	A	222	GLU
1	A	226	GLN
1	A	255	GLN
1	A	270	LEU
2	B	19	LYS
2	B	45	ARG
2	B	70	PHE
2	B	74	GLU
2	B	75	LYS
2	B	83	ASN
1	D	17	ARG
1	D	35	ARG
1	D	58	GLU
1	D	70	HIS
1	D	97	MET
1	D	157	ARG
1	D	215	LEU
1	D	249	VAL
1	D	256	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	268	LYS
1	D	272	LEU
2	E	70	PHE
2	E	74	GLU
2	E	75	LYS
2	E	83	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	70	HIS
1	A	86	ASN
1	A	93	HIS
1	A	114	HIS
1	A	115	GLN
1	A	180	GLN
1	A	188	HIS
1	A	218	GLN
1	A	224	GLN
1	A	242	GLN
2	B	31	HIS
2	B	51	HIS
2	B	83	ASN
2	B	89	GLN
1	D	43	GLN
1	D	70	HIS
1	D	93	HIS
1	D	114	HIS
1	D	180	GLN
1	D	192	HIS
1	D	218	GLN
1	D	224	GLN
1	D	242	GLN
1	D	255	GLN
2	E	51	HIS
2	E	83	ASN

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	274/275 (99%)	0.31	16 (5%) 23 25	11, 21, 55, 107	0
1	D	274/275 (99%)	0.19	10 (3%) 42 47	13, 23, 44, 116	0
2	B	99/100 (99%)	0.30	3 (3%) 50 54	14, 26, 46, 73	0
2	E	99/100 (99%)	0.63	10 (10%) 7 8	16, 31, 57, 67	0
3	C	8/8 (100%)	0.45	0 100 100	14, 18, 20, 20	0
3	F	8/8 (100%)	-0.20	0 100 100	17, 19, 22, 33	0
All	All	762/766 (99%)	0.30	39 (5%) 28 31	11, 23, 48, 116	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	225	THR	10.5
1	A	226	GLN	9.7
1	A	227	ASP	7.0
1	D	196	ASP	6.6
1	A	223	ASP	6.1
1	D	194	ILE	5.2
1	A	220	ASP	5.0
1	A	16	GLY	4.9
2	B	48	LYS	4.8
1	A	219	ARG	4.4
1	D	195	SER	4.1
1	D	197	HIS	4.0
2	E	48	LYS	4.0
1	A	228	THR	4.0
1	D	267	PRO	3.7
1	D	225	THR	3.7
1	A	256	ARG	3.4
1	A	221	GLY	3.3
2	E	75	LYS	3.3

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	218	GLN	3.0
2	E	44	GLU	3.0
1	A	252	GLY	2.9
2	E	74	GLU	2.9
1	A	222	GLU	2.9
1	D	268	LYS	2.8
1	A	273	ARG	2.8
2	E	98	ASP	2.8
2	E	45	ARG	2.7
2	E	49	VAL	2.7
2	E	73	THR	2.6
1	A	271	THR	2.6
1	D	226	GLN	2.6
1	A	224	GLN	2.5
2	B	1	ILE	2.5
2	B	47	GLU	2.5
1	D	177	GLU	2.5
2	E	88	SER	2.4
1	D	62	GLU	2.2
2	E	47	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.