



wwPDB X-ray Structure Validation Summary Report

Jun 25, 2024 – 12:24 AM EDT

PDB ID : 6F77
Title : Crystal structure of the prephenate aminotransferase from *Rhizobium meliloti*
Authors : Cobessi, D.; Giustini, C.; Graindorge, M.; Matringe, M.
Deposited on : 2017-12-07
Resolution : 1.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	97%
1	C	400	11% 87% 7% 5%
1	D	400	4% 95% 5%
1	E	400	94% 5%
1	F	400	% 96%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	400	 <p>A horizontal bar chart representing the quality of chain. The bar is 94% green and 5% yellow. A small red square is at the beginning of the bar. The percentage values '94%' and '5%' are printed below the bar.</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 19872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate aminotransferase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	399	Total 3099	C 1980	N 525	O 577	S 17	0	11	0
1	C	380	Total 2878	C 1839	N 477	O 545	S 17	0	3	0
1	D	399	Total 3004	C 1913	N 498	O 578	S 15	0	4	0
1	E	399	Total 3076	C 1962	N 516	O 581	S 17	0	9	0
1	F	399	Total 3040	C 1937	N 509	O 579	S 15	0	6	0

- Molecule 2 is a protein called Aspartate aminotransferase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P S			
2	B	399	Total 3080	C 1964	N 515	O 583	P S 1 17	0	8	0

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total 16	C 8	N 1	O 6	P 1	0	0
3	C	1	Total 16	C 8	N 1	O 6	P 1	0	0
3	D	1	Total 16	C 8	N 1	O 6	P 1	0	0
3	E	1	Total 16	C 8	N 1	O 6	P 1	0	0
3	F	1	Total 15	C 8	N 1	O 5	P 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	372	Total 372	O 372	0	0
4	B	313	Total 313	O 313	0	0
4	C	130	Total 130	O 130	0	0
4	D	211	Total 211	O 211	0	0
4	E	297	Total 297	O 297	0	0
4	F	293	Total 293	O 293	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aspartate aminotransferase A

Chain A:  97%



- Molecule 1: Aspartate aminotransferase A

Chain C:  11% 87% 7% 5%



- Molecule 1: Aspartate aminotransferase A

Chain D:  4% 95% 5%




- Molecule 1: Aspartate aminotransferase A


Chain E:  94% 5%



- Molecule 1: Aspartate aminotransferase A



● Molecule 2: Aspartate aminotransferase A



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	103.54Å 93.01Å 123.07Å 90.00° 91.36° 90.00°	Depositor
Resolution (Å)	48.12 – 1.79 48.12 – 1.79	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.12-1.79) 98.1 (48.12-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.79Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.174 , 0.209 0.175 , 0.208	Depositor DCC
R_{free} test set	10689 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.381	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.021 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	19872	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.9139e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3202	0.55	0/4344
1	C	0.31	0/2956	0.48	0/4025
1	D	0.33	0/3086	0.50	0/4205
1	E	0.38	0/3170	0.55	0/4309
1	F	0.38	0/3128	0.54	0/4256
2	B	0.38	0/3149	0.54	0/4278
All	All	0.37	0/18691	0.53	0/25417

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3099	0	3107	10	0
1	C	2878	0	2798	23	0
1	D	3004	0	2900	13	0
1	E	3076	0	3040	17	0
1	F	3040	0	2976	12	0
2	B	3080	0	3039	14	0
3	A	16	0	8	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	16	0	7	2	0
3	D	16	0	7	2	0
3	E	16	0	8	1	0
3	F	15	0	6	2	0
4	A	372	0	0	2	0
4	B	313	0	0	2	0
4	C	130	0	0	1	0
4	D	211	0	0	2	0
4	E	297	0	0	2	0
4	F	293	0	0	2	0
All	All	19872	0	17896	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 83 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:240:LYS:HZ1	3:C:600:PLP:C4A	2.06	0.68
1:E:55:ASP:OD2	1:E:59:ARG:NH1	2.28	0.64
1:E:206[A]:MET:CE	1:E:239:SER:HB2	2.28	0.64
1:E:101:LYS:NZ	4:E:701:HOH:O	2.32	0.61
2:B:101:LYS:NZ	4:B:501:HOH:O	2.34	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	408/400 (102%)	401 (98%)	7 (2%)	0	100 100
1	C	379/400 (95%)	373 (98%)	6 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	401/400 (100%)	392 (98%)	9 (2%)	0	100	100
1	E	406/400 (102%)	398 (98%)	8 (2%)	0	100	100
1	F	403/400 (101%)	396 (98%)	7 (2%)	0	100	100
2	B	404/400 (101%)	399 (99%)	5 (1%)	0	100	100
All	All	2401/2400 (100%)	2359 (98%)	42 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/320 (102%)	325 (100%)	1 (0%)	92	91
1	C	297/320 (93%)	291 (98%)	6 (2%)	55	44
1	D	307/320 (96%)	301 (98%)	6 (2%)	55	44
1	E	321/320 (100%)	317 (99%)	4 (1%)	71	65
1	F	315/320 (98%)	313 (99%)	2 (1%)	86	84
2	B	319/319 (100%)	318 (100%)	1 (0%)	92	91
All	All	1885/1919 (98%)	1865 (99%)	20 (1%)	78	68

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	142[A]	VAL
1	E	155[B]	GLU
1	F	325	PHE
1	F	14	SER
1	C	235[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	257	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LLP	B	240	2	23,24,25	2.80	7 (30%)	25,32,34	1.77	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LLP	B	240	2	-	8/16/17/19	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	240	LLP	C4-C4'	9.45	1.64	1.46
2	B	240	LLP	C4'-NZ	6.16	1.47	1.27
2	B	240	LLP	C2'-C2	3.49	1.56	1.50
2	B	240	LLP	C4-C5	-2.46	1.38	1.42
2	B	240	LLP	C5'-C5	2.42	1.57	1.50

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	240	LLP	OP4-C5'-C5	4.00	116.97	109.35
2	B	240	LLP	CD-CE-NZ	3.44	119.35	110.93
2	B	240	LLP	C5'-C5-C6	-3.22	114.07	119.37
2	B	240	LLP	C4-C3-C2	3.19	122.17	120.19
2	B	240	LLP	CD-CG-CB	2.80	123.52	113.62

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	240	LLP	C4-C4'-NZ-CE
2	B	240	LLP	C5'-OP4-P-OP1
2	B	240	LLP	C5'-OP4-P-OP2
2	B	240	LLP	C5'-OP4-P-OP3
2	B	240	LLP	CG-CD-CE-NZ

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	240	LLP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PLP	E	600	-	16,16,16	1.08	2 (12%)	20,23,23	0.86	0
3	PLP	C	600	-	16,16,16	1.14	3 (18%)	20,23,23	1.03	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PLP	A	600	-	16,16,16	1.06	2 (12%)	20,23,23	1.09	1 (5%)
3	PLP	F	600	1	15,15,16	0.97	1 (6%)	20,22,23	1.59	5 (25%)
3	PLP	D	600	-	16,16,16	1.16	1 (6%)	20,23,23	1.01	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	E	600	-	-	5/8/8/8	0/1/1/1
3	PLP	C	600	-	-	5/8/8/8	0/1/1/1
3	PLP	A	600	-	-	5/8/8/8	0/1/1/1
3	PLP	F	600	1	-	3/6/6/8	0/1/1/1
3	PLP	D	600	-	-	5/8/8/8	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	600	PLP	C2-N1	2.61	1.38	1.33
3	D	600	PLP	C2-N1	2.57	1.38	1.33
3	A	600	PLP	C2-N1	2.46	1.38	1.33
3	F	600	PLP	C2-N1	2.31	1.38	1.33
3	A	600	PLP	C4-C4A	2.28	1.51	1.46

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	600	PLP	C6-C5-C4	3.41	120.84	118.16
3	F	600	PLP	O4P-C5A-C5	3.14	115.34	109.35
3	A	600	PLP	O3P-P-O4P	2.99	114.68	106.73
3	F	600	PLP	C4A-C4-C5	-2.77	118.08	120.94
3	D	600	PLP	C5-C6-N1	-2.35	119.90	123.82

There are no chirality outliers.

5 of 23 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	600	PLP	C5A-O4P-P-O1P
3	A	600	PLP	C5A-O4P-P-O2P

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	600	PLP	C5A-O4P-P-O3P
3	C	600	PLP	C5A-O4P-P-O1P
3	C	600	PLP	C5A-O4P-P-O2P

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	600	PLP	1	0
3	C	600	PLP	2	0
3	A	600	PLP	3	0
3	F	600	PLP	2	0
3	D	600	PLP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/400 (99%)	-0.31	0 100 100	9, 17, 38, 57	0
1	C	380/400 (95%)	0.76	45 (11%) 4 3	22, 44, 68, 79	0
1	D	399/400 (99%)	0.09	15 (3%) 40 35	16, 31, 69, 86	0
1	E	399/400 (99%)	-0.25	2 (0%) 91 89	12, 22, 41, 59	0
1	F	399/400 (99%)	-0.26	4 (1%) 82 80	11, 22, 48, 74	0
2	B	398/400 (99%)	-0.21	3 (0%) 86 84	10, 21, 46, 59	0
All	All	2374/2400 (98%)	-0.04	69 (2%) 51 46	9, 25, 57, 86	0

The worst 5 of 69 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	17	ILE	7.7
1	C	198	HIS	5.4
1	C	343	LYS	5.2
1	D	10	ARG	4.8
1	C	340	PRO	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	LLP	B	240	24/25	0.97	0.10	12,17,31,44	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PLP	C	600	16/16	0.93	0.17	24,33,41,43	16
3	PLP	D	600	16/16	0.94	0.18	20,29,36,37	16
3	PLP	E	600	16/16	0.96	0.13	18,27,36,37	0
3	PLP	A	600	16/16	0.97	0.12	16,22,36,38	0
3	PLP	F	600	15/16	0.97	0.09	18,21,33,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.