

Full wwPDB X-ray Structure Validation Report (i)

Dec 4, 2023 – 12:32 PM EST

PDB ID : 8F78

Title: Compound 1 bound to procaspase-6

Authors: Fan, P.; Zhao, Y.; Renslo, A.R.; Arkin, M.R.

Deposited on : 2022-11-18

Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	299	72%	12%	16%
1	В	299	75%	9%	16%
1	С	299	73%	8% •	18%
1	D	299	72%	10%	17%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 8291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Procaspase-6.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	۸	251	Total	С	N	О	S	0	1	0
1	A	231	2029	1296	350	368	15	0	1	
1	В	251	Total	С	N	О	S	0	1	0
1	Ъ	201	2029	1296	350	368	15	U	1	
1	С	246	Total	С	N	О	S	0	0	0
1		240	1977	1260	345	358	14	U	0	
1	D	247	Total	С	N	О	S	0	0	0
1	ש	241	1983	1264	344	361	14	U		

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	163	ALA	CYS	conflict	UNP P55212
A	294	GLU	-	expression tag	UNP P55212
A	295	ASN	-	expression tag	UNP P55212
A	296	LEU	-	expression tag	UNP P55212
A	297	TYR	-	expression tag	UNP P55212
A	298	PHE	-	expression tag	UNP P55212
A	299	GLN	-	expression tag	UNP P55212
В	163	ALA	CYS	conflict	UNP P55212
В	294	GLU	-	expression tag	UNP P55212
В	295	ASN	-	expression tag	UNP P55212
В	296	LEU	_	expression tag	UNP P55212
В	297	TYR	-	expression tag	UNP P55212
В	298	PHE	_	expression tag	UNP P55212
В	299	GLN	-	expression tag	UNP P55212
С	163	ALA	CYS	conflict	UNP P55212
С	294	GLU	-	expression tag	UNP P55212
С	295	ASN	-	expression tag	UNP P55212
С	296	LEU	-	expression tag	UNP P55212
С	297	TYR	-	expression tag	UNP P55212
С	298	PHE	-	expression tag	UNP P55212
С	299	GLN	_	expression tag	UNP P55212



Continued from previous page						
	\mathcal{C}	lontinued	l fr	rom	previous	page

Chain	Residue	Modelled	Actual	Comment	Reference
D	163	ALA	CYS	conflict	UNP P55212
D	294	GLU	-	expression tag	UNP P55212
D	295	ASN	-	expression tag	UNP P55212
D	296	LEU	-	expression tag	UNP P55212
D	297	TYR	-	expression tag	UNP P55212
D	298	PHE	-	expression tag	UNP P55212
D	299	GLN	-	expression tag	UNP P55212

 \bullet Molecule 2 is 5-fluoro-2-({[3-(pyrimidin-2-yl)pyridin-2-yl]amino}methyl)phenol (three-letter code: 2J6) (formula: $C_{16}H_{13}FN_4O)$ (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
2	Λ	1	Total	С	F	N	О	0	1	
2	A	1	44	32	2	8	2	0	1	
9	C	1	Total	С	F	N	О	0	1	
		1	44	32	2	8	2	0	1	

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	D	3	Total Cl 3 3	0	0

• Molecule 4 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	44	Total O 44 44	0	0
4	В	44	Total O 44 44	0	0
4	С	52	Total O 52 52	0	0
4	D	41	Total O 41 41	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Procaspase-6 Chain A: 16% MET SER SER ALA ALA ALA ARG GLY ARG GLY ARG GLY ARG GLY ALA ARG GLY ALA ALA ALA ALA ASN MET THR ASN MET THR ASP ALA ASP ALA ASP ARG GLY ARG ASN MET THR ASP ARG GLY ARG GLY ARG ASN ARG GLY ARG GLY ARG GLY ARG ASN ARG GLY AR • Molecule 1: Procaspase-6 Chain B: 75% 16% • Molecule 1: Procaspase-6 Chain C: 73% 8% 18% • Molecule 1: Procaspase-6 Chain D: 72% 17%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants	102.28Å 102.28Å 321.65Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	85.40 - 2.65	Depositor
resolution (A)	85.40 - 2.65	EDS
% Data completeness	97.7 (85.40-2.65)	Depositor
(in resolution range)	97.7 (85.40-2.65)	EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.23 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.164 , 0.207	Depositor
it, it free	0.168 , 0.205	DCC
R_{free} test set	2753 reflections $(5.12%)$	wwPDB-VP
Wilson B-factor (Å ²)	61.8	Xtriage
Anisotropy	0.880	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 43.3	EDS
L-test for twinning ²	$< L > = 0.52, < L^2> = 0.35$	Xtriage
Estimated twinning fraction	0.489 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8291	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 2J6, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Willi		# Z > 5	RMSZ	# Z > 5	
1	A	0.65	0/2079	0.78	0/2799	
1	В	0.65	0/2079	0.79	0/2799	
1	С	0.64	0/2023	0.79	0/2724	
1	D	0.65	0/2028	0.78	0/2731	
All	All	0.65	0/8209	0.78	0/11053	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2029	0	1994	22	0
1	В	2029	0	1994	18	0
1	С	1977	0	1943	23	0
1	D	1983	0	1951	21	0
2	A	44	0	24	4	0
2	С	44	0	26	7	0
3	A	1	0	0	0	0
3	D	3	0	0	0	0
4	A	44	0	0	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	В	44	0	0	0	0
4	С	52	0	0	2	0
4	D	41	0	0	2	0
All	All	8291	0	7932	81	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 5.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
1 D 101 1 D C 11 1	4 D. 402 HOH O	distance (Å)	overlap (Å)
1:D:164:ARG:HA	4:D:402:HOH:O	1.78	0.82
1:A:51:ASN:HD22	1:A:89:ASN:HD22	1.34	0.75
1:D:123:GLU:OE2	4:D:401:HOH:O	2.09	0.70
1:C:109:ALA:N	1:C:150:SER:OG	2.24	0.70
1:C:201:PRO:CA	1:C:281:MET:HE1	2.22	0.69
1:C:201:PRO:HA	1:C:281:MET:HE1	1.74	0.69
1:C:201:PRO:CB	1:C:281:MET:HE1	2.24	0.67
1:B:51:ASN:HD22	1:B:89:ASN:HD22	1.43	0.66
1:D:76:ARG:NH1	1:D:80:ASP:OD2	2.29	0.65
1:C:198:TYR:HA	2:C:301[A]:2J6:C13	2.26	0.65
1:D:51:ASN:HD22	1:D:89:ASN:HD22	1.52	0.58
1:B:145:GLY:HA2	1:B:152:VAL:HG22	1.87	0.57
1:B:112:ASP:OD2	1:B:291:LYS:HE2	2.04	0.56
1:D:112:ASP:OD2	1:D:291:LYS:HE2	2.06	0.56
1:A:118:PHE:CZ	1:A:139:LEU:HD13	2.42	0.55
1:C:43:ARG:O	1:C:111:ALA:HA	2.07	0.54
1:A:157:ILE:HD11	1:A:286:LEU:HD21	1.90	0.54
1:A:51:ASN:HD22	1:A:89:ASN:ND2	2.03	0.54
1:B:51:ASN:HD22	1:B:89:ASN:ND2	2.06	0.54
1:D:34:ALA:HA	1:D:285:LYS:HE2	1.90	0.54
2:A:301[A]:2J6:N22	2:A:301[A]:2J6:N7	2.46	0.53
1:B:252:VAL:O	1:B:256:VAL:HG23	2.09	0.52
1:C:51:ASN:HD22	1:C:89:ASN:HD22	1.57	0.52
2:C:301[B]:2J6:H2	4:C:408:HOH:O	2.09	0.52
1:A:145:GLY:HA2	1:A:152:VAL:HG22	1.91	0.52
1:D:51:ASN:HD22	1:D:89:ASN:ND2	2.07	0.51
1:A:281[B]:MET:HE2	1:B:275:VAL:CG1	2.41	0.51
1:C:76:ARG:HD3	1:C:80:ASP:OD2	2.11	0.50
1:D:62:PRO:O	1:D:121:HIS:HE1	1.94	0.50
1:C:188:ASN:HD22	1:C:223:VAL:H	1.60	0.50
		Continu	ed on next page



 $Continued\ from\ previous\ page...$

Atom-1	Atom-2	Interatomic	Clash
Atom-1		${\rm distance}({\rm \AA})$	overlap (Å)
1:A:120:SER:O	1:A:162:ALA:HA	2.12	0.49
1:C:126:HIS:HB3	1:C:133:LYS:HB2	1.93	0.49
1:A:126:HIS:HB3	1:A:133:LYS:HB2	1.94	0.49
1:C:188:ASN:HD21	1:C:221:GLU:HB3	1.78	0.49
2:A:301[A]:2J6:H7	1:B:200:LEU:HD23	1.95	0.48
1:A:281[B]:MET:HG2	1:B:277:CYS:HB3	1.95	0.48
1:C:244:GLU:OE1	1:C:246:THR:OG1	2.28	0.48
1:C:216:TYR:OH	1:D:144:LYS:HE3	2.13	0.48
1:D:123:GLU:HG2	1:D:164:ARG:NH1	2.29	0.47
1:C:126:HIS:CG	1:C:133:LYS:HD3	2.49	0.47
1:A:75:THR:HA	1:A:85:VAL:HG11	1.98	0.46
1:A:131:ASP:HB2	1:D:94:GLU:HG3	1.96	0.46
1:B:126:HIS:HB3	1:B:133:LYS:HB2	1.99	0.45
1:B:157:ILE:HD11	1:B:286:LEU:HD21	1.98	0.45
1:A:203:GLY:N	1:A:281[B]:MET:SD	2.85	0.45
1:D:71:ARG:HH11	1:D:89:ASN:ND2	2.15	0.45
2:A:301[B]:2J6:N7	4:A:401:HOH:O	2.26	0.45
1:C:51:ASN:HD22	1:C:89:ASN:ND2	2.14	0.45
1:B:125:ASN:HB3	1:B:200:LEU:HD11	1.98	0.45
1:D:220:ARG:HA	1:D:226:SER:HA	1.98	0.45
1:A:145:GLY:CA	1:A:152:VAL:HG22	2.47	0.44
1:B:120:SER:O	1:B:162:ALA:HA	2.18	0.44
1:C:124:GLY:HA3	4:C:414:HOH:O	2.18	0.43
1:C:201:PRO:CB	1:C:281:MET:CE	2.96	0.43
1:C:48:LEU:HD21	1:C:103:VAL:HG21	2.01	0.43
1:D:211:SER:HA	1:D:228:TYR:CG	2.53	0.43
1:A:47:ALA:HA	1:A:115:VAL:O	2.19	0.43
1:A:252:VAL:O	1:A:256:VAL:HG23	2.18	0.43
1:C:198:TYR:HA	2:C:301[A]:2J6:C14	2.49	0.43
1:C:214:GLU:HB2	2:C:301[A]:2J6:C9	2.49	0.43
1:C:118:PHE:CZ	1:C:139:LEU:HD13	2.53	0.42
1:B:211:SER:HA	1:B:228:TYR:CG	2.55	0.42
1:A:275:VAL:HG11	1:B:281[B]:MET:HE2	2.02	0.42
2:C:301[B]:2J6:N22	2:C:301[B]:2J6:N7	2.67	0.42
1:B:145:GLY:CA	1:B:152:VAL:HG22	2.50	0.41
1:A:211:SER:HA	1:A:228:TYR:CG	2.55	0.41
1:A:116:CYS:O	1:A:158:PHE:HA	2.20	0.41
1:D:145:GLY:HA2	1:D:152:VAL:HG22	2.02	0.41
1:A:31:PHE:CD2	1:B:255:LYS:HE2	2.56	0.41
1:C:198:TYR:HB3	2:C:301[A]:2J6:C2	2.51	0.41
1:A:37:TYR:O	1:A:39:MET:HG2	2.20	0.41



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:58:HIS:CE1	1:D:101:HIS:CG	3.09	0.41
2:A:301[A]:2J6:H7	1:B:200:LEU:CD2	2.51	0.41
1:C:219:HIS:HB2	1:C:227:TRP:CD2	2.56	0.41
2:C:301[B]:2J6:C14	1:D:198:TYR:HA	2.50	0.41
1:D:112:ASP:OD2	1:D:291:LYS:CE	2.69	0.41
1:D:126:HIS:HB3	1:D:133:LYS:HB2	2.03	0.41
1:D:241:SER:HA	1:D:288:PHE:O	2.22	0.40
1:D:227:TRP:CD2	1:D:259:ARG:HD3	2.56	0.40
1:B:115:VAL:HG22	1:B:157:ILE:HB	2.04	0.40
1:A:62:PRO:O	1:A:121:HIS:HE1	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$248/299 \ (83\%)$	234 (94%)	14 (6%)	0	100	100
1	В	248/299 (83%)	234 (94%)	14 (6%)	0	100	100
1	С	242/299 (81%)	228 (94%)	13 (5%)	1 (0%)	34	48
1	D	243/299 (81%)	232 (96%)	11 (4%)	0	100	100
All	All	981/1196 (82%)	928 (95%)	52 (5%)	1 (0%)	51	69

All (1) Ramachandran outliers are listed below:

\mathbf{M}	ol	Chain	Res	Type
1		С	201	PRO



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	220/260~(85%)	216 (98%)	4 (2%)	59 75		
1	В	220/260 (85%)	215 (98%)	5 (2%)	50 68		
1	С	214/260 (82%)	213 (100%)	1 (0%)	88 94		
1	D	215/260 (83%)	211 (98%)	4 (2%)	57 74		
All	All	869/1040 (84%)	855 (98%)	14 (2%)	62 78		

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	VAL
1	A	125	ASN
1	A	150	SER
1	A	299	GLN
1	В	86	LYS
1	В	106	VAL
1	В	146	ASP
1	В	164	ARG
1	В	294	GLU
1	С	76	ARG
1	D	76	ARG
1	D	100	ILE
1	D	106	VAL
1	D	281	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	HIS
1	A	89	ASN
1	A	224	ASN
1	A	230	GLN
1	В	58	HIS



Continued from previous page...

Mol	Chain	Res	Type
1	В	89	ASN
1	В	224	ASN
1	В	230	GLN
1	С	41	HIS
1	С	52	HIS
1	С	89	ASN
1	C C	121	HIS
1	С	167	GLN
1	С	188	ASN
1	С	224	ASN
1	C	230	GLN
1	D	89	ASN
1	D	121	HIS
1	D	224	ASN
1	D	230	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Tuno	Chain	Res	Dag	Dag	Dag	Link	Bo	ond leng	$_{ m ths}$	В	ond ang	gles
MIOI	Mol Type Cha	Chain		LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2			
2	2J6	С	301[A]	-	24,24,24	0.84	1 (4%)	31,32,32	1.83	7 (22%)			
2	2J6	С	301[B]	-	24,24,24	1.12	2 (8%)	31,32,32	2.12	12 (38%)			
2	2J6	A	301[A]	-	24,24,24	1.07	1 (4%)	31,32,32	1.94	10 (32%)			
2	2J6	A	301[B]	-	24,24,24	0.92	1 (4%)	31,32,32	2.18	11 (35%)			

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2J6	С	301[A]	-	-	0/9/9/9	0/3/3/3
2	2J6	С	301[B]	-	-	2/9/9/9	0/3/3/3
2	2J6	A	301[A]	-	-	0/9/9/9	0/3/3/3
2	2J6	A	301[B]	-	-	2/9/9/9	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
2	A	301[A]	2J6	C4-N7	4.20	1.41	1.34
2	A	301[B]	2J6	C4-N7	3.36	1.39	1.34
2	С	301[B]	2J6	C4-N7	3.03	1.39	1.34
2	С	301[B]	2J6	C13-C12	2.86	1.42	1.37
2	С	301[A]	2J6	C4-N7	2.19	1.38	1.34

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	301[B]	2J6	C21-N22-C17	5.27	121.59	116.08
2	A	301[A]	2J6	C19-N18-C17	4.81	121.11	116.08
2	С	301[B]	2J6	F16-C12-C13	4.64	124.88	118.25
2	A	301[B]	2J6	C19-N18-C17	4.43	120.71	116.08
2	С	301[B]	2J6	C19-N18-C17	4.43	120.71	116.08
2	A	301[A]	2J6	C21-N22-C17	4.33	120.61	116.08
2	С	301[A]	2J6	C8-N7-C4	-4.24	117.56	123.14
2	A	301[B]	2J6	C8-N7-C4	-4.10	117.74	123.14
2	A	301[B]	2J6	C3-C4-N7	3.94	123.73	121.84
2	С	301[A]	2J6	C19-N18-C17	3.86	120.11	116.08
2	С	301[A]	2J6	C21-N22-C17	3.84	120.09	116.08
2	С	301[B]	2J6	C10-C9-C14	3.71	122.12	118.16
2	С	301[B]	2J6	F16-C12-C11	-3.32	112.90	118.54



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	301[A]	2J6	C3-C17-N18	3.27	120.65	116.99
2	A	301[B]	2J6	C3-C17-N22	3.12	120.48	116.99
2	С	301[B]	2J6	C20-C19-N18	-3.11	118.34	123.43
2	A	301[A]	2J6	C2-C3-C17	2.96	123.10	118.55
2	A	301[A]	2J6	C20-C19-N18	-2.76	118.92	123.43
2	A	301[B]	2J6	C20-C21-N22	-2.75	118.94	123.43
2	С	301[B]	2J6	C8-N7-C4	-2.72	119.55	123.14
2	С	301[B]	2J6	C21-N22-C17	2.67	118.86	116.08
2	С	301[B]	2J6	C3-C17-N22	2.66	119.97	116.99
2	A	301[A]	2J6	C2-C3-C4	2.59	119.39	117.46
2	С	301[A]	2J6	C20-C19-N18	-2.56	119.24	123.43
2	С	301[B]	2J6	C6-N5-C4	2.43	121.58	116.77
2	С	301[A]	2J6	C20-C21-N22	-2.40	119.51	123.43
2	A	301[A]	2J6	C20-C21-N22	-2.38	119.54	123.43
2	A	301[A]	2J6	C6-N5-C4	2.33	121.39	116.77
2	С	301[A]	2J6	C10-C9-C14	2.31	120.63	118.16
2	A	301[B]	2J6	C6-N5-C4	2.30	121.34	116.77
2	A	301[B]	2J6	C11-C12-C13	-2.29	120.31	123.29
2	A	301[B]	2J6	C20-C19-N18	-2.26	119.73	123.43
2	A	301[B]	2J6	N22-C17-N18	-2.25	120.70	125.57
2	A	301[A]	2J6	C11-C12-C13	-2.20	120.43	123.29
2	С	301[B]	2J6	C2-C3-C4	2.18	119.08	117.46
2	С	301[B]	2J6	C20-C21-N22	-2.12	119.97	123.43
2	С	301[B]	2J6	C9-C8-N7	-2.05	106.69	113.38
2	A	301[B]	2J6	C8-C9-C14	2.04	122.71	120.00
2	С	301[A]	2J6	C6-N5-C4	2.03	120.79	116.77
2	A	301[A]	2J6	N22-C17-N18	-2.01	121.23	125.57

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301[B]	2J6	N7-C8-C9-C14
2	A	301[B]	2J6	N7-C8-C9-C10
2	С	301[B]	2J6	N7-C8-C9-C10
2	С	301[B]	2J6	N7-C8-C9-C14


There are no ring outliers.

4 monomers are involved in 11 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	301[A]	2J6	4	0
2	С	301[B]	2J6	3	0
2	A	301[A]	2J6	3	0
2	A	301[B]	2J6	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	251/299~(83%)	-0.01	5 (1%) 65 60	61, 79, 120, 204	0
1	В	251/299~(83%)	-0.11	3 (1%) 79 77	61, 78, 118, 141	0
1	С	$246/299 \ (82\%)$	-0.11	1 (0%) 92 93	58, 76, 111, 141	0
1	D	$247/299 \ (82\%)$	-0.09	2 (0%) 86 85	58, 76, 118, 176	0
All	All	995/1196 (83%)	-0.08	11 (1%) 80 79	58, 78, 117, 204	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	297	TYR	10.0
1	A	295	ASN	5.3
1	A	296	LEU	5.2
1	A	298	PHE	5.0
1	A	294	GLU	4.2
1	В	295	ASN	3.6
1	В	57	TRP	3.0
1	В	281[A]	MET	2.4
1	D	260	ARG	2.1
1	D	263	PHE	2.1
1	С	78	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	CL	D	302	1/1	0.87	0.08	107,107,107,107	0
2	2J6	С	301[B]	22/22	0.91	0.17	37,61,70,75	22
2	2J6	С	301[A]	22/22	0.91	0.17	37,56,67,70	22
2	2J6	A	301[B]	22/22	0.94	0.16	47,70,74,79	22
2	2J6	A	301[A]	22/22	0.94	0.16	42,67,73,78	22
3	CL	D	301	1/1	0.96	0.07	94,94,94,94	0
3	CL	A	302	1/1	0.97	0.12	73,73,73,73	0
3	CL	D	303	1/1	0.99	0.08	75,75,75,75	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

