

# Full wwPDB X-ray Structure Validation Report (i)

#### Nov 7, 2023 – 01:41 PM JST

PDB ID	:	5F8V
Title	:	Crystal structure of PLP bound phosphoserine aminotransferase (PSAT) from
		Trichomonas vaginalis
Authors	:	Singh, R.K.; Gourinath, S.
Deposited on	:	2015-12-09
Resolution	:	2.14  Å(reported)
Authors Deposited on Resolution	: : :	Singh, R.K.; Gourinath, S. 2015-12-09 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	379	94%	••
1	В	379	2% 95%	••
1	С	379	92%	• •
1	D	379	% 93%	• •
1	Е	379	2% 94%	•••
1	F	379	% <b>88%</b>	•• 9%



Mol	Chain	Length	Quality of chain	
1	G	379	94% •	•
1	Н	379	<sup>2%</sup> 93%	•



## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 23601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		A	Atom	s			ZeroOcc	AltConf	Trace
1		967	Total	С	Ν	0	Р	S	0	0	0
1	A	307	2933	1878	499	531	1	24	0	0	0
1	F	264	Total	С	Ν	0	Р	S	0	0	0
1	Ľ	504	2898	1852	495	526	1	24	0	0	0
1	В	367	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	0	0
1	D	507	2895	1854	492	524	1	24	0	0	0
1	С	363	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	0	0
1	U	505	2830	1809	483	513	1	24	0	0	0
1	Л	267	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	0	0
1	D	507	2886	1848	495	518	1	24	0	0	U
1	F	343	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	0	0
1	Ľ	040	2680	1723	452	481	1	23	0	0	0
1	C	367	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	0	0
1	G	507	2887	1854	495	513	1	24	0	0	0
1	Ц	366	Total	С	Ν	0	Р	S	0	0	0
	11	500	2843	1825	483	510	1	24		0	0

• Molecule 1 is a protein called Aminotransferase, class V family protein.

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	372	LEU	-	expression tag	UNP A2DW27
А	373	GLU	-	expression tag	UNP A2DW27
А	374	HIS	-	expression tag	UNP A2DW27
А	375	HIS	-	expression tag	UNP A2DW27
А	376	HIS	-	expression tag	UNP A2DW27
А	377	HIS	-	expression tag	UNP A2DW27
А	378	HIS	-	expression tag	UNP A2DW27
А	379	HIS	-	expression tag	UNP A2DW27
E	372	LEU	-	expression tag	UNP A2DW27
E	373	GLU	-	expression tag	UNP A2DW27
Е	374	HIS	-	expression tag	UNP A2DW27
E	375	HIS	-	expression tag	UNP A2DW27
E	376	HIS	-	expression tag	UNP A2DW27



Chain	Residue	Modelled	Actual	Comment	Reference
Е	377	HIS	_	expression tag	UNP A2DW27
Е	378	HIS	-	expression tag	UNP A2DW27
Е	379	HIS	_	expression tag	UNP A2DW27
В	372	LEU	-	expression tag	UNP A2DW27
В	373	GLU	_	expression tag	UNP A2DW27
В	374	HIS	-	expression tag	UNP A2DW27
В	375	HIS	-	expression tag	UNP A2DW27
В	376	HIS	-	expression tag	UNP A2DW27
В	377	HIS	-	expression tag	UNP A2DW27
В	378	HIS	-	expression tag	UNP A2DW27
В	379	HIS	-	expression tag	UNP A2DW27
С	372	LEU	-	expression tag	UNP A2DW27
С	373	GLU	-	expression tag	UNP A2DW27
С	374	HIS	-	expression tag	UNP A2DW27
С	375	HIS	-	expression tag	UNP A2DW27
С	376	HIS	-	expression tag	UNP A2DW27
С	377	HIS	-	expression tag	UNP A2DW27
С	378	HIS	-	expression tag	UNP A2DW27
С	379	HIS	-	expression tag	UNP A2DW27
D	372	LEU	-	expression tag	UNP A2DW27
D	373	GLU	-	expression tag	UNP A2DW27
D	374	HIS	-	expression tag	UNP A2DW27
D	375	HIS	-	expression tag	UNP A2DW27
D	376	HIS	-	expression tag	UNP A2DW27
D	377	HIS	-	expression tag	UNP A2DW27
D	378	HIS	-	expression tag	UNP A2DW27
D	379	HIS	-	expression tag	UNP A2DW27
F	372	LEU	-	expression tag	UNP A2DW27
F	373	GLU	-	expression tag	UNP A2DW27
F	374	HIS	-	expression tag	UNP A2DW27
F	375	HIS	-	expression tag	UNP A2DW27
F	376	HIS	-	expression tag	UNP A2DW27
F	377	HIS	-	expression tag	UNP A2DW27
F	378	HIS	-	expression tag	UNP A2DW27
F	379	HIS	-	expression tag	UNP A2DW27
G	372	LEU	-	expression tag	UNP A2DW27
G	373	GLU	-	expression tag	UNP A2DW27
G	374	HIS	-	expression tag	UNP A2DW27
G	375	HIS	-	expression tag	UNP A2DW27
G	376	HIS	-	expression tag	UNP A2DW27
G	377	HIS	-	expression tag	UNP A2DW27
G	378	HIS	-	expression tag	UNP A2DW27



5F	'8V

Chain	Residue	Modelled	Actual	Comment	Reference
G	379	HIS	-	expression tag	UNP A2DW27
Н	372	LEU	-	expression tag	UNP A2DW27
Н	373	GLU	-	expression tag	UNP A2DW27
Н	374	HIS	-	expression tag	UNP A2DW27
Н	375	HIS	-	expression tag	UNP A2DW27
Н	376	HIS	-	expression tag	UNP A2DW27
Н	377	HIS	-	expression tag	UNP A2DW27
Н	378	HIS	-	expression tag	UNP A2DW27
Н	379	HIS	-	expression tag	UNP A2DW27

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	122	Total O 122 122	0	0
2	Е	50	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 50 & 50 \end{array}$	0	0
2	В	98	Total O 98 98	0	0
2	С	41	Total         O           41         41	0	0
2	D	126	Total O 126 126	0	0
2	F	77	Total O 77 77	0	0
2	G	147	Total O 147 147	0	0
2	Н	88	Total         O           88         88	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminotransferase, class V family protein











## 4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	51.97Å 235.43Å 134.52Å	Deperitor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.02^{\circ}$ $90.00^{\circ}$	Depositor	
$\mathbf{P}_{\text{oscolution}}(\hat{\mathbf{A}})$	37.42 - 2.14	Depositor	
Resolution (A)	37.42 - 2.14	EDS	
% Data completeness	99.4 (37.42-2.14)	Depositor	
(in resolution range)	99.4 (37.42-2.14)	EDS	
$R_{merge}$	(Not available)	Depositor	
$R_{sym}$	0.15	Depositor	
$< I/\sigma(I) > 1$	$2.99 (at 2.14 \text{\AA})$	Xtriage	
Refinement program	REFMAC 5.8.0103	Depositor	
D D	0.209 , $0.239$	Depositor	
$\mathbf{n},  \mathbf{n}_{free}$	0.212 , $0.238$	DCC	
$R_{free}$ test set	8877 reflections $(5.01%)$	wwPDB-VP	
Wilson B-factor $(Å^2)$	21.8	Xtriage	
Anisotropy	0.162	Xtriage	
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , $10.2$	EDS	
L-test for $twinning^2$	$<  L  > = 0.47, < L^2 > = 0.30$	Xtriage	
Estimated twinning fraction	0.288 for h,-k,-l	Xtriage	
Pererted twinning freation	0.759 for H, K, L	Deperitor	
Reported twinning fraction	0.241 for -h,-k,l	Depositor	
Outliers	2 of 177272 reflections $(0.001\%)$	Xtriage	
$F_o, F_c$ correlation	0.94	EDS	
Total number of atoms	23601	wwPDB-VP	
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 43.65 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7105e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for a centric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bo	ond angles
MIOI			# Z  > 5	RMSZ	# Z  > 5
1	А	0.41	0/2988	0.63	0/4049
1	В	0.40	0/2948	0.62	0/3998
1	С	0.39	0/2879	0.57	0/3907
1	D	0.42	0/2939	0.64	0/3986
1	Е	0.39	0/2949	0.58	0/3995
1	F	0.39	0/2727	0.60	1/3705~(0.0%)
1	G	0.41	0/2942	0.65	0/3993
1	Н	0.39	0/2895	0.60	0/3931
All	All	0.40	0/23267	0.61	1/31564~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	F	201	GLN	N-CA-C	7.29	130.68	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	201	GLN	Peptide



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2933	0	2857	5	0
1	В	2895	0	2794	3	0
1	С	2830	0	2722	5	0
1	D	2886	0	2784	4	0
1	Е	2898	0	2826	2	0
1	F	2680	0	2562	6	0
1	G	2887	0	2795	2	0
1	Н	2843	0	2724	5	0
2	А	122	0	0	0	0
2	В	98	0	0	0	0
2	С	41	0	0	0	0
2	D	126	0	0	0	0
2	Е	50	0	0	0	0
2	F	77	0	0	0	0
2	G	147	0	0	0	0
2	Н	88	0	0	0	0
All	All	23601	0	22064	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:ARG:HD2	1:A:217:ASP:OD1	1.99	0.61
1:B:93:HIS:HB2	1:B:149:ALA:HB2	1.86	0.57
1:F:136:VAL:HG22	1:F:299:TRP:CZ2	2.47	0.49
1:F:176:ILE:HD12	1:F:196:ILE:HD11	1.95	0.49
1:D:63:LEU:HD22	1:D:187:ILE:HD12	1.96	0.48
1:A:63:LEU:HD22	1:A:187:ILE:HD12	1.95	0.48
1:G:63:LEU:HD22	1:G:187:ILE:HD12	1.96	0.47
1:G:282:TYR:CZ	1:G:305:MET:HG2	2.49	0.47
1:E:200:ALA:HB3	1:E:206:LEU:O	2.14	0.47
1:D:93:HIS:HB2	1:D:149:ALA:HB2	1.97	0.46
1:D:282:TYR:CZ	1:D:305:MET:HG2	2.50	0.46



A 4 amo 1	A.t.a.m. D	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:H:193:VAL:CG1	1:H:196:ILE:HD13	2.46	0.46
1:A:81:GLN:O	1:A:85:ILE:HG12	2.16	0.46
1:H:93:HIS:HB2	1:H:149:ALA:HB2	1.98	0.46
1:F:96:VAL:HG12	1:F:149:ALA:HB3	1.99	0.45
1:F:170:VAL:HG22	1:F:171:PRO:HD2	1.98	0.45
1:H:193:VAL:HG11	1:H:196:ILE:HD13	1.97	0.45
1:H:176:ILE:HD12	1:H:196:ILE:HD11	1.99	0.45
1:F:202:LLP:NZ	1:F:202:LLP:O3	2.50	0.44
1:A:282:TYR:CZ	1:A:305:MET:HG2	2.52	0.44
1:B:48:TRP:CE2	1:B:248:THR:HG22	2.53	0.44
1:A:93:HIS:HB2	1:A:149:ALA:HB2	2.00	0.44
1:C:200:ALA:HB3	1:C:206:LEU:O	2.18	0.43
1:D:72:LEU:HD21	1:D:237:VAL:HG22	2.01	0.43
1:C:325:VAL:HG21	1:C:337:LYS:HB3	2.01	0.42
1:C:96:VAL:HG12	1:C:149:ALA:HB3	2.01	0.42
1:C:48:TRP:CE2	1:C:248:THR:HG22	2.54	0.41
1:H:125:VAL:HG21	1:H:151:PHE:CE2	2.55	0.41
1:C:63:LEU:HD22	1:C:187:ILE:HD12	2.03	0.41
1:E:93:HIS:HB2	1:E:149:ALA:HB2	2.03	0.41
1:B:81:GLN:O	1:B:85:ILE:HG12	2.21	0.41
1:F:158:THR:OG1	1:F:202:LLP:O3	2.28	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	364/379~(96%)	356~(98%)	8 (2%)	0	100	100
1	В	364/379~(96%)	353~(97%)	11 (3%)	0	100	100
1	С	360/379~(95%)	353 (98%)	7 (2%)	0	100	100
1	D	364/379~(96%)	351 (96%)	13 (4%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	Ε	361/379~(95%)	357~(99%)	4 (1%)	0	100	100
1	F	336/379~(89%)	328~(98%)	8 (2%)	0	100	100
1	G	364/379~(96%)	357~(98%)	7~(2%)	0	100	100
1	Н	363/379~(96%)	351~(97%)	11 (3%)	1 (0%)	41	36
All	All	2876/3032~(95%)	2806~(98%)	69~(2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Н	369	PRO

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percenti	$\mathbf{les}$
1	А	309/320~(97%)	308 (100%)	1 (0%)	92 94	1
1	В	299/320~(93%)	298 (100%)	1 (0%)	92 94	1
1	С	290/320~(91%)	287~(99%)	3 (1%)	76 79	)
1	D	296/320~(92%)	291~(98%)	5(2%)	60 63	3
1	Ε	305/320~(95%)	302~(99%)	3~(1%)	76 79	)
1	$\mathbf{F}$	273/320~(85%)	272 (100%)	1 (0%)	91 93	3
1	G	297/320~(93%)	291~(98%)	6(2%)	55 57	7
1	Н	$28\overline{6}/320~(89\%)$	283~(99%)	3 (1%)	76 79	)
All	All	2355/2560~(92%)	2332 (99%)	23 (1%)	76 79	)

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	32	ARG
1	Е	283	GLU
1	Е	323	LYS



Mol	Chain	Res	Type
1	Е	335	THR
1	В	294	ARG
1	С	74	VAL
1	С	122	VAL
1	С	335	THR
1	D	132	ASN
1	D	158	THR
1	D	203	ASN
1	D	252	TYR
1	D	292	VAL
1	F	158	THR
1	G	5	ARG
1	G	74	VAL
1	G	158	THR
1	G	169	ASP
1	G	206	LEU
1	G	332	LYS
1	Н	16	VAL
1	Н	158	THR
1	Н	279	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	30	ASN
1	G	240	ASN
1	Н	70	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



Mol	Type	Chain	Dog	Link	Bo	ond leng	ths	B	ond ang	gles
WIOI	туре	Ullalli	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
1	LLP	С	202	1	23,24,25	2.94	5 (21%)	$25,\!32,\!34$	1.49	4 (16%)
1	LLP	F	202	1	23,24,25	2.86	5 (21%)	25,32,34	1.67	10 (40%)
1	LLP	D	202	1	23,24,25	2.78	5 (21%)	25,32,34	1.63	6 (24%)
1	LLP	В	202	1	23,24,25	2.85	5 (21%)	25,32,34	1.54	5 (20%)
1	LLP	А	202	1	23,24,25	2.74	5 (21%)	25,32,34	1.51	5 (20%)
1	LLP	Е	202	1	23,24,25	2.83	5 (21%)	25,32,34	1.64	7 (28%)
1	LLP	G	202	1	23,24,25	2.75	5 (21%)	25,32,34	1.53	5 (20%)
1	LLP	Н	202	1	23,24,25	2.80	5 (21%)	25,32,34	1.80	7 (28%)

expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	С	202	1	-	4/16/17/19	0/1/1/1
1	LLP	F	202	1	-	5/16/17/19	0/1/1/1
1	LLP	D	202	1	-	4/16/17/19	0/1/1/1
1	LLP	В	202	1	-	4/16/17/19	0/1/1/1
1	LLP	А	202	1	-	5/16/17/19	0/1/1/1
1	LLP	Е	202	1	-	5/16/17/19	0/1/1/1
1	LLP	G	202	1	-	8/16/17/19	0/1/1/1
1	LLP	Н	202	1	-	8/16/17/19	0/1/1/1

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
1	С	202	LLP	C3-C2	8.90	1.49	1.40
1	F	202	LLP	C3-C2	8.44	1.49	1.40
1	Е	202	LLP	C3-C2	8.38	1.49	1.40
1	В	202	LLP	C3-C2	7.90	1.48	1.40
1	Н	202	LLP	C3-C2	7.78	1.48	1.40
1	D	202	LLP	C3-C2	7.74	1.48	1.40
1	G	202	LLP	C3-C2	7.10	1.48	1.40
1	А	202	LLP	C3-C2	6.78	1.47	1.40
1	А	202	LLP	C4-C5	6.76	1.50	1.42



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	202	LLP	C4-C5	6.71	1.50	1.42
1	Н	202	LLP	C4-C5	6.70	1.50	1.42
1	D	202	LLP	C4-C5	6.39	1.50	1.42
1	В	202	LLP	C4-C5	6.30	1.49	1.42
1	F	202	LLP	C4-C5	6.30	1.49	1.42
1	Е	202	LLP	C4-C5	6.29	1.49	1.42
1	С	202	LLP	C4-C5	6.24	1.49	1.42
1	G	202	LLP	C4'-NZ	6.05	1.47	1.27
1	Н	202	LLP	C4'-NZ	6.04	1.47	1.27
1	А	202	LLP	C4'-NZ	6.00	1.47	1.27
1	F	202	LLP	C4'-NZ	6.00	1.47	1.27
1	D	202	LLP	C4'-NZ	5.93	1.47	1.27
1	В	202	LLP	C4'-NZ	5.85	1.46	1.27
1	С	202	LLP	C4'-NZ	5.62	1.46	1.27
1	В	202	LLP	C4-C3	5.60	1.49	1.40
1	С	202	LLP	C4-C3	5.54	1.49	1.40
1	Е	202	LLP	C4'-NZ	5.50	1.45	1.27
1	Е	202	LLP	C4-C3	5.31	1.48	1.40
1	А	202	LLP	C4-C3	5.03	1.48	1.40
1	D	202	LLP	C4-C3	4.90	1.48	1.40
1	F	202	LLP	C4-C3	4.83	1.48	1.40
1	G	202	LLP	C4-C3	4.75	1.47	1.40
1	Н	202	LLP	C4-C3	4.51	1.47	1.40
1	G	202	LLP	C4-C4'	3.50	1.53	1.46
1	С	202	LLP	C4-C4'	3.46	1.53	1.46
1	В	202	LLP	C4-C4'	3.44	1.53	1.46
1	А	202	LLP	C4-C4'	3.32	1.52	1.46
1	F	202	LLP	C4-C4'	3.22	1.52	1.46
1	Н	202	LLP	C4-C4'	3.20	1.52	1.46
1	D	202	LLP	C4-C4'	3.15	1.52	1.46
1	Е	202	LLP	C4-C4'	2.98	1.52	1.46

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
1	G	202	LLP	C3-C4-C5	-4.28	114.97	118.26
1	Н	202	LLP	C4-C4'-NZ	-3.86	106.60	124.31
1	Е	202	LLP	C4-C4'-NZ	-3.71	107.30	124.31
1	Е	202	LLP	C4-C3-C2	-3.58	117.97	120.19
1	С	202	LLP	C3-C4-C5	-3.54	115.54	118.26
1	D	202	LLP	C3-C4-C5	-3.46	115.60	118.26
1	Н	202	LLP	C5-C4-C4'	3.45	127.22	121.56



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	$Ideal(^{o})$
1	D	202	LLP	OP2-P-OP4	-3.42	97.62	106.73
1	А	202	LLP	C3-C4-C5	-3.42	115.63	118.26
1	В	202	LLP	C3-C4-C5	-3.21	115.80	118.26
1	F	202	LLP	C3-C4-C5	-3.14	115.85	118.26
1	С	202	LLP	C4-C3-C2	-3.08	118.28	120.19
1	Н	202	LLP	C3-C4-C5	-2.95	115.99	118.26
1	В	202	LLP	OP3-P-OP4	-2.87	99.09	106.73
1	D	202	LLP	C4-C4'-NZ	-2.85	111.22	124.31
1	В	202	LLP	C4-C4'-NZ	-2.82	111.36	124.31
1	F	202	LLP	C4-C4'-NZ	-2.77	111.57	124.31
1	F	202	LLP	OP3-P-OP4	-2.71	99.53	106.73
1	А	202	LLP	C4-C4'-NZ	-2.65	112.13	124.31
1	Н	202	LLP	OP3-P-OP4	-2.64	99.70	106.73
1	Е	202	LLP	C6-N1-C2	2.57	123.92	119.17
1	Н	202	LLP	C6-N1-C2	2.55	123.89	119.17
1	F	202	LLP	C6-N1-C2	2.54	123.88	119.17
1	D	202	LLP	C6-N1-C2	2.52	123.84	119.17
1	С	202	LLP	C4-C4'-NZ	-2.45	113.05	124.31
1	В	202	LLP	C6-N1-C2	2.45	123.71	119.17
1	А	202	LLP	CD-CE-NZ	2.43	116.90	110.93
1	F	202	LLP	OP3-P-OP2	2.41	116.84	107.64
1	F	202	LLP	CD-CG-CB	2.34	121.91	113.62
1	G	202	LLP	CD-CE-NZ	2.34	116.65	110.93
1	F	202	LLP	O3-C3-C2	2.33	122.57	117.49
1	С	202	LLP	C6-N1-C2	2.30	123.44	119.17
1	G	202	LLP	C4-C4'-NZ	-2.27	113.88	124.31
1	Е	202	LLP	OP3-P-OP4	-2.25	100.75	106.73
1	А	202	LLP	C6-N1-C2	2.22	123.27	119.17
1	G	202	LLP	C6-N1-C2	2.20	123.24	119.17
1	В	202	LLP	C4-C3-C2	-2.18	118.84	120.19
1	Е	202	LLP	C5-C4-C4'	2.15	125.09	121.56
1	Н	202	LLP	O3-C3-C2	2.15	122.17	117.49
1	F	202	LLP	C4-C3-C2	-2.14	118.86	120.19
1	A	202	LLP	C5-C4-C4 <sup>7</sup>	2.14	125.07	121.56
1	F	202	LLP	$CD-CE-N\overline{Z}$	2.13	116.15	110.93
1	H	$20\overline{2}$	LLP	$C4-C3-\overline{C2}$	-2.12	118.88	120.19
1	D	202	LLP	C5-C4-C4'	2.09	125.00	121.56
1	Е	202	LLP	OP3-P-OP2	2.07	115.56	107.64
1	G	202	LLP	C5-C4-C4 <sup>7</sup>	2.06	124.95	121.56
1	F	202	LLP	C5-C4-C4'	2.06	124.95	121.56
1	D	202	LLP	OP3-P-OP2	2.05	115.47	107.64
1	Ε	202	LLP	O3-C3-C2	2.03	121.91	117.49

Continued from previous page...



There are no chirality outliers.

All (43) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	202	LLP	C4-C4'-NZ-CE
1	Е	202	LLP	C4-C4'-NZ-CE
1	Е	202	LLP	C-CA-CB-CG
1	В	202	LLP	C4-C4'-NZ-CE
1	С	202	LLP	C4-C4'-NZ-CE
1	D	202	LLP	C4-C4'-NZ-CE
1	F	202	LLP	C4-C4'-NZ-CE
1	F	202	LLP	O-C-CA-CB
1	G	202	LLP	C4-C4'-NZ-CE
1	G	202	LLP	C4-C5-C5'-OP4
1	Н	202	LLP	C5-C4-C4'-NZ
1	Н	202	LLP	C4-C4'-NZ-CE
1	А	202	LLP	CG-CD-CE-NZ
1	В	202	LLP	CG-CD-CE-NZ
1	С	202	LLP	CG-CD-CE-NZ
1	F	202	LLP	CG-CD-CE-NZ
1	Н	202	LLP	C3-C4-C4'-NZ
1	Ε	202	LLP	CA-CB-CG-CD
1	F	202	LLP	CA-CB-CG-CD
1	D	202	LLP	CA-CB-CG-CD
1	G	202	LLP	CE-CD-CG-CB
1	А	202	LLP	CE-CD-CG-CB
1	Ε	202	LLP	C3-C4-C4'-NZ
1	G	202	LLP	CG-CD-CE-NZ
1	Н	202	LLP	CE-CD-CG-CB
1	G	202	LLP	C-CA-CB-CG
1	G	202	LLP	C6-C5-C5'-OP4
1	В	202	LLP	CD-CE-NZ-C4'
1	С	202	LLP	CD-CE-NZ-C4'
1	D	202	LLP	CG-CD-CE-NZ
1	Н	202	LLP	CG-CD-CE-NZ
1	В	202	LLP	N-CA-CB-CG
1	D	202	LLP	C3-C4-C4'-NZ
1	Ε	202	LLP	C5-C4-C4'-NZ
1	Н	202	LLP	CD-CE-NZ-C4'
1	G	$20\overline{2}$	LLP	C5'-OP4-P-OP1
1	Н	202	LLP	CA-CB-CG-CD
1	A	202	LLP	C-CA-CB-CG
1	A	202	LLP	C3-C4-C4'-NZ
1	G	202	LLP	C3-C4-C4'-NZ



$\alpha$ $\cdot$ $\cdot$ $\cdot$	C	•	
Continued	trom	nremous	naae
Contentaca	<i>J</i> 10110	proceeduo	page

Mol	Chain	Res	Type	Atoms
1	F	202	LLP	CE-CD-CG-CB
1	С	202	LLP	N-CA-CB-CG
1	Н	202	LLP	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	F	202	LLP	2	0

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	366/379~(96%)	-0.34	0 100 100	11, 18, 29, 39	0
1	В	366/379~(96%)	-0.21	6 (1%) 72 77	11, 19, 35, 57	0
1	С	362/379~(95%)	-0.04	1 (0%) 94 95	16,31,51,61	0
1	D	366/379~(96%)	-0.23	4 (1%) 80 84	12, 18, 35, 44	0
1	Ε	363/379~(95%)	-0.02	8 (2%) 62 68	14, 29, 54, 69	0
1	F	342/379~(90%)	-0.10	4 (1%) 79 83	16, 28, 45, 55	0
1	G	366/379~(96%)	-0.33	0 100 100	12, 18, 28, 34	0
1	Η	365/379~(96%)	-0.04	6 (1%) 72 77	14, 27, 48, 60	0
All	All	2896/3032~(95%)	-0.16	29 (1%) 82 86	11, 23, 45, 69	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	339	HIS	6.8
1	В	340	VAL	5.8
1	D	343	GLY	4.4
1	Н	320	ALA	3.4
1	D	342	VAL	3.0
1	В	341	SER	2.9
1	Ε	318	LEU	2.9
1	Е	310	PRO	2.6
1	Н	330	GLN	2.5
1	F	328	CYS	2.5
1	D	340	VAL	2.4
1	D	146	GLU	2.4
1	Е	311	ASP	2.4
1	В	342	VAL	2.4
1	Е	285	ILE	2.4
1	Ε	313	TYR	2.3



	9	1	1 0	
Mol	Chain	Res	Type	RSRZ
1	Е	327	PHE	2.2
1	F	308	PHE	2.2
1	Н	342	VAL	2.2
1	F	339	HIS	2.2
1	В	343	GLY	2.2
1	F	340	VAL	2.1
1	Н	364	ALA	2.1
1	С	323	LYS	2.1
1	В	318	LEU	2.1
1	Е	330	GLN	2.0
1	Н	321	ASP	2.0
1	Н	318	LEU	2.0
1	E	131	ALA	2.0

Continued from previous page...

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
1	LLP	F	202	24/25	0.90	0.15	24,26,30,31	0
1	LLP	С	202	24/25	0.93	0.15	26,29,31,31	0
1	LLP	Е	202	24/25	0.93	0.14	25,30,32,33	0
1	LLP	G	202	24/25	0.93	0.12	17,22,23,24	0
1	LLP	А	202	24/25	0.94	0.11	16,22,23,24	0
1	LLP	Н	202	24/25	0.94	0.13	20,23,24,25	0
1	LLP	D	202	24/25	0.96	0.10	16,18,20,20	0
1	LLP	В	202	24/25	0.96	0.10	18,20,22,22	0

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

There are no ligands in this entry.



## 6.5 Other polymers (i)

There are no such residues in this entry.

