



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 21, 2023 – 06:20 AM JST

PDB ID : 7F9M
Title : Crystal structure of the variable region of Plasmodium RIFIN #4 (PF3D7_1000500) in complex with LAIR1 (with T67L, N69S and A77T mutations)
Authors : Xie, Y.; Song, H.; Li, X.; Qi, J.; Gao, G.F.
Deposited on : 2021-07-04
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	
1	B	175	
2	C	112	
2	D	112	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rifin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	158	1169	730	207	226	6	0	0	0
1	B	158	1169	730	207	226	6	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	149	HIS	-	expression tag	UNP A0A143ZWD5
A	150	MET	-	expression tag	UNP A0A143ZWD5
A	151	HIS	-	expression tag	UNP A0A143ZWD5
A	152	HIS	-	expression tag	UNP A0A143ZWD5
A	153	HIS	-	expression tag	UNP A0A143ZWD5
A	154	HIS	-	expression tag	UNP A0A143ZWD5
A	155	HIS	-	expression tag	UNP A0A143ZWD5
A	156	HIS	-	expression tag	UNP A0A143ZWD5
B	149	HIS	-	expression tag	UNP A0A143ZWD5
B	150	MET	-	expression tag	UNP A0A143ZWD5
B	151	HIS	-	expression tag	UNP A0A143ZWD5
B	152	HIS	-	expression tag	UNP A0A143ZWD5
B	153	HIS	-	expression tag	UNP A0A143ZWD5
B	154	HIS	-	expression tag	UNP A0A143ZWD5
B	155	HIS	-	expression tag	UNP A0A143ZWD5
B	156	HIS	-	expression tag	UNP A0A143ZWD5

- Molecule 2 is a protein called Leukocyte-associated immunoglobulin-like receptor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	99	779	487	133	157	2	0	0	0
2	D	99	779	487	133	157	2	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	14	HIS	-	expression tag	UNP Q6GTX8
C	15	MET	-	expression tag	UNP Q6GTX8
C	16	HIS	-	expression tag	UNP Q6GTX8
C	17	HIS	-	expression tag	UNP Q6GTX8
C	18	HIS	-	expression tag	UNP Q6GTX8
C	19	HIS	-	expression tag	UNP Q6GTX8
C	20	HIS	-	expression tag	UNP Q6GTX8
C	21	HIS	-	expression tag	UNP Q6GTX8
C	67	LEU	THR	engineered mutation	UNP Q6GTX8
C	69	SER	ASN	engineered mutation	UNP Q6GTX8
C	77	THR	ALA	engineered mutation	UNP Q6GTX8
C	123	ALA	-	expression tag	UNP Q6GTX8
C	124	ALA	-	expression tag	UNP Q6GTX8
C	125	ALA	-	expression tag	UNP Q6GTX8
D	14	HIS	-	expression tag	UNP Q6GTX8
D	15	MET	-	expression tag	UNP Q6GTX8
D	16	HIS	-	expression tag	UNP Q6GTX8
D	17	HIS	-	expression tag	UNP Q6GTX8
D	18	HIS	-	expression tag	UNP Q6GTX8
D	19	HIS	-	expression tag	UNP Q6GTX8
D	20	HIS	-	expression tag	UNP Q6GTX8
D	21	HIS	-	expression tag	UNP Q6GTX8
D	67	LEU	THR	engineered mutation	UNP Q6GTX8
D	69	SER	ASN	engineered mutation	UNP Q6GTX8
D	77	THR	ALA	engineered mutation	UNP Q6GTX8
D	123	ALA	-	expression tag	UNP Q6GTX8
D	124	ALA	-	expression tag	UNP Q6GTX8
D	125	ALA	-	expression tag	UNP Q6GTX8



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.78Å 73.78Å 345.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.52 – 2.90 47.52 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (47.52-2.90) 99.5 (47.52-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.91Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.254 , 0.310 0.254 , 0.309	Depositor DCC
R_{free} test set	1119 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	93.3	Xtrriage
Anisotropy	0.547	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3896	wwPDB-VP
Average B, all atoms (Å ²)	105.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/1180	0.76	2/1596 (0.1%)
1	B	0.52	1/1180 (0.1%)	0.71	0/1596
2	C	0.56	1/798 (0.1%)	0.72	0/1085
2	D	0.52	0/798	0.69	0/1085
All	All	0.53	2/3956 (0.1%)	0.72	2/5362 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	49	CYS	CB-SG	-5.73	1.72	1.81
1	B	292	LYS	CD-CE	5.39	1.64	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	266	LEU	CA-CB-CG	6.16	129.48	115.30
1	A	283	LEU	CB-CG-CD2	-5.53	101.61	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1169	0	1216	64	0
1	B	1169	0	1216	56	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	779	0	748	26	0
2	D	779	0	748	32	0
All	All	3896	0	3928	168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:62:ARG:NH1	2:D:96:ALA:O	1.65	1.30
2:D:62:ARG:NH2	2:D:98:PRO:O	1.68	1.26
1:A:179:ALA:HA	1:A:309:ILE:HD11	1.24	1.13
2:D:62:ARG:HH22	2:D:98:PRO:N	1.53	1.06
1:B:198:ILE:HD11	1:B:290:ALA:HA	1.49	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/175 (89%)	139 (89%)	14 (9%)	3 (2%)	8	28
1	B	156/175 (89%)	140 (90%)	16 (10%)	0	100	100
2	C	97/112 (87%)	92 (95%)	4 (4%)	1 (1%)	15	45
2	D	97/112 (87%)	92 (95%)	4 (4%)	1 (1%)	15	45
All	All	506/574 (88%)	463 (92%)	38 (8%)	5 (1%)	15	45

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	267	ARG
1	A	221	SER
1	A	266	LEU
2	D	63	GLU
2	C	43	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/137 (90%)	119 (96%)	5 (4%)	31	65
1	B	124/137 (90%)	121 (98%)	3 (2%)	49	79
2	C	89/100 (89%)	85 (96%)	4 (4%)	27	61
2	D	89/100 (89%)	84 (94%)	5 (6%)	21	52
All	All	426/474 (90%)	409 (96%)	17 (4%)	31	65

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	87	ARG
2	D	121	LYS
1	B	285	ARG
2	C	64	SER
2	C	76	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	230	ASN
1	B	295	ASN
1	B	303	GLN
1	A	257	GLN
1	A	214	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/175 (90%)	0.11	4 (2%) 57 55	79, 95, 120, 155	0
1	B	158/175 (90%)	0.12	3 (1%) 66 65	81, 102, 118, 168	0
2	C	99/112 (88%)	0.43	4 (4%) 38 33	81, 100, 143, 210	0
2	D	99/112 (88%)	0.93	16 (16%) 1 1	84, 109, 169, 200	0
All	All	514/574 (89%)	0.33	27 (5%) 26 22	79, 101, 138, 210	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	166	VAL	12.2
2	C	25	ASP	6.3
2	D	26	LEU	6.3
2	D	53	VAL	4.1
2	D	123	ALA	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.