



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2023 – 11:11 AM EST

PDB ID : 8FAA
Title : Crystal structure of Xanthomonas campestris GH35 beta-galactosidase
Authors : Godoy, A.S.; Polikarpov, I.
Deposited on : 2022-11-25
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	
1	B	511	
1	C	511	
1	D	511	
1	E	511	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	511	<p>2% 84% 15%</p>
1	G	511	<p>2% 85% 14%</p>
1	H	511	<p>% 87% 12%</p>
1	I	511	<p>3% 86% 13%</p>
1	J	511	<p>2% 87% 12%</p>
1	K	511	<p>3% 86% 13%</p>
1	L	511	<p>5% 86% 12%</p>
1	M	511	<p>2% 87% 12%</p>
1	N	511	<p>4% 85% 14%</p>
1	O	511	<p>5% 82% 17%</p>
1	P	511	<p>% 87% 12%</p>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 65632 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	510	3921	2491	707	718	5	0	0	0
1	B	510	3921	2491	707	718	5	0	0	0
1	C	510	3921	2491	707	718	5	0	0	0
1	D	510	3921	2491	707	718	5	0	0	0
1	E	510	3921	2491	707	718	5	0	0	0
1	F	510	3921	2491	707	718	5	0	0	0
1	G	510	3921	2491	707	718	5	0	0	0
1	H	510	3921	2491	707	718	5	0	0	0
1	I	510	3921	2491	707	718	5	0	0	0
1	J	510	3921	2491	707	718	5	0	0	0
1	K	510	3921	2491	707	718	5	0	0	0
1	L	510	3921	2491	707	718	5	0	0	0
1	M	510	3921	2491	707	718	5	0	0	0
1	N	510	3921	2491	707	718	5	0	0	0
1	O	510	3921	2491	707	718	5	0	0	0
1	P	510	3921	2491	707	718	5	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP Q8P844
B	2	MET	-	initiating methionine	UNP Q8P844
C	2	MET	-	initiating methionine	UNP Q8P844
D	2	MET	-	initiating methionine	UNP Q8P844
E	2	MET	-	initiating methionine	UNP Q8P844
F	2	MET	-	initiating methionine	UNP Q8P844
G	2	MET	-	initiating methionine	UNP Q8P844
H	2	MET	-	initiating methionine	UNP Q8P844
I	2	MET	-	initiating methionine	UNP Q8P844
J	2	MET	-	initiating methionine	UNP Q8P844
K	2	MET	-	initiating methionine	UNP Q8P844
L	2	MET	-	initiating methionine	UNP Q8P844
M	2	MET	-	initiating methionine	UNP Q8P844
N	2	MET	-	initiating methionine	UNP Q8P844
O	2	MET	-	initiating methionine	UNP Q8P844
P	2	MET	-	initiating methionine	UNP Q8P844

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	242	Total O 242 242	0	0
2	B	157	Total O 157 157	0	0
2	C	186	Total O 186 186	0	0
2	D	186	Total O 186 186	0	0
2	E	200	Total O 200 200	0	0
2	F	203	Total O 203 203	0	0
2	G	196	Total O 196 196	0	0
2	H	188	Total O 188 188	0	0
2	I	215	Total O 215 215	0	0
2	J	198	Total O 198 198	0	0
2	K	185	Total O 185 185	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	L	209	Total 209	O 209	0	0
2	M	129	Total 129	O 129	0	0
2	N	113	Total 113	O 113	0	0
2	O	94	Total 94	O 94	0	0
2	P	195	Total 195	O 195	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-galactosidase



- Molecule 1: Beta-galactosidase




- Molecule 1: Beta-galactosidase









- Molecule 1: Beta-galactosidase




- Molecule 1: Beta-galactosidase




- Molecule 1: Beta-galactosidase





- Molecule 1: Beta-galactosidase









• Molecule 1: Beta-galactosidase




• Molecule 1: Beta-galactosidase





• Molecule 1: Beta-galactosidase







● Molecule 1: Beta-galactosidase




● Molecule 1: Beta-galactosidase




● Molecule 1: Beta-galactosidase





● Molecule 1: Beta-galactosidase







● Molecule 1: Beta-galactosidase



● Molecule 1: Beta-galactosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	128.27Å 162.99Å 235.48Å 90.00° 92.83° 90.00°	Depositor
Resolution (Å)	49.10 – 2.50 49.12 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (49.10-2.50) 99.1 (49.12-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 2.48Å)	Xtrriage
Refinement program	PHENIX v1	Depositor
R, R_{free}	0.211 , 0.251 0.211 , 0.250	Depositor DCC
R_{free} test set	16767 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	30.4	Xtrriage
Anisotropy	0.453	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.020 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	65632	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.71 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.5908e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/4033	0.61	0/5508
1	B	0.34	0/4033	0.59	0/5508
1	C	0.34	0/4033	0.60	0/5508
1	D	0.37	1/4033 (0.0%)	0.63	2/5508 (0.0%)
1	E	0.36	0/4033	0.61	0/5508
1	F	0.37	0/4033	0.61	1/5508 (0.0%)
1	G	0.36	0/4033	0.62	1/5508 (0.0%)
1	H	0.35	0/4033	0.60	0/5508
1	I	0.39	0/4033	0.64	1/5508 (0.0%)
1	J	0.35	0/4033	0.59	0/5508
1	K	0.39	0/4033	0.61	1/5508 (0.0%)
1	L	0.42	4/4033 (0.1%)	0.69	7/5508 (0.1%)
1	M	0.32	0/4033	0.57	0/5508
1	N	0.33	0/4033	0.58	1/5508 (0.0%)
1	O	0.39	2/4033 (0.0%)	0.60	1/5508 (0.0%)
1	P	0.34	0/4033	0.59	0/5508
All	All	0.36	7/64528 (0.0%)	0.61	15/88128 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	F	0	2
All	All	0	3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	4	GLU	CB-CG	-9.03	1.34	1.52
1	L	456	ASP	CB-CG	6.26	1.65	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	458	ARG	CB-CG	5.91	1.68	1.52
1	L	458	ARG	CA-CB	5.80	1.66	1.53
1	D	239	ARG	CG-CD	-5.58	1.38	1.51

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	458	ARG	CD-NE-CZ	9.30	136.62	123.60
1	L	458	ARG	NE-CZ-NH1	9.11	124.86	120.30
1	D	239	ARG	CB-CG-CD	-8.85	88.58	111.60
1	L	472	ASP	CB-CG-OD2	8.73	126.15	118.30
1	L	472	ASP	CB-CG-OD1	-8.48	110.66	118.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	239	ARG	Sidechain
1	F	71	ARG	Sidechain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3921	0	3780	55	1
1	B	3921	0	3780	52	0
1	C	3921	0	3780	42	0
1	D	3921	0	3780	41	1
1	E	3921	0	3780	47	0
1	F	3921	0	3780	56	0
1	G	3921	0	3780	56	0
1	H	3921	0	3780	44	0
1	I	3921	0	3780	71	0
1	J	3921	0	3780	47	0
1	K	3921	0	3780	52	0
1	L	3921	0	3780	74	0
1	M	3921	0	3780	41	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	N	3921	0	3780	55	0
1	O	3921	0	3780	67	0
1	P	3921	0	3780	46	0
2	A	242	0	0	16	4
2	B	157	0	0	17	0
2	C	186	0	0	12	1
2	D	186	0	0	8	1
2	E	200	0	0	20	1
2	F	203	0	0	20	1
2	G	196	0	0	16	1
2	H	188	0	0	12	1
2	I	215	0	0	26	2
2	J	198	0	0	12	0
2	K	185	0	0	15	1
2	L	209	0	0	20	2
2	M	129	0	0	12	0
2	N	113	0	0	23	0
2	O	94	0	0	22	0
2	P	195	0	0	10	1
All	All	65632	0	60480	816	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 816 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:239:ARG:HD2	1:L:458:ARG:HB3	1.27	1.13
1:G:364:ARG:NH2	2:G:1301:HOH:O	1.81	1.06
1:I:141:GLN:NE2	2:I:603:HOH:O	1.89	1.02
1:P:428:ARG:NH2	2:P:602:HOH:O	1.93	1.01
1:B:428:ARG:NH2	2:B:601:HOH:O	1.95	0.98

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:696:HOH:O	2:K:729:HOH:O[2_655]	1.59	0.61
2:A:794:HOH:O	2:G:1474:HOH:O[2_555]	1.96	0.24
2:H:716:HOH:O	2:L:604:HOH:O[1_655]	1.98	0.22

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:LYS:O	1:D:239:ARG:NH2[1_455]	2.03	0.17
2:A:606:HOH:O	2:P:692:HOH:O[1_565]	2.04	0.16

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/511 (99%)	491 (97%)	17 (3%)	0	100	100
1	B	508/511 (99%)	489 (96%)	19 (4%)	0	100	100
1	C	508/511 (99%)	489 (96%)	19 (4%)	0	100	100
1	D	508/511 (99%)	485 (96%)	22 (4%)	1 (0%)	47	68
1	E	508/511 (99%)	491 (97%)	16 (3%)	1 (0%)	47	68
1	F	508/511 (99%)	494 (97%)	14 (3%)	0	100	100
1	G	508/511 (99%)	490 (96%)	18 (4%)	0	100	100
1	H	508/511 (99%)	491 (97%)	17 (3%)	0	100	100
1	I	508/511 (99%)	490 (96%)	16 (3%)	2 (0%)	34	54
1	J	508/511 (99%)	493 (97%)	15 (3%)	0	100	100
1	K	508/511 (99%)	494 (97%)	14 (3%)	0	100	100
1	L	508/511 (99%)	491 (97%)	17 (3%)	0	100	100
1	M	508/511 (99%)	494 (97%)	14 (3%)	0	100	100
1	N	508/511 (99%)	491 (97%)	17 (3%)	0	100	100
1	O	508/511 (99%)	491 (97%)	17 (3%)	0	100	100
1	P	508/511 (99%)	493 (97%)	15 (3%)	0	100	100
All	All	8128/8176 (99%)	7857 (97%)	267 (3%)	4 (0%)	100	100

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	200	ALA
1	I	201	ALA
1	D	239	ARG
1	E	195	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	381/382 (100%)	371 (97%)	10 (3%)	46	72
1	B	381/382 (100%)	370 (97%)	11 (3%)	42	69
1	C	381/382 (100%)	369 (97%)	12 (3%)	40	67
1	D	381/382 (100%)	368 (97%)	13 (3%)	37	63
1	E	381/382 (100%)	366 (96%)	15 (4%)	32	57
1	F	381/382 (100%)	367 (96%)	14 (4%)	34	60
1	G	381/382 (100%)	376 (99%)	5 (1%)	69	87
1	H	381/382 (100%)	370 (97%)	11 (3%)	42	69
1	I	381/382 (100%)	372 (98%)	9 (2%)	49	74
1	J	381/382 (100%)	372 (98%)	9 (2%)	49	74
1	K	381/382 (100%)	372 (98%)	9 (2%)	49	74
1	L	381/382 (100%)	369 (97%)	12 (3%)	40	67
1	M	381/382 (100%)	371 (97%)	10 (3%)	46	72
1	N	381/382 (100%)	374 (98%)	7 (2%)	59	81
1	O	381/382 (100%)	370 (97%)	11 (3%)	42	69
1	P	381/382 (100%)	372 (98%)	9 (2%)	49	74
All	All	6096/6112 (100%)	5929 (97%)	167 (3%)	44	71

5 of 167 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	429	LEU
1	N	253	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	180	ARG
1	M	14	ARG
1	O	148	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	N	210	GLN
1	O	329	HIS
1	N	211	GLN
1	O	103	GLN
1	P	141	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	510/511 (99%)	-0.25	5 (0%) 82 84	15, 25, 40, 66	0
1	B	510/511 (99%)	-0.05	10 (1%) 65 68	21, 31, 48, 68	0
1	C	510/511 (99%)	-0.18	7 (1%) 75 77	19, 28, 43, 69	0
1	D	510/511 (99%)	-0.17	13 (2%) 57 61	19, 27, 46, 71	0
1	E	510/511 (99%)	-0.28	9 (1%) 68 71	18, 25, 43, 64	0
1	F	510/511 (99%)	-0.19	8 (1%) 72 74	16, 27, 46, 70	0
1	G	510/511 (99%)	-0.19	9 (1%) 68 71	17, 26, 45, 72	0
1	H	510/511 (99%)	-0.22	4 (0%) 86 87	20, 29, 43, 68	0
1	I	510/511 (99%)	-0.09	13 (2%) 57 61	17, 29, 47, 72	0
1	J	510/511 (99%)	-0.11	11 (2%) 62 65	19, 29, 49, 69	0
1	K	510/511 (99%)	-0.11	16 (3%) 49 52	19, 30, 45, 74	0
1	L	510/511 (99%)	0.01	24 (4%) 31 33	18, 28, 50, 78	0
1	M	510/511 (99%)	0.00	12 (2%) 59 62	25, 37, 55, 76	0
1	N	510/511 (99%)	0.22	19 (3%) 41 45	25, 42, 59, 84	0
1	O	510/511 (99%)	0.30	24 (4%) 31 33	27, 39, 56, 78	0
1	P	510/511 (99%)	-0.17	6 (1%) 79 80	19, 29, 46, 69	0
All	All	8160/8176 (99%)	-0.09	190 (2%) 60 63	15, 30, 50, 84	0

The worst 5 of 190 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	471	VAL	10.8
1	L	472	ASP	10.7
1	K	472	ASP	8.7
1	L	457	THR	8.2
1	D	454	ALA	8.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.