



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 12:19 PM BST

PDB ID : 1FC1
Title : CRYSTALLOGRAPHIC REFINEMENT AND ATOMIC MODELS OF A HUMAN FC FRAGMENT AND ITS COMPLEX WITH FRAGMENT B OF PROTEIN A FROM STAPHYLOCOCCUS AUREUS AT 2.9-AND 2.8-ANGSTROMS RESOLUTION
Authors : Deisenhofer, J.
Deposited on : 1981-05-21
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| Clashscore | 141614 | 2172 (2.90-2.90) |
| Ramachandran outliers | 138981 | 2115 (2.90-2.90) |
| Sidechain outliers | 138945 | 2117 (2.90-2.90) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 224 | |
| 1 | B | 224 | |
| 2 | C | 9 | |
| 2 | D | 9 | |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 2 | MAN | C | 3 | X | - | - | - |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|------------|-------------|--------------|------------|------------------|-----------------|----------------|-------------------------|
| 2 | FUC | C | 9 | X | - | - | - |
| 2 | MAN | D | 3 | X | - | - | - |
| 2 | FUC | D | 9 | X | - | - | - |

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 3532 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FC FRAGMENT.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|---------------|-----------|----------|----------|--------|---------|---------|-------|
| | | | Total | C | N | O | S | | | |
| 1 | A | 207 | Total 1656 | C 1054 | N 282 | O 313 | S 7 | 60 | 0 | 1 |
| 1 | B | 207 | Total 1656 | C 1054 | N 282 | O 313 | S 7 | 217 | 0 | 1 |

There are 14 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| A | 272 | GLN | GLU | conflict | UNP P01857 |
| A | 283 | GLN | GLU | conflict | UNP P01857 |
| A | 294 | GLN | GLU | conflict | UNP P01857 |
| A | 312 | ASN | ASP | conflict | UNP P01857 |
| A | 315 | ASP | ASN | conflict | UNP P01857 |
| A | 356 | GLU | ASP | conflict | UNP P01857 |
| A | 358 | MET | LEU | conflict | UNP P01857 |
| B | 272 | GLN | GLU | conflict | UNP P01857 |
| B | 283 | GLN | GLU | conflict | UNP P01857 |
| B | 294 | GLN | GLU | conflict | UNP P01857 |
| B | 312 | ASN | ASP | conflict | UNP P01857 |
| B | 315 | ASP | ASN | conflict | UNP P01857 |
| B | 356 | GLU | ASP | conflict | UNP P01857 |
| B | 358 | MET | LEU | conflict | UNP P01857 |

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|------------|--------------|-----------------|--------------|----|---|----|----------------|----------------|--------------|
| 2 | C | 9 | Total | C | N | O | 18 | 0 | 0 |
| | | | 110 | 62 | 4 | 44 | | | |
| 2 | D | 9 | Total | C | N | O | 55 | 0 | 0 |
| | | | 110 | 62 | 4 | 44 | | | |

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: FC FRAGMENT



- Molecule 1: FC FRAGMENT




- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyr

anose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alph
a-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

| Property | Value | Source |
|--|---|-----------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 80.40Å 146.40Å 50.40Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | (Not available) – 2.90 | Depositor |
| % Data completeness (in resolution range) | (Not available) ((Not available)-2.90) | Depositor |
| R_{merge} | (Not available) | Depositor |
| R_{sym} | (Not available) | Depositor |
| Refinement program | unknown | Depositor |
| R, R_{free} | (Not available) , (Not available) | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| Total number of atoms | 3532 | wwPDB-VP |
| Average B, all atoms (Å ²) | 21.0 | wwPDB-VP |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, GAL, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------------|-------------|---------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | A | 1.01 | 4/1702 (0.2%) | 1.22 | 2/2318 (0.1%) |
| 1 | B | 1.01 | 4/1702 (0.2%) | 1.21 | 1/2318 (0.0%) |
| All | All | 1.01 | 8/3404 (0.2%) | 1.22 | 3/4636 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | A | 0 | 17 |
| 1 | B | 0 | 22 |
| All | All | 0 | 39 |

The worst 5 of 8 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 1 | B | 381 | TRP | NE1-CE2 | -7.99 | 1.27 | 1.37 |
| 1 | A | 277 | TRP | NE1-CE2 | -7.94 | 1.27 | 1.37 |
| 1 | B | 277 | TRP | NE1-CE2 | -7.76 | 1.27 | 1.37 |
| 1 | B | 313 | TRP | NE1-CE2 | -7.71 | 1.27 | 1.37 |
| 1 | A | 417 | TRP | NE1-CE2 | -7.63 | 1.27 | 1.37 |

All (3) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 1 | B | 401 | ASP | CB-CA-C | -5.71 | 98.98 | 110.40 |
| 1 | A | 297 | ASN | N-CA-CB | -5.52 | 100.66 | 110.60 |
| 1 | A | 285 | HIS | CA-CB-CG | -5.42 | 104.39 | 113.60 |

There are no chirality outliers.

5 of 39 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 1 | A | 259 | VAL | Mainchain |
| 1 | A | 276 | ASN | Sidechain |
| 1 | A | 280 | ASP | Mainchain |
| 1 | A | 287 | ALA | Mainchain |
| 1 | A | 309 | LEU | Mainchain |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 1656 | 0 | 1629 | 51 | 4 |
| 1 | B | 1656 | 0 | 1629 | 36 | 1 |
| 2 | C | 110 | 0 | 94 | 5 | 0 |
| 2 | D | 110 | 0 | 94 | 1 | 0 |
| All | All | 3532 | 0 | 3446 | 86 | 4 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|------------------|--------------------------|-------------------|
| 1:B:274:LYS:HB3 | 1:B:324:SER:HB2 | 1.47 | 0.96 |
| 1:A:297:ASN:HB3 | 1:A:299:THR:HG22 | 1.61 | 0.81 |
| 1:A:328:LEU:HD21 | 1:A:332:ILE:HG13 | 1.64 | 0.77 |
| 1:A:250:THR:HG22 | 1:A:257:PRO:HB3 | 1.68 | 0.76 |
| 1:A:296:TYR:HD2 | 2:C:1:NAG:H62 | 1.52 | 0.75 |

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------------|--------------------------|-------------------|
| 1:A:389:ASN:OD1 | 1:B:389:ASN:ND2[4_566] | 1.69 | 0.51 |

Continued on next page...

Continued from previous page...

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------------|--------------------------|-------------------|
| 1:A:268:HIS:CG | 1:A:285:HIS:NE2[2_674] | 1.81 | 0.39 |
| 1:A:268:HIS:ND1 | 1:A:285:HIS:NE2[2_674] | 2.01 | 0.19 |
| 1:A:268:HIS:CB | 1:A:285:HIS:NE2[2_674] | 2.16 | 0.04 |

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|---------|----------|-------------|----|
| 1 | A | 205/224 (92%) | 179 (87%) | 18 (9%) | 8 (4%) | 3 | 12 |
| 1 | B | 205/224 (92%) | 179 (87%) | 17 (8%) | 9 (4%) | 2 | 10 |
| All | All | 410/448 (92%) | 358 (87%) | 35 (8%) | 17 (4%) | 3 | 11 |

5 of 17 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 297 | ASN |
| 1 | B | 268 | HIS |
| 1 | B | 292 | ARG |
| 1 | B | 297 | ASN |
| 1 | A | 285 | HIS |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1 | A | 193/207 (93%) | 186 (96%) | 7 (4%) | 35 | 69 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1 | B | 193/207 (93%) | 186 (96%) | 7 (4%) | 35 | 69 |
| All | All | 386/414 (93%) | 372 (96%) | 14 (4%) | 35 | 69 |

5 of 14 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 434 | ASN |
| 1 | B | 284 | VAL |
| 1 | B | 390 | ASN |
| 1 | A | 425 | CYS |
| 1 | B | 376 | ASP |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 438 | GLN |
| 1 | B | 283 | GLN |
| 1 | B | 418 | GLN |
| 1 | A | 421 | ASN |
| 1 | B | 419 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 2 | NAG | C | 1 | 1,2 | 14,14,15 | 0.72 | 0 | 17,19,21 | 1.79 | 3 (17%) |
| 2 | NAG | C | 2 | 2 | 14,14,15 | 0.99 | 0 | 17,19,21 | 1.55 | 2 (11%) |
| 2 | MAN | C | 3 | 2 | 11,11,12 | 0.52 | 0 | 15,15,17 | 1.48 | 2 (13%) |
| 2 | MAN | C | 4 | 2 | 11,11,12 | 0.56 | 0 | 15,15,17 | 1.30 | 1 (6%) |
| 2 | NAG | C | 5 | 2 | 14,14,15 | 0.95 | 0 | 17,19,21 | 1.66 | 1 (5%) |
| 2 | GAL | C | 6 | 2 | 11,11,12 | 0.55 | 0 | 15,15,17 | 1.64 | 4 (26%) |
| 2 | MAN | C | 7 | 2 | 11,11,12 | 0.76 | 0 | 15,15,17 | 1.54 | 1 (6%) |
| 2 | NAG | C | 8 | 2 | 14,14,15 | 0.79 | 0 | 17,19,21 | 1.82 | 1 (5%) |
| 2 | FUC | C | 9 | 2 | 10,10,11 | 0.78 | 0 | 14,14,16 | 1.23 | 1 (7%) |
| 2 | NAG | D | 1 | 1,2 | 14,14,15 | 0.76 | 0 | 17,19,21 | 1.51 | 2 (11%) |
| 2 | NAG | D | 2 | 2 | 14,14,15 | 0.92 | 0 | 17,19,21 | 1.38 | 1 (5%) |
| 2 | MAN | D | 3 | 2 | 11,11,12 | 0.60 | 0 | 15,15,17 | 1.81 | 2 (13%) |
| 2 | MAN | D | 4 | 2 | 11,11,12 | 0.62 | 0 | 15,15,17 | 1.54 | 1 (6%) |
| 2 | NAG | D | 5 | 2 | 14,14,15 | 0.83 | 0 | 17,19,21 | 1.67 | 1 (5%) |
| 2 | GAL | D | 6 | 2 | 11,11,12 | 0.60 | 0 | 15,15,17 | 1.30 | 1 (6%) |
| 2 | MAN | D | 7 | 2 | 11,11,12 | 0.83 | 1 (9%) | 15,15,17 | 1.60 | 1 (6%) |
| 2 | NAG | D | 8 | 2 | 14,14,15 | 0.92 | 1 (7%) | 17,19,21 | 1.66 | 1 (5%) |
| 2 | FUC | D | 9 | 2 | 10,10,11 | 0.74 | 0 | 14,14,16 | 1.19 | 1 (7%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 2 | NAG | C | 1 | 1,2 | - | 4/6/23/26 | 0/1/1/1 |
| 2 | NAG | C | 2 | 2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | MAN | C | 3 | 2 | 1/1/4/5 | 0/2/19/22 | 0/1/1/1 |
| 2 | MAN | C | 4 | 2 | - | 0/2/19/22 | 0/1/1/1 |
| 2 | NAG | C | 5 | 2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | GAL | C | 6 | 2 | - | 1/2/19/22 | 0/1/1/1 |
| 2 | MAN | C | 7 | 2 | - | 0/2/19/22 | 0/1/1/1 |
| 2 | NAG | C | 8 | 2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | FUC | C | 9 | 2 | 2/2/4/5 | - | 0/1/1/1 |
| 2 | NAG | D | 1 | 1,2 | - | 4/6/23/26 | 0/1/1/1 |
| 2 | NAG | D | 2 | 2 | - | 1/6/23/26 | 0/1/1/1 |
| 2 | MAN | D | 3 | 2 | 1/1/4/5 | 0/2/19/22 | 0/1/1/1 |
| 2 | MAN | D | 4 | 2 | - | 0/2/19/22 | 0/1/1/1 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|-----------|---------|
| 2 | NAG | D | 5 | 2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | GAL | D | 6 | 2 | - | 1/2/19/22 | 0/1/1/1 |
| 2 | MAN | D | 7 | 2 | - | 0/2/19/22 | 0/1/1/1 |
| 2 | NAG | D | 8 | 2 | - | 0/6/23/26 | 0/1/1/1 |
| 2 | FUC | D | 9 | 2 | 2/2/4/5 | - | 0/1/1/1 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|------|-------------|----------|
| 2 | D | 8 | NAG | C1-C2 | 2.37 | 1.55 | 1.52 |
| 2 | D | 7 | MAN | C2-C3 | 2.24 | 1.55 | 1.52 |

The worst 5 of 27 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|------|-------------|----------|
| 2 | C | 8 | NAG | C1-O5-C5 | 7.12 | 121.84 | 112.19 |
| 2 | D | 8 | NAG | C1-O5-C5 | 5.91 | 120.20 | 112.19 |
| 2 | D | 5 | NAG | C1-O5-C5 | 5.70 | 119.91 | 112.19 |
| 2 | D | 3 | MAN | C1-O5-C5 | 5.66 | 119.86 | 112.19 |
| 2 | C | 5 | NAG | C1-O5-C5 | 5.42 | 119.54 | 112.19 |

5 of 6 chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 2 | C | 9 | FUC | C5 |
| 2 | C | 9 | FUC | C1 |
| 2 | D | 9 | FUC | C5 |
| 2 | D | 9 | FUC | C1 |
| 2 | D | 3 | MAN | C1 |

5 of 11 torsion outliers are listed below:


| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------------|
| 2 | C | 1 | NAG | C8-C7-N2-C2 |
| 2 | C | 1 | NAG | O7-C7-N2-C2 |
| 2 | D | 1 | NAG | C8-C7-N2-C2 |
| 2 | D | 1 | NAG | O7-C7-N2-C2 |
| 2 | D | 1 | NAG | O5-C5-C6-O6 |


There are no ring outliers.

3 monomers are involved in 6 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 2 | C | 1 | NAG | 2 | 0 |
| 2 | D | 6 | GAL | 1 | 0 |
| 2 | C | 6 | GAL | 3 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.