

wwPDB X-ray Structure Validation Summary Report (i)

May 15, 2020 - 07:04 pm BST

PDB ID	:	5FDX
Title	:	Structure of DDR1 receptor tyrosine kinase in complex with D2164 inhibitor
		at 2.65 Angstroms resolution.
Authors	:	Bartual, S.G.; Pinkas, D.M.; Wang, Z.; Ding, K.; Mahajan, P.; Kupinska,
		K.; Mukhopadhyay, S.; Strain-Damerell, C.; Chalk, R.; Borkowska, O.; Talon,
		R.; Kopec, J.; Williams, E.; Tallant, C.; Chaikuad, A.; Sorell, F.; Newman, J.;
		Burgess-Brown, N.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.M.; Bountra,
		C.; Bullock, A.; Structural Genomics Consortium (SGC)
Deposited on	:	2015-12-16
$\operatorname{Resolution}$:	2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac		20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)	:	Parkinson et al. (1996) 2.11
vandation ripenne (wwrDD-vr)	•	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R_{free}	130704	1332(2.68-2.64)
Clashscore	141614	1374(2.68-2.64)
Ramachandran outliers	138981	1349(2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	337	% • 74%	9%	•	17%
1	В	337	73%	8%		19%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	280	Total	С	Ν	Ο	S	0	0	0
	200	2194	1409	383	384	18	0	0	0	
1	р	274	Total	С	Ν	Ο	S	0	0	0
	D	274	2117	1360	364	376	17			0

• Molecule 1 is a protein called Epithelial discoidin domain-containing receptor 1.

Chain	Residue	Modelled	Actual	Comment	Reference
А	577	MET	-	initiating methionine	UNP Q08345
А	578	GLY	_	expression tag	UNP Q08345
А	579	HIS	-	expression tag	UNP Q08345
А	580	HIS	-	expression tag	UNP Q08345
А	581	HIS	-	expression tag	UNP Q08345
А	582	HIS	-	expression tag	UNP Q08345
А	583	HIS	-	expression tag	UNP Q08345
А	584	HIS	-	expression tag	UNP Q08345
А	585	SER	-	expression tag	UNP Q08345
А	586	SER	-	expression tag	UNP Q08345
А	587	GLY	-	expression tag	UNP Q08345
А	588	VAL	-	expression tag	UNP Q08345
А	589	ASP	-	expression tag	UNP Q08345
А	590	LEU	-	expression tag	UNP Q08345
А	591	GLY	-	expression tag	UNP Q08345
А	592	THR	-	expression tag	UNP Q08345
А	593	GLU	-	expression tag	UNP Q08345
А	594	ASN	-	expression tag	UNP Q08345
А	595	LEU	_	expression tag	UNP Q08345
А	596	TYR	-	expression tag	UNP Q08345
А	597	PHE	-	expression tag	UNP Q08345
A	598	GLN	-	expression tag	UNP Q08345
А	599	SER	-	expression tag	UNP Q08345
А	600	MET	-	expression tag	UNP Q08345
В	577	MET	-	initiating methionine	UNP Q08345

There are 48 discrepancies between the modelled and reference sequences:


Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
В	578	GLY	-	expression tag	UNP Q08345
В	579	HIS	-	expression tag	UNP Q08345
В	580	HIS	-	expression tag	UNP Q08345
В	581	HIS	-	expression tag	UNP Q08345
В	582	HIS	-	expression tag	UNP Q08345
В	583	HIS	-	expression tag	UNP Q08345
В	584	HIS	-	expression tag	UNP Q08345
В	585	SER	-	expression tag	UNP Q08345
В	586	SER	-	expression tag	UNP Q08345
В	587	GLY	-	expression tag	UNP Q08345
В	588	VAL	-	expression tag	UNP Q08345
В	589	ASP	-	expression tag	UNP Q08345
В	590	LEU	-	expression tag	UNP Q08345
В	591	GLY	-	expression tag	UNP Q08345
В	592	THR	-	expression tag	UNP Q08345
В	593	GLU	-	expression tag	UNP Q08345
В	594	ASN	-	expression tag	UNP Q08345
В	595	LEU	-	expression tag	UNP Q08345
В	596	TYR	-	expression tag	UNP Q08345
В	597	PHE	-	expression tag	UNP Q08345
В	598	GLN	-	expression tag	UNP Q08345
В	599	SER	-	expression tag	UNP Q08345
В	600	MET	-	expression tag	UNP Q08345

Continued from previous page...

• Molecule 2 is 3-[(4-methylpiperazin-1-yl)methyl]- {N}-[(4 {R})-4-methyl-2-pyrimidin-5-yl-3,4-dihydro-1 {H}-isoquinolin-7-yl]-5-(trifluoromethyl)benzamide (three-letter code: 5X1) (formula: $C_{28}H_{31}F_3N_6O$).








Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
0	Λ	1	Total	С	F	Ν	Ο	0	0
		1	38	28	3	6	1	0	
0	р	D 1	Total	С	F	Ν	Ο	0	0
	D		38	28	3	6	1	0	


• Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
3	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 4 2 2 \end{array}$	0	0

• Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} \text{C} \text{O} \\ 7 4 3 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	75	Total O 75 75	0	0
5	В	66	Total O 66 66	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epithelial discoidin domain-containing receptor 1



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	45.55Å 120.34 Å 63.02 Å	Depositor
a, b, c, α , β , γ	90.00° 94.87° 90.00°	Depositor
Resolution (Å)	43.44 - 2.65	Depositor
Resolution (A)	43.44 - 2.65	EDS
% Data completeness	98.4 (43.44-2.65)	Depositor
(in resolution range)	98.4(43.44-2.65)	EDS
R _{merge}	0.09	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.38 (at 2.65 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
D D.	0.194 , 0.273	Depositor
R, R_{free}	0.214 , 0.294	DCC
R_{free} test set	961 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	43.1	Xtriage
Anisotropy	0.719	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , 71.8	EDS
L-test for twinning ²	$ \langle L \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4569	wwPDB-VP
Average B, all atoms $(Å^2)$	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO, $5\mathrm{X1}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.52	0/2245	0.69	0/3042	
1	В	0.50	0/2166	0.65	0/2936	
All	All	0.51	0/4411	0.67	0/5978	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2194	0	2134	14	0
1	В	2117	0	2025	10	0
2	А	38	0	0	0	0
2	В	38	0	0	0	0
3	А	16	0	24	0	0
3	В	4	0	6	0	0
4	А	7	0	10	0	0
4	В	14	0	20	2	0
5	А	75	0	0	0	0
5	В	66	0	0	0	0
All	All	4569	0	4219	24	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:800:GLN:CB	1:A:801:GLY:HA3	2.04	0.88
1:B:799:VAL:HG13	1:B:802:ARG:HB2	1.72	0.70
1:A:882:GLU:HA	1:A:885:LEU:HD12	1.81	0.61
1:A:800:GLN:CB	1:A:801:GLY:CA	2.79	0.59
1:A:799:VAL:HA	1:A:802:ARG:HH11	1.70	0.57

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	274/337~(81%)	257 (94%)	16 (6%)	1 (0%)	34 48
1	В	268/337~(80%)	256~(96%)	11 (4%)	1 (0%)	34 48
All	All	542/674~(80%)	513~(95%)	27~(5%)	2~(0%)	34 48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	799	VAL
1	В	631	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	224/288~(78%)	212~(95%)	12~(5%)	22 34		
1	В	211/288 (73%)	200~(95%)	11 (5%)	23 36		
All	All	435/576~(76%)	412 (95%)	23~(5%)	22 35		

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

 $5~{\rm of}~23$ residues with a non-rotameric side chain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	903	ARG
1	В	664	ASN
1	В	882	GLU
1	А	907	GLU
1	В	670	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	716	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	B	ond leng	gths	B	ond ang	gles
	Type	Cham	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	А	1005	-	3,3,3	0.48	0	2,2,2	0.20	0
3	EDO	А	1004	-	3,3,3	0.38	0	2,2,2	0.59	0
2	5X1	В	1001	-	42,42,42	2.60	13 (30%)	$57,\!61,\!61$	1.99	16 (28%)
3	EDO	В	1002	-	3,3,3	0.42	0	2,2,2	0.28	0
3	EDO	А	1003	-	3,3,3	0.53	0	2,2,2	0.31	0
4	PEG	В	1003	-	$6,\!6,\!6$	0.68	0	$5,\!5,\!5$	0.33	0
2	5X1	А	1001	-	42,42,42	2.74	13 (30%)	57,61,61	1.75	15 (26%)
4	PEG	А	1006	-	$6,\!6,\!6$	0.58	0	$5,\!5,\!5$	0.19	0
4	PEG	В	1004	-	$6,\!6,\!6$	0.60	0	$5,\!5,\!5$	0.28	0
3	EDO	А	1002	-	3,3,3	0.45	0	2,2,2	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	А	1005	-	-	1/1/1/1	-
3	EDO	А	1004	-	-	0/1/1/1	-
2	5X1	В	1001	-	-	2/22/44/44	0/5/5/5
3	EDO	В	1002	-	-	1/1/1/1	-
3	EDO	А	1003	-	-	0/1/1/1	-
4	PEG	В	1003	-	-	2/4/4/4	-
2	5X1	A	1001	-	-	2/22/44/44	0/5/5/5
4	PEG	А	1006	-	-	3/4/4/4	-
4	PEG	В	1004	-	-	3/4/4/4	-
3	EDO	А	1002	-	_	1/1/1/1	_

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	А	1001	5X1	CAF-CBE	10.43	1.53	1.39
2	В	1001	5X1	CAF-CBE	9.96	1.52	1.39
2	А	1001	5X1	CAM-CBC	8.09	1.53	1.39
2	В	1001	5X1	CAM-CBC	7.10	1.51	1.39

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
2	А	1001	5X1	CAX-NAW	5.24	1.49	1.35

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	1001	5X1	CAS-NBI-CAT	7.21	122.86	112.59
2	В	1001	5X1	CBC-CAS-NBI	-4.43	107.50	112.09
2	А	1001	5X1	CAS-NBI-CAT	4.14	118.49	112.59
2	А	1001	5X1	CBC-CAS-NBI	-4.13	107.81	112.09
2	А	1001	5X1	CAH-NAU-CAG	4.13	121.08	115.80

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	1001	5X1	CAZ-CAR-NBH-CAQ
2	В	1001	5X1	CAZ-CAR-NBH-CAP
4	А	1006	PEG	O1-C1-C2-O2
4	В	1004	PEG	O2-C3-C4-O4
3	В	1002	EDO	O1-C1-C2-O2

There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	1004	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ >2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	280/337~(83%)	-0.23	2 (0%) 87 87	30,53,81,98	0
1	В	274/337~(81%)	-0.01	8 (2%) 51 48	39, 58, 102, 132	0
All	All	554/674~(82%)	-0.12	10 (1%) 68 65	30, 56, 92, 132	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	692	VAL	3.8
1	В	735	GLY	3.2
1	А	621	PHE	3.0
1	В	773	LEU	2.7
1	В	652	VAL	2.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	\mathbf{RSR}	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
4	PEG	В	1004	7/7	0.77	0.33	76,79,80,81	0
4	PEG	А	1006	7/7	0.80	0.16	68,71,72,72	0
3	EDO	В	1002	4/4	0.83	0.17	$61,\!62,\!65,\!66$	0
3	EDO	А	1005	4/4	0.85	0.15	$65,\!66,\!68,\!69$	0
3	EDO	А	1003	4/4	0.90	0.17	45,47,52,55	0
4	PEG	В	1003	7/7	0.90	0.20	$48,\!55,\!59,\!59$	0
2	5X1	В	1001	38/38	0.93	0.21	41,46,58,59	0
3	EDO	А	1004	4/4	0.96	0.21	45,45,48,51	0
2	5X1	А	1001	38/38	0.96	0.17	$31,\!40,\!48,\!48$	0
3	EDO	А	1002	4/4	0.96	0.14	$36,\!38,\!39,\!40$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

