



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 7, 2020 – 01:58 am BST

PDB ID : 6FDX
Title : Crystal structure of T. brucei PDE-B1 catalytic domain with inhibitor NPD-1086
Authors : Singh, A.K.; Brown, D.G.
Deposited on : 2017-12-27
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	 3% 81% 9% • 8%
1	B	360	 3% 82% 9% • 8%

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5518 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	333	2636	1673	445	500	18	0	0	0
1	B	332	2624	1667	441	498	18	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	559	GLY	-	expression tag	UNP Q8WQX9
A	560	SER	-	expression tag	UNP Q8WQX9
A	561	HIS	-	expression tag	UNP Q8WQX9
A	562	MET	-	expression tag	UNP Q8WQX9
A	563	ALA	-	expression tag	UNP Q8WQX9
A	564	SER	-	expression tag	UNP Q8WQX9
B	559	GLY	-	expression tag	UNP Q8WQX9
B	560	SER	-	expression tag	UNP Q8WQX9
B	561	HIS	-	expression tag	UNP Q8WQX9
B	562	MET	-	expression tag	UNP Q8WQX9
B	563	ALA	-	expression tag	UNP Q8WQX9
B	564	SER	-	expression tag	UNP Q8WQX9


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is (4aS,8aR)-2-[1-(2-aminoquinazolin-4-yl)piperidin-4-yl]-4-(3,4-dimethoxyphenyl)-1,2,4a,5,8,8a-hexahydrophthalazin-1-one (three-letter code: D5Z) (formula: C₂₉H₃₂N₆O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 38 29 6 3	0	0
4	B	1	Total C N O 38 29 6 3	0	0

- Molecule 5 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 3 1 2	0	0
5	A	1	Total C O 3 1 2	0	0
5	A	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0
5	B	1	Total C O 3 1 2	0	0

- Molecule 6 is GUANIDINE (three-letter code: GAI) (formula: CH₅N₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N 4 1 3	0	0
6	A	1	Total C N 4 1 3	0	0
6	B	1	Total C N 4 1 3	0	0
6	B	1	Total C N 4 1 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	76	Total O 76 76	0	0
7	B	68	Total O 68 68	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphodiesterase



- Molecule 1: Phosphodiesterase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.10Å 116.39Å 68.31Å 90.00° 108.67° 90.00°	Depositor
Resolution (Å)	57.10 – 2.31 54.05 – 2.31	Depositor EDS
% Data completeness (in resolution range)	97.1 (57.10-2.31) 97.1 (54.05-2.31)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.32Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.169 , 0.232 0.183 , 0.225	Depositor DCC
R_{free} test set	2602 reflections (7.22%)	wwPDB-VP
Wilson B-factor (Å ²)	51.5	Xtrriage
Anisotropy	0.613	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5518	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, FMT, GAI, D5Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.73	0/2687	0.80	0/3634
1	B	0.73	0/2675	0.79	1/3620 (0.0%)
All	All	0.73	0/5362	0.80	1/7254 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	804	SER	C-N-CA	5.19	134.68	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2636	0	2598	30	0
1	B	2624	0	2585	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	38	0	0	0	0
4	B	38	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	9	0	3	0	0
5	B	9	0	3	0	0
6	A	8	0	8	0	0
6	B	8	0	8	0	0
7	A	76	0	0	1	0
7	B	68	0	0	0	0
All	All	5518	0	5205	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:608:THR:HG22	1:A:687:ARG:HH21	1.21	1.01
1:A:589:ILE:O	1:A:593:GLU:HG3	1.69	0.91
1:A:608:THR:HG22	1:A:687:ARG:NH2	1.89	0.87
1:A:586:VAL:HG22	1:A:587:THR:HG23	1.61	0.83
1:A:666:VAL:HG22	1:A:667:PRO:CD	2.16	0.75
1:A:666:VAL:HG22	1:A:667:PRO:HD2	1.70	0.72
1:B:801:ASP:HB3	1:B:804:SER:HB2	1.73	0.71
1:B:602:LEU:N	1:B:602:LEU:HD22	2.10	0.66
1:B:602:LEU:N	1:B:603:PRO:CD	2.60	0.64
1:A:602:LEU:HD12	1:A:602:LEU:O	1.99	0.63
1:B:643:GLN:N	1:B:643:GLN:OE1	2.23	0.58
1:B:643:GLN:H	1:B:643:GLN:CD	2.00	0.58
1:A:666:VAL:HG22	1:A:667:PRO:HD3	1.88	0.56
1:A:605:PHE:CD1	1:A:605:PHE:N	2.73	0.56
1:A:668:TYR:CD1	1:A:840:VAL:HG13	2.42	0.55
1:A:709:HIS:O	7:A:1101:HOH:O	2.18	0.55
1:A:753:LEU:O	1:A:759:ASP:OD2	2.25	0.55
1:B:601:GLU:C	1:B:602:LEU:HD22	2.26	0.54
1:B:633:TYR:CZ	1:B:637:LEU:HD11	2.43	0.54
1:A:590:THR:HG23	1:A:592:VAL:H	1.75	0.51
1:A:759:ASP:OD1	1:A:762:ASP:N	2.45	0.49
1:B:605:PHE:N	1:B:605:PHE:CD1	2.80	0.49
1:A:586:VAL:O	1:A:586:VAL:HG13	2.12	0.49
1:A:771:LEU:O	1:A:771:LEU:HD23	2.14	0.48
1:B:641:LEU:H	1:B:642:PRO:CD	2.27	0.48
1:B:641:LEU:N	1:B:642:PRO:CD	2.78	0.47
1:B:704:ILE:HD12	1:B:760:VAL:HG21	1.95	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:591:LYS:HA	1:A:594:ARG:HB3	1.97	0.46
1:B:602:LEU:HB2	1:B:634:ARG:NH2	2.31	0.46
1:A:831:ASP:OD1	1:A:834:ARG:NH1	2.49	0.46
1:A:608:THR:CG2	1:A:687:ARG:NH2	2.73	0.46
1:B:602:LEU:N	1:B:603:PRO:HD2	2.30	0.46
1:A:650:GLU:O	1:A:654:ASN:ND2	2.49	0.45
1:B:663:TYR:OH	1:B:674:VAL:HG11	2.17	0.45
1:B:852:LYS:HG3	1:B:858:VAL:HG21	1.99	0.44
1:B:883:ALA:N	1:B:884:PRO:HD2	2.33	0.44
1:B:637:LEU:HD21	1:B:649:ASP:OD1	2.19	0.42
1:A:774:ARG:HA	1:A:774:ARG:HD2	1.60	0.42
1:B:680:THR:HG21	1:B:824:SER:CB	2.50	0.42
1:A:816:ILE:HD13	1:A:816:ILE:HA	1.91	0.41
1:B:795:PHE:CG	1:B:816:ILE:HG13	2.55	0.41
1:B:915:LEU:HA	1:B:915:LEU:HD23	1.78	0.41
1:A:834:ARG:HD2	1:B:853:GLU:HG2	2.03	0.41
1:A:671:PHE:O	1:A:675:VAL:HG23	2.21	0.41
1:B:878:ILE:HG23	1:B:904:ILE:CG2	2.50	0.41
1:A:589:ILE:N	1:A:589:ILE:HD13	2.35	0.41
1:A:697:LEU:O	1:A:701:VAL:HG23	2.21	0.41
1:A:668:TYR:CE1	1:A:840:VAL:HG13	2.56	0.40
1:A:594:ARG:NH1	1:A:594:ARG:HG2	2.36	0.40
1:A:590:THR:CG2	1:A:592:VAL:HB	2.52	0.40
1:A:608:THR:O	1:A:608:THR:CG2	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	331/360 (92%)	305 (92%)	22 (7%)	4 (1%)	13 13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	330/360 (92%)	316 (96%)	12 (4%)	2 (1%)	25	30
All	All	661/720 (92%)	621 (94%)	34 (5%)	6 (1%)	17	19

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	805	ASP
1	A	587	THR
1	A	594	ARG
1	A	893	CYS
1	B	893	CYS
1	A	588	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/311 (93%)	273 (95%)	15 (5%)	23	32
1	B	287/311 (92%)	276 (96%)	11 (4%)	33	46
All	All	575/622 (92%)	549 (96%)	26 (4%)	27	38

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	589	ILE
1	A	590	THR
1	A	594	ARG
1	A	605	PHE
1	A	649	ASP
1	A	650	GLU
1	A	666	VAL
1	A	709	HIS
1	A	719	SER
1	A	774	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	822	ASP
1	A	865	SER
1	A	866	LYS
1	A	899	TRP
1	A	906	SER
1	B	586	VAL
1	B	602	LEU
1	B	605	PHE
1	B	621	THR
1	B	634	ARG
1	B	636	LEU
1	B	647	CYS
1	B	804	SER
1	B	822	ASP
1	B	831	ASP
1	B	906	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	895	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	GAI	A	1008	-	3,3,3	3.20	1 (33%)	3,3,3	1.58	1 (33%)
5	FMT	A	1004	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	B	1008	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	B	1006	-	0,2,2	0.00	-	0,1,1	0.00	-
5	FMT	B	1005	-	0,2,2	0.00	-	0,1,1	0.00	-
4	D5Z	A	1003	-	43,43,43	0.58	1 (2%)	51,62,62	1.26	5 (9%)
6	GAI	B	1007	-	3,3,3	0.22	0	3,3,3	0.58	0
5	FMT	A	1005	-	0,2,2	0.00	-	0,1,1	0.00	-
6	GAI	B	1003	-	3,3,3	3.00	1 (33%)	3,3,3	1.37	1 (33%)
6	GAI	A	1007	-	3,3,3	1.13	0	3,3,3	0.37	0
4	D5Z	B	1004	-	43,43,43	0.65	1 (2%)	51,62,62	1.18	4 (7%)
5	FMT	A	1006	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	D5Z	B	1004	-	-	3/16/53/53	0/6/6/6
4	D5Z	A	1003	-	-	2/16/53/53	0/6/6/6

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1008	GAI	C-N3	-5.35	1.26	1.36
6	B	1003	GAI	C-N3	-4.87	1.27	1.36
4	B	1004	D5Z	C7-N2	2.29	1.50	1.47
4	A	1003	D5Z	C7-N2	2.10	1.50	1.47

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	D5Z	C17-C10-N3	5.51	131.69	120.97
4	B	1004	D5Z	C17-C10-N3	4.13	129.01	120.97
4	A	1003	D5Z	N6-C11-N4	-2.96	123.27	127.22
4	B	1004	D5Z	C11-N4-C10	2.74	121.50	113.91
4	A	1003	D5Z	C11-N4-C10	2.73	121.48	113.91
4	B	1004	D5Z	N6-C11-N4	-2.70	123.61	127.22
4	A	1003	D5Z	O2-C20-N2	2.36	124.51	121.64
6	A	1008	GAI	N3-C-N2	2.31	121.64	116.13
4	B	1004	D5Z	C17-C10-N4	-2.25	117.87	122.66
6	B	1003	GAI	N3-C-N2	2.04	121.00	116.13
4	A	1003	D5Z	C17-C10-N4	-2.02	118.36	122.66

There are no chirality outliers.



All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1003	D5Z	C17-C10-N3-C9
4	A	1003	D5Z	N4-C10-N3-C9
4	B	1004	D5Z	C17-C10-N3-C9
4	B	1004	D5Z	N4-C10-N3-C9
4	B	1004	D5Z	C28-C2-O1-C1

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	333/360 (92%)	0.21	10 (3%) 50 57	41, 63, 110, 129	0
1	B	332/360 (92%)	0.36	11 (3%) 46 53	30, 61, 106, 130	0
All	All	665/720 (92%)	0.29	21 (3%) 47 55	30, 62, 108, 130	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	586	VAL	4.7
1	B	602	LEU	4.5
1	A	868	MET	4.3
1	B	601	GLU	4.1
1	B	605	PHE	3.9
1	B	599	VAL	3.8
1	A	858	VAL	3.7
1	A	856	VAL	3.7
1	A	852	LYS	3.4
1	A	859	LEU	3.3
1	B	603	PRO	2.8
1	A	857	GLU	2.7
1	B	607	VAL	2.6
1	B	600	CYS	2.4
1	B	589	ILE	2.4
1	A	586	VAL	2.3
1	B	620	SER	2.3
1	A	633	TYR	2.1
1	A	637	LEU	2.1
1	A	602	LEU	2.0
1	B	634	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	FMT	A	1005	3/3	0.77	0.12	79,79,84,91	0
3	MG	A	1002	1/1	0.84	0.09	37,37,37,37	0
5	FMT	B	1006	3/3	0.84	0.44	77,77,89,94	0
6	GAI	B	1007	4/4	0.87	0.15	65,70,73,79	0
6	GAI	B	1003	4/4	0.87	0.15	84,85,88,90	0
5	FMT	A	1004	3/3	0.88	0.21	72,72,78,84	0
5	FMT	A	1006	3/3	0.89	0.20	62,62,68,78	0
5	FMT	B	1008	3/3	0.89	0.20	84,84,88,93	0
5	FMT	B	1005	3/3	0.91	0.21	76,76,80,85	0
6	GAI	A	1007	4/4	0.92	0.17	72,77,80,81	0
6	GAI	A	1008	4/4	0.92	0.17	79,83,88,93	0
4	D5Z	A	1003	38/38	0.93	0.16	54,69,77,81	0
2	ZN	A	1001	1/1	0.95	0.12	56,56,56,56	0
4	D5Z	B	1004	38/38	0.96	0.17	46,54,79,80	0
2	ZN	B	1001	1/1	0.99	0.18	45,45,45,45	0
3	MG	B	1002	1/1	1.00	0.17	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around D5Z A 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.