



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2023 – 06:09 PM EDT

PDB ID : 8FFE  
Title : Crystal structure of LRP6 E1E2 domains bound to YW210.09 Fab and engineered XWnt8 peptide  
Authors : Jude, K.M.; Tsutsumi, N.; Waghray, D.; Garcia, K.C.  
Deposited on : 2022-12-08  
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.32.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	613	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 90%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 20px;">2%      90%      10%</p>
2	H	276	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 79%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 20px;">3%      79%      8%      13%</p>
3	L	225	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 88%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 20px;">2%      88%      8%      .</p>
4	B	4	<div style="display: flex; align-items: center;"> <div style="width: 25%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 75%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 20px;">25%      75%</p>
5	C	5	<div style="display: flex; align-items: center;"> <div style="width: 20%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 80%; height: 10px; background-color: yellow; margin-right: 5px;"></div> </div> <p style="margin-left: 20px;">20%      80%</p>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
6	D	3	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FUC	B	4	-	-	-	X
6	MAN	D	2	-	-	-	X

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 9116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Low-density lipoprotein receptor-related protein 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	613	4889	3096	842	927	24	0	4	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	632	HIS	-	expression tag	UNP O75581


- Molecule 2 is a protein called YW210.09 Fab heavy chain with engineered XWnt8 NC peptide and linker.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	239	1796	1136	303	349	8	0	2	0

- Molecule 3 is a protein called YW210.09 Fab light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	215	1639	1024	270	339	6	0	0	0

- Molecule 4 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	B	4	48	28	2	18	0	0	0

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	C	5	59	34	2	23	0	0	0

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
6	D	3	39	22	2	15	0	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	A	1	6	3	3	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	H	1	Total	C	O	0	0
			6	3	3		
7	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	3	Total	Na	0	0
			3	3		

- Molecule 9 is SUCCINIC ACID (three-letter code: SIN) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	L	1	Total	C	O	0	0
			8	4	4		


- Molecule 10 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
10	A	346	Total 347	O 347	0	1
10	H	130	Total 130	O 130	0	0
10	L	116	Total 116	O 116	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Low-density lipoprotein receptor-related protein 6



- Molecule 2: YW210.09 Fab heavy chain with engineered XWnt8 NC peptide and linker



- Molecule 3: YW210.09 Fab light chain




- Molecule 4: alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose







- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  20% 80%



- Molecule 6: alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  67% 33%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	125.21Å 91.39Å 104.06Å 90.00° 103.67° 90.00°	Depositor
Resolution (Å)	44.37 – 1.72 44.64 – 1.72	Depositor EDS
% Data completeness (in resolution range)	90.6 (44.37-1.72) 83.5 (44.64-1.72)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.35 (at 1.72Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.185 , 0.231 0.184 , 0.229	Depositor DCC
$R_{free}$ test set	5472 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.3	Xtrriage
Anisotropy	0.469	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9116	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, NA, MAN, SIN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/5009	0.67	0/6819
2	H	0.48	0/1847	0.63	0/2520
3	L	0.45	0/1674	0.63	1/2277 (0.0%)
All	All	0.46	0/8530	0.66	1/11616 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	170	ASP	CB-CG-OD1	5.92	123.63	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4889	0	4728	36	0
2	H	1796	0	1733	11	0
3	L	1639	0	1587	9	0
4	B	48	0	43	1	0
5	C	59	0	52	0	0
6	D	39	0	34	0	0
7	A	30	0	40	7	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	H	6	0	8	0	0
7	L	6	0	8	0	0
8	A	3	0	0	0	0
9	L	8	0	4	0	0
10	A	347	0	0	2	0
10	H	130	0	0	0	0
10	L	116	0	0	3	0
All	All	9116	0	8237	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:VAL:HA	7:A:702:GOL:H32	1.72	0.70
1:A:46:VAL:HG12	1:A:47:VAL:HG23	1.73	0.70
1:A:327:ALA:H	7:A:701:GOL:H12	1.60	0.67
1:A:235:ASP:HA	1:A:252:LYS:HD2	1.79	0.65
1:A:621:ILE:HG21	1:A:628:ILE:HD12	1.85	0.58
1:A:611:ARG:NH1	10:A:803:HOH:O	2.37	0.56
3:L:105:GLU:HG2	3:L:106:ILE:N	2.19	0.56
1:A:476:LEU:HB2	1:A:605:TYR:O	2.06	0.55
2:H:1195:ILE:HD13	2:H:1210:LYS:HA	1.88	0.55
1:A:420:THR:HG22	1:A:427:ILE:HG12	1.90	0.54
1:A:513:LYS:HZ3	7:A:708:GOL:H11	1.73	0.53
1:A:100:LEU:HD11	1:A:109:LEU:HD11	1.90	0.53
1:A:78:THR:HG22	1:A:86:VAL:HA	1.90	0.53
3:L:142:ARG:NH1	10:L:402:HOH:O	2.42	0.53
1:A:187:LEU:HD11	1:A:196:LEU:HD11	1.92	0.52
2:H:1126:PRO:HG3	2:H:1138:LEU:HB3	1.91	0.51
1:A:513:LYS:NZ	7:A:708:GOL:H11	2.26	0.51
1:A:427:ILE:HG21	1:A:460:MET:HE1	1.91	0.51
3:L:186:TYR:CE2	3:L:209:ARG:HD3	2.45	0.51
1:A:591:PRO:HB2	1:A:610:LEU:HD23	1.94	0.50
2:H:1047:TRP:CZ2	2:H:1049:GLY:HA2	2.46	0.50
1:A:330:LEU:HD21	7:A:701:GOL:H2	1.94	0.50
2:H:1082:MET:HE1	2:H:1090:TYR:CZ	2.48	0.49
1:A:134:TRP:CD1	1:A:135:GLN:HG3	2.47	0.49
1:A:252:LYS:HD3	1:A:253:TYR:CZ	2.47	0.49
1:A:374:ILE:O	1:A:386:ARG:HA	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:40:PRO:HG2	3:L:165:GLU:HG3	1.95	0.47
1:A:197:TYR:OH	1:A:252:LYS:HG3	2.13	0.47
1:A:397:VAL:HG12	1:A:398:VAL:HG23	1.97	0.46
1:A:517:MET:HE1	10:A:1062:HOH:O	2.14	0.45
1:A:28:ARG:HD3	1:A:52:ASP:OD1	2.17	0.45
1:A:436:MET:HG3	1:A:601:HIS:CD2	2.52	0.44
1:A:330:LEU:HD23	1:A:343:SER:HA	2.00	0.44
2:H:1119:PRO:HB3	2:H:1145:TYR:HB3	2.00	0.44
1:A:436:MET:HG3	1:A:601:HIS:HD2	1.83	0.43
2:H:1030:THR:O	2:H:1053:TYR:HB3	2.18	0.43
1:A:343:SER:HB2	7:A:701:GOL:H32	2.00	0.43
3:L:187:GLU:O	3:L:209:ARG:NH2	2.52	0.43
1:A:21:PRO:HB3	1:A:233:PHE:CD2	2.54	0.42
1:A:361:HIS:CE1	2:H:236:ARG:HD3	2.53	0.42
1:A:579:LYS:HE3	7:A:702:GOL:H2	2.01	0.42
2:H:1012:VAL:HG11	2:H:1082(C):LEU:HD13	2.01	0.42
1:A:331:LEU:HD11	1:A:578:LEU:HG	2.00	0.42
3:L:24:ARG:NE	3:L:70:ASP:OD1	2.53	0.42
1:A:139:GLN:HB3	1:A:157:TRP:CE2	2.55	0.41
3:L:105:GLU:HG3	3:L:173:TYR:OH	2.20	0.41
2:H:1143:LYS:HE2	10:L:405:HOH:O	2.20	0.41
3:L:107:LYS:NZ	10:L:403:HOH:O	2.44	0.41
2:H:1094:ARG:HG2	2:H:1095:GLY:O	2.21	0.41
1:A:487:THR:HG22	4:B:1:NAG:H82	2.02	0.41
1:A:572:LEU:HD12	1:A:573:PRO:HD2	2.03	0.41
2:H:1100(G):TYR:OH	3:L:94:THR:OG1	2.26	0.41
1:A:95:LEU:HB2	1:A:115:GLU:HB3	2.02	0.40
1:A:182:TYR:HB3	1:A:202:LYS:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	615/613 (100%)	570 (93%)	42 (7%)	3 (0%)	29	13
2	H	235/276 (85%)	231 (98%)	4 (2%)	0	100	100
3	L	213/225 (95%)	209 (98%)	4 (2%)	0	100	100
All	All	1063/1114 (95%)	1010 (95%)	50 (5%)	3 (0%)	47	24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	490	GLY
1	A	336[A]	ARG
1	A	336[B]	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	528/529 (100%)	518 (98%)	10 (2%)	57	39
2	H	197/218 (90%)	194 (98%)	3 (2%)	65	49
3	L	188/196 (96%)	185 (98%)	3 (2%)	62	47
All	All	913/943 (97%)	897 (98%)	16 (2%)	59	41

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	ASP
1	A	77	ARG
1	A	162	LYS
1	A	165	ARG
1	A	179	SER
1	A	243	SER
1	A	252	LYS
1	A	436	MET
1	A	565	ARG
1	A	608	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	H	1058	ASN
2	H	1129	LYS
2	H	1206	LYS
3	L	50	SER
3	L	123	GLU
3	L	202	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	B	1	1,4	14,14,15	0.56	0	17,19,21	0.59	0
4	FUC	B	2	4	10,10,11	0.65	0	14,14,16	0.90	1 (7%)
4	NAG	B	3	4	14,14,15	0.24	0	17,19,21	0.39	0
4	FUC	B	4	4	10,10,11	1.15	0	14,14,16	1.10	2 (14%)
5	NAG	C	1	1,5	14,14,15	0.64	1 (7%)	17,19,21	0.46	0
5	NAG	C	2	5	14,14,15	0.35	0	17,19,21	0.47	0
5	BMA	C	3	5	11,11,12	0.86	0	15,15,17	0.91	1 (6%)
5	FUC	C	4	5	10,10,11	0.88	1 (10%)	14,14,16	0.79	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	FUC	C	5	5	10,10,11	0.95	0	14,14,16	0.79	1 (7%)
6	NAG	D	1	1,6	14,14,15	0.36	0	17,19,21	0.58	0
6	MAN	D	2	6	11,11,12	0.94	0	15,15,17	1.01	1 (6%)
6	NAG	D	3	6	14,14,15	0.56	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	1	1,4	-	0/6/23/26	0/1/1/1
4	FUC	B	2	4	-	-	0/1/1/1
4	NAG	B	3	4	-	2/6/23/26	0/1/1/1
4	FUC	B	4	4	-	-	0/1/1/1
5	NAG	C	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	C	2	5	-	0/6/23/26	0/1/1/1
5	BMA	C	3	5	-	2/2/19/22	0/1/1/1
5	FUC	C	4	5	-	-	0/1/1/1
5	FUC	C	5	5	-	-	0/1/1/1
6	NAG	D	1	1,6	-	0/6/23/26	0/1/1/1
6	MAN	D	2	6	-	0/2/19/22	0/1/1/1
6	NAG	D	3	6	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1	NAG	O5-C1	-2.21	1.40	1.43
5	C	4	FUC	C2-C3	2.03	1.55	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	2	MAN	C1-O5-C5	2.49	115.56	112.19
5	C	3	BMA	C1-O5-C5	2.37	115.41	112.19
5	C	5	FUC	O2-C2-C1	2.28	113.81	109.15
4	B	4	FUC	C1-O5-C5	2.23	117.84	112.78
4	B	2	FUC	C1-O5-C5	2.22	117.80	112.78
4	B	4	FUC	C1-C2-C3	2.01	112.14	109.67

There are no chirality outliers.



All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	3	NAG	C8-C7-N2-C2
4	B	3	NAG	O7-C7-N2-C2
5	C	3	BMA	C4-C5-C6-O6
5	C	3	BMA	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
9	SIN	L	302	-	7,7,7	0.99	0	8,8,8	1.45	1 (12%)
7	GOL	A	704	-	5,5,5	1.08	0	5,5,5	0.78	0
7	GOL	L	301	-	5,5,5	0.82	0	5,5,5	1.03	0
7	GOL	H	1301	-	5,5,5	1.33	0	5,5,5	0.72	0
7	GOL	A	702	-	5,5,5	1.86	2 (40%)	5,5,5	1.19	0
7	GOL	A	703	-	5,5,5	1.03	0	5,5,5	0.89	0
7	GOL	A	701	-	5,5,5	1.56	2 (40%)	5,5,5	0.79	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GOL	A	708	-	5,5,5	1.54	2 (40%)	5,5,5	1.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	SIN	L	302	-	-	4/5/5/5	-
7	GOL	A	704	-	-	1/4/4/4	-
7	GOL	L	301	-	-	2/4/4/4	-
7	GOL	H	1301	-	-	2/4/4/4	-
7	GOL	A	702	-	-	1/4/4/4	-
7	GOL	A	703	-	-	1/4/4/4	-
7	GOL	A	701	-	-	1/4/4/4	-
7	GOL	A	708	-	-	1/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	702	GOL	C1-C2	2.82	1.63	1.51
7	A	702	GOL	C3-C2	2.80	1.63	1.51
7	A	708	GOL	C3-C2	2.50	1.62	1.51
7	A	701	GOL	C1-C2	2.37	1.61	1.51
7	A	708	GOL	C1-C2	2.26	1.61	1.51
7	A	701	GOL	C3-C2	2.12	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	L	302	SIN	O4-C4-C3	2.38	121.67	114.03

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	708	GOL	O1-C1-C2-C3
7	A	703	GOL	C1-C2-C3-O3
7	A	704	GOL	O1-C1-C2-C3
7	L	301	GOL	O1-C1-C2-C3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
7	A	702	GOL	O1-C1-C2-O2
7	H	1301	GOL	O2-C2-C3-O3
7	A	701	GOL	O2-C2-C3-O3
9	L	302	SIN	C2-C3-C4-O3
9	L	302	SIN	C2-C3-C4-O4
7	H	1301	GOL	C1-C2-C3-O3
7	L	301	GOL	O1-C1-C2-O2
9	L	302	SIN	O2-C1-C2-C3
9	L	302	SIN	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	702	GOL	2	0
7	A	701	GOL	3	0
7	A	708	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	613/613 (100%)	-0.03	14 (2%) 60 65	23, 34, 61, 112	0
2	H	239/276 (86%)	-0.05	9 (3%) 40 45	24, 34, 66, 102	0
3	L	215/225 (95%)	-0.05	5 (2%) 60 65	27, 36, 59, 100	0
All	All	1067/1114 (95%)	-0.04	28 (2%) 56 60	23, 35, 63, 112	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	83	THR	8.0
1	A	80	PHE	7.2
1	A	81	ASN	6.6
1	A	20	ALA	5.3
3	L	210	GLY	4.9
1	A	84	GLU	4.1
2	H	239	ALA	3.5
1	A	46	VAL	3.4
3	L	212	CYS	3.4
3	L	126	LYS	3.4
2	H	237	ALA	3.2
2	H	1129	LYS	3.2
1	A	22	LEU	3.1
1	A	174	PHE	2.7
2	H	1056[A]	TYR	2.7
2	H	236	ARG	2.6
2	H	1000	SER	2.6
3	L	169	LYS	2.4
2	H	1134	GLY	2.4
1	A	82	LYS	2.3
1	A	218	ALA	2.3
1	A	628	ILE	2.2
3	L	83	PHE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	H	238	ILE	2.1
1	A	39	GLY	2.1
1	A	85	SER	2.1
1	A	632	HIS	2.0
2	H	1128	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	FUC	B	4	10/11	0.28	0.42	65,86,91,99	0
6	NAG	D	1	14/15	0.68	0.16	73,78,88,88	0
5	BMA	C	3	11/12	0.69	0.25	59,71,77,78	0
6	MAN	D	2	11/12	0.73	0.41	95,97,103,105	0
6	NAG	D	3	14/15	0.77	0.19	69,86,89,89	0
4	NAG	B	1	14/15	0.78	0.12	61,76,87,88	0
4	NAG	B	3	14/15	0.84	0.16	85,93,96,99	0
5	FUC	C	4	10/11	0.88	0.35	57,61,68,72	0
4	FUC	B	2	10/11	0.89	0.25	82,92,96,97	0
5	FUC	C	5	10/11	0.92	0.16	43,47,49,50	0
5	NAG	C	2	14/15	0.92	0.20	41,52,61,63	0
5	NAG	C	1	14/15	0.93	0.13	35,43,49,50	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain B:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	GOL	H	1301	6/6	0.59	0.29	51,55,60,62	0
7	GOL	A	708	6/6	0.65	0.33	35,46,50,53	0
7	GOL	A	701	6/6	0.69	0.23	38,44,46,51	0
9	SIN	L	302	8/8	0.78	0.16	46,54,62,64	0
7	GOL	A	704	6/6	0.83	0.17	41,47,48,51	0
7	GOL	A	703	6/6	0.85	0.22	44,45,51,52	0
7	GOL	A	702	6/6	0.87	0.26	31,36,42,47	0
8	NA	A	706	1/1	0.92	0.10	27,27,27,27	0
7	GOL	L	301	6/6	0.92	0.22	47,48,50,53	0
8	NA	A	705	1/1	0.96	0.04	32,32,32,32	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
8	NA	A	707	1/1	0.98	0.09	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.