



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 6, 2023 – 01:56 AM EST

PDB ID : 5FGU  
Title : Structure of Sda1 nuclease apoprotein as an EGFP fixed-arm fusion  
Authors : Moon, A.F.; Krahn, J.M.; Xun, L.; Cuneo, M.J.; Pedersen, L.C.  
Deposited on : 2015-12-21  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	565	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4283 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein, Extracellular streptodornase D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	495	3892	2475	663	743	11	0	15	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P42212
A	-1	SER	-	expression tag	UNP P42212
A	0	HIS	-	expression tag	UNP P42212
A	?	-	SER	chromophore	UNP P42212
A	?	-	TYR	chromophore	UNP P42212
A	66	CR2	GLY	chromophore	UNP P42212
A	72	ALA	SER	engineered mutation	UNP P42212
A	230	ALA	-	linker	UNP P42212
A	231	ALA	-	linker	UNP P42212
A	232	ALA	-	linker	UNP P42212
A	1188	GLY	HIS	engineered mutation	UNP Q675N6

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C O	0	0
			4	2 2		
3	A	1	Total	C O	0	0
			4	2 2		
3	A	1	Total	C O	0	0
			4	2 2		
3	A	1	Total	C O	0	0
			4	2 2		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	327	Total O 329 329	0	4

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein, Extracellular streptodornase D



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.41Å 139.37Å 120.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.28 – 1.90 38.28 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (38.28-1.90) 95.8 (38.28-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.177 , 0.201 0.177 , 0.200	Depositor DCC
$R_{free}$ test set	2734 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.3	Xtrriage
Anisotropy	0.896	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.015 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.027 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4283	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, EDO, SO4, CR2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/3979	0.64	4/5393 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73[A]	ARG	NE-CZ-NH2	-8.83	115.89	120.30
1	A	73[B]	ARG	NE-CZ-NH2	-8.83	115.89	120.30
1	A	73[A]	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	A	73[B]	ARG	NE-CZ-NH1	5.83	123.22	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3892	0	3701	11	0
2	A	30	0	0	0	0
3	A	16	0	24	0	0
4	A	16	0	12	0	0
5	A	329	0	0	5	0
All	All	4283	0	3737	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73[B]:ARG:NH1	5:A:1502:HOH:O	2.22	0.72
1:A:1077:ARG:NH1	1:A:1083:GLY:O	2.28	0.65
1:A:77:HIS:NE2	1:A:1107:GLU:OE2	2.35	0.51
1:A:79:LYS:NZ	5:A:1508[A]:HOH:O	2.43	0.50
1:A:73[A]:ARG:NH2	5:A:1511:HOH:O	2.45	0.48
1:A:1351:VAL:HG22	1:A:1382:HIS:CE1	2.50	0.47
1:A:1140:PHE:CD1	1:A:1147:ARG:HA	2.50	0.47
1:A:36:ASP:OD1	5:A:1501:HOH:O	2.21	0.45
1:A:1063:LEU:CD2	1:A:1243[A]:LYS:HE3	2.50	0.41
1:A:161:ILE:HG13	1:A:185:ASN:HB2	2.03	0.40
1:A:198:ASN:OD1	5:A:1503:HOH:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	498/565 (88%)	489 (98%)	9 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	405/492 (82%)	403 (100%)	2 (0%)	88 89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	LEU
1	A	163	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CR2	A	66	1	20,20,21	3.27	9 (45%)	25,27,29	2.45	13 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	A	66	1	-	2/6/25/26	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CR2	CA2-C2	9.95	1.58	1.48
1	A	66	CR2	CA1-C1	5.39	1.55	1.49
1	A	66	CR2	C1-N3	4.79	1.44	1.37
1	A	66	CR2	CG2-CB2	4.56	1.55	1.46
1	A	66	CR2	CA3-N3	-3.31	1.40	1.47
1	A	66	CR2	CB2-CA2	-3.00	1.32	1.35
1	A	66	CR2	C2-N3	2.40	1.45	1.39
1	A	66	CR2	CE2-CD2	2.39	1.43	1.38
1	A	66	CR2	O2-C2	-2.29	1.18	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CR2	O2-C2-CA2	-4.75	128.29	130.96
1	A	66	CR2	CB2-CA2-N2	4.52	135.09	128.83
1	A	66	CR2	CG2-CB2-CA2	-3.92	125.14	129.94
1	A	66	CR2	CB2-CA2-C2	-3.85	117.68	122.28
1	A	66	CR2	C1-CA1-N1	-3.17	105.83	112.85
1	A	66	CR2	CA1-C1-N2	3.12	128.45	124.28
1	A	66	CR2	CD2-CG2-CD1	2.84	121.84	117.64
1	A	66	CR2	CE2-CD2-CG2	-2.80	117.60	121.25
1	A	66	CR2	O3-C3-CA3	-2.72	118.17	126.39
1	A	66	CR2	C2-N3-C1	-2.71	106.66	107.99
1	A	66	CR2	CA1-C1-N3	-2.69	118.91	122.52
1	A	66	CR2	C2-CA2-N2	-2.52	107.17	108.93
1	A	66	CR2	O2-C2-N3	2.31	128.94	124.35

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	CR2	C2-CA2-CB2-CG2
1	A	66	CR2	N2-CA2-CB2-CG2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	1401	-	4,4,4	0.16	0	6,6,6	0.14	0
3	EDO	A	1410	-	3,3,3	0.39	0	2,2,2	0.37	0
4	ACT	A	1411	-	3,3,3	0.69	0	3,3,3	1.66	1 (33%)
4	ACT	A	1414	-	3,3,3	0.74	0	3,3,3	1.47	0
3	EDO	A	1407	-	3,3,3	0.43	0	2,2,2	0.40	0
2	SO4	A	1404	-	4,4,4	0.18	0	6,6,6	0.26	0
2	SO4	A	1405	-	4,4,4	0.15	0	6,6,6	0.09	0
2	SO4	A	1402	-	4,4,4	0.14	0	6,6,6	0.19	0
4	ACT	A	1413	-	3,3,3	0.74	0	3,3,3	1.34	0
4	ACT	A	1412	-	3,3,3	0.87	0	3,3,3	1.42	0
3	EDO	A	1409	-	3,3,3	0.48	0	2,2,2	0.20	0
2	SO4	A	1403	-	4,4,4	0.11	0	6,6,6	0.28	0
3	EDO	A	1408	-	3,3,3	0.49	0	2,2,2	0.26	0
2	SO4	A	1406	-	4,4,4	0.32	0	6,6,6	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1408	-	-	0/1/1/1	-
3	EDO	A	1410	-	-	0/1/1/1	-
3	EDO	A	1409	-	-	0/1/1/1	-
3	EDO	A	1407	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1411	ACT	OXT-C-CH3	2.15	124.04	115.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	494/565 (87%)	0.35	57 (11%) <b>4</b> <b>5</b>	18, 29, 74, 127	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1143	ASN	6.4
1	A	1355	ASN	5.9
1	A	1357	ASP	5.8
1	A	1349	SER	5.7
1	A	1385	THR	5.5
1	A	1354	TYR	5.2
1	A	1383	HIS	4.4
1	A	1342	TYR	4.2
1	A	1144	THR	4.2
1	A	1338	GLN	4.2
1	A	1083	GLY	4.1
1	A	192	PRO	4.0
1	A	1336	SER	4.0
1	A	1384	SER	4.0
1	A	1085	SER	3.8
1	A	1337	LEU	3.4
1	A	1353	TRP	3.4
1	A	1343	VAL	3.2
1	A	1087	LYS	3.2
1	A	1142	LYS	3.1
1	A	1084	GLN	2.9
1	A	1359	MET	2.9
1	A	1088[A]	PRO	2.9
1	A	1141	PHE	2.9
1	A	1177	ASN	2.8
1	A	1341	VAL	2.8
1	A	1082	SER	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1077	ARG	2.8
1	A	1358	SER	2.7
1	A	1377	LEU	2.7
1	A	1145	ASN	2.7
1	A	1121	ALA	2.6
1	A	1339	ASP	2.6
1	A	1173	LEU	2.6
1	A	1371	MET	2.6
1	A	1172	LEU	2.6
1	A	191	GLY	2.6
1	A	1382	HIS	2.5
1	A	1378	THR	2.5
1	A	1174	SER	2.5
1	A	157	GLN	2.4
1	A	1216	ASP	2.3
1	A	1152	ASP	2.3
1	A	1159	ASN	2.3
1	A	1175	ASN	2.3
1	A	1350	ASP	2.3
1	A	1215	ASN	2.3
1	A	1075	THR	2.3
1	A	194	LEU	2.2
1	A	1086	THR	2.2
1	A	1176	GLY	2.2
1	A	1120	SER	2.1
1	A	1074	THR	2.1
1	A	1340	THR	2.1
1	A	1346	ASN	2.1
1	A	1381	LYS	2.0
1	A	1158	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CR2	A	66	19/20	0.96	0.10	20,24,30,34	0



### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	1408	4/4	0.74	0.39	50,56,61,61	0
2	SO4	A	1405	5/5	0.79	0.24	89,92,97,99	5
4	ACT	A	1413	4/4	0.80	0.18	51,57,57,61	0
3	EDO	A	1410	4/4	0.82	0.25	46,52,57,58	0
3	EDO	A	1409	4/4	0.86	0.15	49,51,57,58	0
4	ACT	A	1414	4/4	0.87	0.25	58,64,67,70	0
3	EDO	A	1407	4/4	0.89	0.16	47,49,49,58	0
4	ACT	A	1412	4/4	0.93	0.10	45,47,55,60	0
4	ACT	A	1411	4/4	0.94	0.13	37,42,44,63	0
2	SO4	A	1404	5/5	0.96	0.13	36,37,59,68	0
2	SO4	A	1401	5/5	0.96	0.27	67,70,74,81	0
2	SO4	A	1403	5/5	0.97	0.14	36,47,52,63	0
2	SO4	A	1402	5/5	0.98	0.17	43,47,65,71	0
2	SO4	A	1406	5/5	0.98	0.06	26,29,37,39	5

### 6.5 Other polymers [i](#)

There are no such residues in this entry.