

# wwPDB X-ray Structure Validation Summary Report (i)

#### Aug 10, 2020 – 02:08 PM BST

PDB ID : 1FJR

Title : CRYSTAL STRUCTURE OF THE ECTODOMAIN OF METHUSELAH

Authors: West Jr., A.P.; Llamas, L.L.; Snow, P.M.; Benzer, S.; Bjorkman, P.J.

Deposited on : 2000-08-08

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain					
1	A	195	69%	24%	• • • •			
1	В	195	69%	24%				
2	С	2	50%	50%				



# 2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 3369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called METHUSELAH ECTODOMAIN.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	188	Total	С	N	Ο	S	0	0	0
1	A	100	1528	965	262	287	14	0		
1	B	188	Total	С	N	О	S	0	0	0
1	D	100	1528	965	262	287	14		U	U


• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	С	2	Total 28	C 16	N 2	O 10	0	0	0


• Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	A	1	Total C N O	0	Ω	
	Λ	1	14 8 1 5	U	U	
3	A	1	Total C N O	0	0	
	Λ	1	14 8 1 5	U	U	
3	В	1	Total C N O	0	0	
)	Б	1	14 8 1 5	0	U	
3	В	1	Total C N O	0	0	
)	Б	1	14 8 1 5	0	0	
2	В	1	Total C N O	0	0	
	D	1	14 8 1 5		U	

 $\bullet$  Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 





$\mathbf{Mol}$	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	В	1	Total O S 5 4 1	0	0

 $\bullet$  Molecule 5 is LEAD (II) ION (three-letter code: PB) (formula: Pb).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	2	Total Pb 2 2	0	0
5	A	2	Total Pb 2 2	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	100	Total O 100 100	0	0
6	В	101	Total O 101 101	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: METHUSELAH ECTODOMAIN



 $\bullet \ \, \text{Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose} \\$ 





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 21 21 21	Depositor	
Cell constants	68.94Å 73.00Å 115.11Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	20.00 - 2.30	Depositor	
% Data completeness	99.5 (20.00-2.30)	Depositor	
(in resolution range)	33.9 (20.00 2.30)	Depositor	
$R_{merge}$	0.07	Depositor	
$R_{sym}$	(Not available)	Depositor	
Refinement program	CNS 1.0	Depositor	
$R, R_{free}$	0.212 , $0.227$	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3369	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP	



## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PB, NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.44	0/1563	0.68	1/2115 (0.0%)	
1	В	0.41	0/1563	0.67	1/2115 (0.0%)	
All	All	0.42	0/3126	0.68	2/4230 (0.0%)	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	94	LEU	CA-CB-CG	5.96	129.00	115.30
1	В	94	LEU	CA-CB-CG	5.89	128.84	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1528	0	1456	57	0
1	В	1528	0	1456	54	0
2	С	28	0	25	1	0
3	A	28	0	26	0	0
3	В	42	0	39	2	0
4	A	5	0	0	0	0
4	В	5	0	0	0	0

Continued on next page...



Continued	trom	nromanne	naae
-	110116	picolous	puyc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	2	0	0	0	0
5	В	2	0	0	0	0
6	A	100	0	0	1	0
6	В	101	0	0	2	0
All	All	3369	0	3002	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance}  ({\rm \AA}) \end{array}$	$egin{array}{c} { m Clash} \ { m overlap} \ ({ m \AA}) \end{array}$
1:B:73:HIS:HB2	1:B:112:PHE:HE1	1.39	0.87
1:B:51:LYS:HD2	1:B:51:LYS:H	1.41	0.85
1:A:151:ARG:HH11	1:A:184:ASN:HD22	1.26	0.83
1:B:67:VAL:HG11	1:B:116:LEU:HD13	1.61	0.83
1:B:71:CYS:H	1:B:166:GLN:NE2	1.77	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	$\mathbf{Outliers}$	Perce	$\mathbf{ntiles}$
1	A	186/195~(95%)	177 (95%)	8 (4%)	1 (0%)	29	35
1	В	186/195~(95%)	177 (95%)	9 (5%)	0	100	100
All	All	372/390~(95%)	354 (95%)	17 (5%)	1 (0%)	41	50

#### All (1) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	54	ARG

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric C		Outliers	Percentiles
1	A	171/177 (97%)	161 (94%)	10 (6%)	20 27
1	В	171/177 (97%)	162 (95%)	9 (5%)	22 31
All	All	342/354 (97%)	323 (94%)	19 (6%)	21 29

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	GLN
1	В	20	GLN
1	В	68	ARG
1	A	94	LEU
1	В	94	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	${f Res}$	$\mathbf{Type}$
1	В	20	GLN
1	В	50	GLN
1	В	166	GLN
1	A	183	HIS
1	A	184	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Chain	Dog	Link	Bo	nd leng	ths	В	ond ang	cles
MIGI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	С	1	1,2	14,14,15	0.53	0	17,19,21	0.77	1 (5%)
2	NAG	С	2	2	14,14,15	0.52	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	С	2	2	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

$\mathbf{Mol}$	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
2	С	1	NAG	C2-N2-C7	-2.33	119.59	122.90

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	2	NAG	O5-C5-C6-O6
2	С	1	NAG	C4-C5-C6-O6
2	С	1	NAG	O5-C5-C6-O6




There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	1	NAG	1	0
2	С	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



### 5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain		Res	Link	Вс	Bond lengths			Bond angles		
MIOI	Type	Chain	res	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
3	NAG	A	310	1	14,14,15	0.79	1 (7%)	17,19,21	0.71	0	
3	NAG	В	320	1	14,14,15	0.58	0	17,19,21	0.75	1 (5%)	
4	SO4	A	402	5	4,4,4	0.28	0	6,6,6	0.16	0	
3	NAG	В	310	1	14,14,15	0.70	0	17,19,21	0.69	1 (5%)	
3	NAG	В	300	1	14,14,15	0.69	0	17,19,21	0.75	0	
4	SO4	В	401	-	4,4,4	0.30	0	6,6,6	0.15	0	
3	NAG	A	300	1	14,14,15	0.58	0	17,19,21	0.72	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	${f Torsions}$	Rings
3	NAG	В	320	1	-	2/6/23/26	0/1/1/1
3	NAG	В	310	1	=	0/6/23/26	0/1/1/1
3	NAG	A	310	1	-	0/6/23/26	0/1/1/1
3	NAG	В	300	1	-	2/6/23/26	0/1/1/1
3	NAG	A	300	1	-	1/6/23/26	0/1/1/1

#### All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	Ideal(A)
3	A	310	NAG	C1-C2	2.04	1.55	1.52

#### All (2) bond angle outliers are listed below:

N.	Iol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
	3	В	310	NAG	C2-N2-C7	-2.07	119.96	122.90
	3	В	320	NAG	C2-N2-C7	-2.04	120.00	122.90

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	В	320	NAG	O5-C5-C6-O6
3	В	320	NAG	C4-C5-C6-O6
3	В	300	NAG	C1-C2-N2-C7
3	A	300	NAG	O5-C5-C6-O6
3	В	300	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	320	NAG	1	0
3	В	310	NAG	1	0

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

