

# wwPDB EM Validation Summary Report (i)

Dec 10, 2022 – 09:36 am GMT

PDR ID		5FMF
I DD ID	•	
EMDB ID	:	EMD-3114
Title	:	the P-lobe of RNA polymerase II pre-initiation complex
Authors	:	Murakami, K.; Tsai, K.; Kalisman, N.; Bushnell, D.A.; Asturias, F.J.; Korn-
		berg, R.D.
Deposited on	:	2015-11-03
Resolution	:	6.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	0.0.1. dev 43
:	4.02b-467
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	1.9.9
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.31.3
	: : : : :

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 6.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f EM} {f structures} \ (\#{f Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of ch	nain		
-	-	400	66%			
	1	492	35%	52%		8% • •
		1 - 1	60%	_		_
2	2	174	64%		28%	6% •
			37%			
3	A	1733	53%	24%	••	18%
			52%			
4	В	1224	61%		29%	• 6%
			34%			
5	С	266	73%		22%	5%
			57%			
6	D	178	67%		26%	6% •
			43%			
7	E	214	74%		24	•% •
			27%			
8	F	84	60%		37%	•



Mol	Chain	Length	Quality of c	hain	
0	C	171	54%		
9	G	1/1	67%	27%	5% •
10	Н	146	57%	29% 5%	9%
	-	110	52%		
	1	119	76%	20%	•
12	J	65	57%	32%	11%
			46%		
13	K	115	71%	28%	•
14	L	46	35% 41%	22%	•
			63%		
15	М	116	69%	29%	•
16	N	72	56%		
	11		61%	_	•
17	0	122	53%	25% · 17	7%
18	Р	345	74%	130/	15%
10	1	010	53%	1570 •	1378
19	Q	180	59%	36%	••
20	В	160	38%	110/ 269/	
20	10	100	44%	1170 • 2076	
21	S	123	14% 49%	28%	7% •
22	Т	72	38%		_
	1	12	61%		•
23	U	150	43%	51%	••
24	V	174	59%		60/
24	v	114	92%	55%	0% •
25	W	62	90%		10%
26	x	63	83%		10%
	11	00	27%		10%
27	Y	778	33% 38%	• 26%	



# 2 Entry composition (i)

There are 29 unique types of molecules in this entry. The entry contains 55363 atoms, of which 691 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DNA REPAIR HELICASE RAD25, SSL2.

Mol	Chain	Residues		At	AltConf	Trace			
1	1	473	Total 3797	C 2390	N 664	0 720	S 23	0	0

• Molecule 2 is a protein called TRANSCRIPTION ELONGATION FACTOR S-II, DST1.

Mol	Chain	Residues		Ate	$\mathbf{oms}$	AltConf	Trace		
2	2	174	Total 666	С 454	N 99	O 108	${ m S}{ m 5}$	0	104

• Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB1.

Mol	Chain	Residues		A	AltConf	Trace			
3	А	1422	Total 11174	C 7036	N 1954	O 2122	S 62	0	0

• Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB2.

Mol	Chain	Residues		А	AltConf	Trace			
4	В	1156	Total 9140	C 5781	N 1606	O 1697	S 56	0	0

• Molecule 5 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB3.

Mol	Chain	Residues		At	AltConf	Trace			
5	С	266	Total 2095	C 1317	N 348	0 417	S 13	0	0

• Molecule 6 is a protein called RNA POLYMERASE II PRE-INITIATION COMPLEX, RPB4.

Mol	Chain	Residues		At	$\mathbf{oms}$	AltConf	Trace		
6	D	178	Total 1434	C 887	N 257	O 288	${ m S} { m 2}$	0	0



• Molecule 7 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUB-UNIT RPABC1, RPB5.

Mol	Chain	Residues		At	AltConf	Trace			
7	Е	214	Total 1752	C 1111	N 309	O 321	S 11	0	0

• Molecule 8 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUB-UNIT RPABC2, RPB6.

Mol	Chain	Residues		At	oms		AltConf	Trace	
8	F	84	Total 679	C 434	N 115	0 127	${ m S} { m 3}$	0	0

• Molecule 9 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB7.

Mol	Chain	Residues		At	oms			AltConf	Trace
9	G	171	Total 1340	C 861	N 222	0 249	S 8	0	0

• Molecule 10 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC3, RPB8.

Mol	Chain	Residues		At	oms		AltConf	Trace	
10	Н	133	Total 1068	C 673	N 180	0 211	S 4	0	0

• Molecule 11 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB9.

Mol	Chain	Residues		$\mathbf{A}^{\dagger}$	toms		AltConf	Trace	
11	Ι	119	Total 971	C 596	N 179	0 186	S 10	0	0

• Molecule 12 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC5, RPB10.

Mol	Chain	Residues		Atc	$\mathbf{ms}$		AltConf	Trace	
19	т	65	Total	С	Ν	Ο	$\mathbf{S}$	0	0
12	J	05	532	339	93	94	6	0	0

• Molecule 13 is a protein called DNA-DIRECTED RNA POLYMERASE II SUBUNIT RPB11.



Mol	Chain	Residues		At	oms	AltConf	Trace		
13	K	115	Total 920	C 590	N 157	0 171	${ m S} { m 2}$	0	1

• Molecule 14 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC4, RPB12.

Mol	Chain	Residues		Atc	$\mathbf{ms}$	AltConf	Trace		
14	L	46	Total 363	C 224	N 72	O 63	$\frac{S}{4}$	0	0

• Molecule 15 is a protein called RNA POLYMERASE II PRE-INITIATION COMPLEX, TOA1.

Mol	Chain	Residues		At	oms		AltConf	Trace	
15	М	116	Total 956	C 599	N 159	0 195	${ m S} { m 3}$	0	0

• Molecule 16 is a DNA chain called NON-TEMPLATE STRAND DNA.

Mol	Chain	Residues		$\mathbf{A}^{\dagger}$	toms	AltConf	Trace		
16	Ν	72	Total 1500	С 711	N 300	0 417	Р 72	0	0

• Molecule 17 is a protein called TRANSCRIPTION INITIATION FACTOR IIA SUBUNIT 2, TOA2.

Mol	Chain	Residues		At	oms		AltConf	Trace	
17	О	101	Total 792	C 500	N 132	0 156	${S \atop 4}$	0	0

• Molecule 18 is a protein called TRANSCRIPTION INITIATION FACTOR IIB, SUA7.

Mol	Chain	Residues		At	oms			AltConf	Trace
18	Р	293	Total 2269	C 1437	N 386	0 428	S 18	0	0

• Molecule 19 is a protein called TATA-BOX-BINDING PROTEIN, TBP.

Mol	Chain	Residues		At	oms		AltConf	Trace	
19	Q	180	Total 1416	C 921	N 242	0 247	S 6	0	0

• Molecule 20 is a protein called TRANSCRIPTION INITIATION FACTOR IIE SUBUNIT



ALPHA, TFA1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
20	R	118	Total 960	C 625	N 158	0 172	${ m S}{ m 5}$	0	0

• Molecule 21 is a protein called TRANSCRIPTION INITIATION FACTOR IIE SUBUNIT BETA, TFA2.

Mol	Chain	Residues	Atoms				AltConf	Trace	
21	S	121	Total 1005	C 650	N 164	0 188	${ m S} { m 3}$	0	0

• Molecule 22 is a DNA chain called TEMPLATE STRAND DNA.

Mol	Chain	Residues	Atoms				AltConf	Trace	
22	Т	72	Total 1452	C 700	N 233	0 447	Р 72	0	0

• Molecule 23 is a protein called RNA POLYMERASE II PRE-INITIATION COMPLEX, TFG1.

Mol	Chain	Residues	Atoms				AltConf	Trace	
23	U	150	Total 1248	C 793	N 220	0 232	${ m S} { m 3}$	0	0

• Molecule 24 is a protein called TRANSCRIPTION INITIATION FACTOR IIF SUBUNIT BETA, TFG2.

Mol	Chain	Residues	Atoms				AltConf	Trace	
24	V	174	Total 1455	C 930	N 250	O 269	S 6	0	0

• Molecule 25 is a protein called RNA POLYMERASE II TRANSCRIPTION FACTOR B SUBUNIT 2, TFB2.

Mol	Chain	Residues	Atoms			AltConf	Trace	
25	W	62	Total 518	C 334	N 83	O 101	0	0

• Molecule 26 is a protein called RNA POLYMERASE II TRANSCRIPTION FACTOR B SUBUNIT 5, TFB5.



Mol	Chain	Residues	Atoms				AltConf	Trace	
26	Х	63	Total 499	C 316	N 88	O 93	$\begin{array}{c} \mathrm{S} \\ \mathrm{2} \end{array}$	0	0

• Molecule 27 is a protein called DNA REPAIR HELICASE RAD3.

Mol	Chain	Residues			Atoms					
27	Y	578	Total 5351	C 2975	Н 691	N 790	O 862	S 33	0	0

• Molecule 28 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	AltConf
28	2	1	Total Mg 1 1	0
28	А	1	Total Mg 1 1	0

• Molecule 29 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	AltConf
29	2	1	Total Zn 1 1	0
29	А	2	Total Zn 2 2	0
29	В	1	Total Zn 1 1	0
29	С	1	Total Zn 1 1	0
29	Ι	2	Total Zn 2 2	0
29	J	1	Total Zn 1 1	0
29	L	1	Total Zn 1 1	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA REPAIR HELICASE RAD25, SSL2

















 $\bullet$  Molecule 7: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC1, RPB5



 $\bullet$  Molecule 8: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC2, RPB6





 $\bullet$  Molecule 10: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC3, RPB8











• Molecule 19: TATA-BOX-BINDING PROTEIN, TBP





• Molecule 22: TEMPLATE STRAND DNA





#### ASP GLU ASP GLU ASP GLU CLL MET GLU GLU



# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	7578	Depositor
Resolution determination method	Not provided	
CTF correction method	SPARX	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	22500	Depositor
Image detector	GATAN K2 (4k x 4k)	Depositor
Maximum map value	9.151	Depositor
Minimum map value	-0.003	Depositor
Average map value	0.142	Depositor
Map value standard deviation	0.652	Depositor
Recommended contour level	3.99	Depositor
Map size (Å)	399.76, 399.76, 399.76	wwPDB
Map dimensions	152, 152, 152	wwPDB
Map angles $(^{\circ})$	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.63, 2.63, 2.63	Depositor



# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	B	ond lengths	E	Bond angles
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	1	4.34	6/3867~(0.2%)	2.05	41/5221~(0.8%)
2	2	1.31	4/571~(0.7%)	1.64	7/765~(0.9%)
3	А	0.57	4/11374~(0.0%)	0.81	11/15384~(0.1%)
4	В	0.49	1/9316~(0.0%)	0.74	4/12564~(0.0%)
5	С	0.49	0/2133	0.78	2/2891~(0.1%)
6	D	0.51	0/1444	0.83	2/1935~(0.1%)
7	Е	0.48	0/1788	0.72	0/2406
8	F	0.62	0/691	0.81	0/933
9	G	0.52	0/1368	0.81	0/1844
10	Н	0.51	0/1086	0.80	0/1470
11	Ι	0.47	0/989	0.78	0/1331
12	J	0.54	0/541	0.88	1/727~(0.1%)
13	Κ	0.47	0/938	0.71	0/1267
14	L	0.54	0/365	0.95	0/485
15	М	0.40	0/970	0.57	0/1310
16	Ν	1.27	20/1692~(1.2%)	1.41	34/2613~(1.3%)
17	0	0.42	0/800	0.63	0/1080
18	Р	4.21	7/2297~(0.3%)	2.23	41/3090~(1.3%)
19	Q	0.41	0/1443	0.62	0/1942
20	R	9.51	2/978~(0.2%)	3.28	11/1321~(0.8%)
21	S	5.35	7/1017~(0.7%)	3.30	36/1356~(2.7%)
22	Т	1.38	45/1618~(2.8%)	1.50	49/2491~(2.0%)
23	U	0.93	4/1273~(0.3%)	0.92	9/1710~(0.5%)
24	V	1.09	3/1481~(0.2%)	0.88	10/1988~(0.5%)
25	W	0.67	0/527	0.68	0/704
26	Х	0.60	0/504	0.69	0/679
27	Y	0.88	11/4735~(0.2%)	0.84	19/6369~(0.3%)
All	All	2.13	114/55806~(0.2%)	1.23	277/75876~(0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected



Mol	Chain	#Chirality outliers	#Planarity outliers
1	1	1	1
2	2	0	2
3	А	0	2
4	В	0	1
18	Р	2	0
20	R	1	0
21	S	0	1
27	Y	0	2
All	All	4	9

by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

The worst 5 of 114 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
20	R	155	PRO	N-CD	213.23	4.46	1.47
20	R	119	PRO	N-CD	204.92	4.34	1.47
1	1	753	PRO	N-CD	203.03	4.32	1.47
18	Р	292	PRO	N-CD	198.01	4.25	1.47
1	1	746	PRO	N-CD	168.34	3.83	1.47

The worst 5 of 277 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
20	R	155	PRO	CA-N-CD	-66.71	18.11	111.50
20	R	119	PRO	N-CD-CG	-58.68	15.18	103.20
21	S	189	PRO	CA-N-CD	-53.80	36.18	111.50
1	1	753	PRO	N-CD-CG	-48.67	30.19	103.20
18	Р	292	PRO	N-CD-CG	-48.02	31.17	103.20

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	1	690	ILE	CB
18	Р	269	ILE	CB
18	Р	277	ILE	CB
20	R	126	ILE	CB

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	1	661	SER	Peptide



Mol	Chain	Res	Type	Group
2	2	269	PHE	Peptide,Sidechain
3	А	234	MET	Mainchain
3	А	95	PHE	Mainchain
4	В	43	LEU	Mainchain

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	3797	0	3727	916	0
2	2	666	0	546	251	0
3	А	11174	0	11217	712	0
4	В	9140	0	9107	421	0
5	С	2095	0	2051	43	0
6	D	1434	0	1460	69	0
7	Е	1752	0	1776	30	0
8	F	679	0	699	94	0
9	G	1340	0	1353	174	0
10	Н	1068	0	1040	21	0
11	Ι	971	0	927	16	0
12	J	532	0	542	14	0
13	Κ	920	0	929	19	0
14	L	363	0	386	24	0
15	М	956	0	916	32	0
16	Ν	1500	0	805	235	0
17	0	792	0	806	49	0
18	Р	2269	0	2346	1242	0
19	Q	1416	0	1491	154	0
20	R	960	0	969	345	0
21	S	1005	0	1024	580	0
22	Т	1452	0	822	273	0
23	U	1248	0	1229	236	0
24	V	1455	0	1465	465	0
25	W	518	0	514	27	0
26	Х	499	0	525	12	0
27	Y	4660	691	4743	807	0
28	2	1	0	0	0	0
28	А	1	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
29	2	1	0	0	0	0
29	А	2	0	0	0	0
29	В	1	0	0	0	0
29	С	1	0	0	0	0
29	Ι	2	0	0	0	0
29	J	1	0	0	0	0
29	L	1	0	0	0	0
All	All	54672	691	53415	6017	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 56.

The worst 5 of 6017 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom_1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
22:T:134:DT:C6	24:V:323:ARG:HD2	1.24	1.69
18:P:279:VAL:HG23	18:P:302:LEU:CG	1.19	1.66
20:R:34:PHE:CE2	20:R:134:LEU:HD11	1.25	1.65
18:P:137:CYS:SG	18:P:142:LEU:HB3	1.34	1.64
27:Y:132:LYS:HD3	27:Y:155:LEU:CD1	1.24	1.64

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	1	467/492~(95%)	436 (93%)	26~(6%)	5 (1%)	14 52
2	2	68/174~(39%)	51 (75%)	10 (15%)	7 (10%)	0 8
3	А	1414/1733 (82%)	1250 (88%)	113 (8%)	51 (4%)	3 25
4	В	1140/1224 (93%)	1018 (89%)	86 (8%)	36 (3%)	4 26



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
5	С	264/266~(99%)	242 (92%)	20 (8%)	2(1%)	19	60
6	D	174/178~(98%)	148 (85%)	18 (10%)	8 (5%)	2	21
7	Е	212/214~(99%)	195 (92%)	13 (6%)	4 (2%)	8	38
8	F	82/84~(98%)	75~(92%)	7 (8%)	0	100	100
9	G	169/171~(99%)	158 (94%)	8 (5%)	3(2%)	8	40
10	Н	129/146~(88%)	106 (82%)	14 (11%)	9~(7%)	1	14
11	Ι	117/119~(98%)	98 (84%)	16 (14%)	3(3%)	5	31
12	J	63/65~(97%)	51 (81%)	9 (14%)	3~(5%)	2	20
13	К	113/115~(98%)	109 (96%)	4 (4%)	0	100	100
14	L	44/46~(96%)	27 (61%)	9 (20%)	8 (18%)	0	2
15	М	110/116~(95%)	103 (94%)	7 (6%)	0	100	100
17	Ο	97/122~(80%)	93 (96%)	4 (4%)	0	100	100
18	Р	287/345~(83%)	235 (82%)	33 (12%)	19 (7%)	1	15
19	Q	178/180 (99%)	170 (96%)	5 (3%)	3 (2%)	9	42
20	R	112/160~(70%)	100 (89%)	10 (9%)	2 (2%)	8	40
21	S	105/123~(85%)	94 (90%)	7 (7%)	4 (4%)	3	24
23	U	142/150~(95%)	131 (92%)	6 (4%)	5 (4%)	3	25
24	V	166/174~(95%)	145 (87%)	16 (10%)	5 (3%)	4	28
25	W	60/62~(97%)	60 (100%)	0	0	100	100
26	X	61/63~(97%)	58 (95%)	3 (5%)	0	100	100
27	Y	558/778 (72%)	515 (92%)	35 (6%)	8 (1%)	11	46
All	All	6332/7300 (87%)	5668 (90%)	479 (8%)	185 (3%)	7	29

5 of 185 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1	447	GLN
1	1	458	SER
2	2	275	LYS
2	2	292	PRO
3	А	74	MET



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	1	415/431~(96%)	370~(89%)	45 (11%)	6	24
2	2	62/151~(41%)	55~(89%)	7 (11%)	6	22
3	А	1240/1520~(82%)	1066 (86%)	174 (14%)	3	17
4	В	985/1061 (93%)	868 (88%)	117 (12%)	5	20
5	С	234/234~(100%)	206 (88%)	28 (12%)	5	20
6	D	160/160~(100%)	129 (81%)	31 (19%)	1	8
7	Е	196/196~(100%)	175 (89%)	21 (11%)	6	24
8	F	74/74~(100%)	67 (90%)	7 (10%)	8	28
9	G	152/152~(100%)	135 (89%)	17 (11%)	6	22
10	Н	117/128 (91%)	103 (88%)	14 (12%)	5	20
11	Ι	113/113~(100%)	106 (94%)	7 (6%)	18	43
12	J	60/60~(100%)	49 (82%)	11 (18%)	1	10
13	Κ	99/99~(100%)	87 (88%)	12 (12%)	5	20
14	L	40/40 (100%)	27~(68%)	13 (32%)	0	2
15	М	107/107~(100%)	104 (97%)	3 (3%)	43	65
17	О	91/108 (84%)	83 (91%)	8 (9%)	10	31
18	Р	256/299~(86%)	215 (84%)	41 (16%)	2	13
19	Q	152/152~(100%)	143 (94%)	9 (6%)	19	45
20	R	108/149~(72%)	88 (82%)	20 (18%)	1	9
21	S	116/118 (98%)	69~(60%)	47 (40%)	0	0
23	U	136/136~(100%)	133 (98%)	3 (2%)	52	71
24	V	163/163~(100%)	158 (97%)	5 (3%)	40	62
25	W	57/57~(100%)	57 (100%)	0	100	100
26	Х	57/57~(100%)	57 (100%)	0	100	100
27	Y	521/707 (74%)	504 (97%)	17 (3%)	38	61
All	All	5711/6472~(88%)	5054 (88%)	657 (12%)	9	21



5 of 657 residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
11	Ι	35	VAL
20	R	133	GLN
12	J	42	LYS
11	Ι	31	THR
18	Р	255	SER

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 109 such side chains are listed below:

Mol	Chain	$\mathbf{Res}$	Type
10	Н	35	GLN
18	Р	84	ASN
27	Y	60	GLN
11	Ι	83	ASN
15	М	59	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
21	S	7
27	Y	7
24	V	3
23	U	3
15	М	2
3	А	2
6	D	1
1	1	1
4	В	1
2	2	1

The worst 5 of 28 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	V	227:SER	С	291:SER	Ν	84.31
1	М	61:ASN	С	228:ASP	Ν	43.11
1	D	76:LYS	С	118:THR	Ν	35.68
1	S	184:TYR	С	185:ASP	Ν	14.79
1	U	313:ALA	С	323:GLY	N	13.13



# 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-3114. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

## 6.1 Orthogonal projections (i)

#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

## 6.2 Central slices (i)

#### 6.2.1 Primary map



X Index: 76



Y Index: 76



Z Index: 76

The images above show central slices of the map in three orthogonal directions.

### 6.3 Largest variance slices (i)

#### 6.3.1 Primary map



X Index: 69

Y Index: 99

Z Index: 88

The images above show the largest variance slices of the map in three orthogonal directions.

#### 6.4 Orthogonal surface views (i)

#### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.99. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



# 6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



# 7 Map analysis (i)

This section contains the results of statistical analysis of the map.

## 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



## 7.2 Volume estimate (i)



The volume at the recommended contour level is  $637 \text{ nm}^3$ ; this corresponds to an approximate mass of 575 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



## 7.3 Rotationally averaged power spectrum (i)



\*Reported resolution corresponds to spatial frequency of 0.167  ${\rm \AA^{-1}}$ 



# 8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



# 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-3114 and PDB model 5FMF. Per-residue inclusion information can be found in section 3 on page 9.

## 9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 3.99 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



### 9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

#### 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3.99).



## 9.4 Atom inclusion (i)



At the recommended contour level, 46% of all backbone atoms, 44% of all non-hydrogen atoms, are inside the map.



1.0

0.0 <0.0

## 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (3.99) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4368	0.0620
1	0.2962	0.0410
2	0.2866	0.0410
А	0.5002	0.0770
В	0.4001	0.0490
С	0.6132	0.0640
D	0.3741	0.0700
Ε	0.5307	0.0900
F	0.6778	0.0690
G	0.4183	0.0690
Н	0.5760	0.0720
Ι	0.4408	0.0600
J	0.5841	0.0660
K	0.5138	0.0540
L	0.6553	0.0600
М	0.3348	0.0730
Ν	0.3933	0.0730
0	0.2420	0.0580
Р	0.1213	0.0240
Q	0.4330	0.0640
R	0.4707	0.0730
S	0.5091	0.1020
Т	0.5021	0.1110
U	0.3739	0.0320
V	0.3862	0.0590
W	0.0923	0.0590
Х	0.1870	0.0240
Y	0.5868	0.0570

