



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 8, 2024 – 04:33 am GMT

PDB ID : 6FM4
Title : The crystal structure of *S. aureus* Gyrase complex with ID-130 and DNA
Authors : Ombrato, R.; Garofalo, B.; Mangano, G.; Mancini, F.
Deposited on : 2018-01-30
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	B	692	74% (green), 22% (yellow), .. (grey)
1	D	692	76% (green), 19% (yellow), .. (grey)
2	E	19	58% (green), 42% (yellow)
3	F	18	67% (green), 33% (yellow)

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 11536 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA gyrase subunit B,DNA gyrase subunit B,DNA gyrase subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	672	Total 5324	C 3318	N 958	O 1023	S 25	0	1	0
1	D	670	Total 5291	C 3298	N 952	O 1016	S 25	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	409	MET	LEU	conflict	UNP P66937
B	544	THR	-	linker	UNP P66937
B	545	GLY	-	linker	UNP P66937
B	1123	PHE	TYR	engineered mutation	UNP Q99XG5
D	409	MET	LEU	conflict	UNP P66937
D	544	THR	-	linker	UNP P66937
D	545	GLY	-	linker	UNP P66937
D	1123	PHE	TYR	engineered mutation	UNP Q99XG5

- Molecule 2 is a DNA chain called DNA (5'-5UA*D(P*GP*CP*CP*GP*TP*AP*GP*GP*G P*CP*CP*CP*TP*AP*CP*GP*GP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	E	19	Total 389	C 183	N 72	O 115	P 19	0	0	0


- Molecule 3 is a DNA chain called DNA (5'-5UA*D(P*GP*CP*CP*GP*TP*AP*GP*GP*G P*CP*CP*CP*TP*AP*CP*GP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	F	18	Total 369	C 173	N 70	O 108	P 18	0	0	0

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mn	1	0
			1	1		
4	E	1	Total	Mn	1	0
			1	1		
4	F	1	Total	Mn	1	0
			1	1		

- Molecule 5 is 5'-O-CARBOXY-2'-DEOXYADENOSINE (three-letter code: 5UA) (formula: C₁₁H₁₃N₅O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	E	1	Total	C	N	O	21	0
			21	11	5	5		
5	F	1	Total	C	N	O	21	0
			21	11	5	5		

- Molecule 6 is {N}-[3-(4-isoquinolin-1-ylpiperazin-1-yl)propyl]benzamide (three-letter code: DU5) (formula: C₂₃H₂₆N₄O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	E	1	56	46	8	2	0	1


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	21	Total	O	0	0
			21	21		
7	D	37	Total	O	0	0
			37	37		
7	E	3	Total	O	0	0
			3	3		
7	F	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA gyrase subunit B,DNA gyrase subunit B,DNA gyrase subunit A



- Molecule 1: DNA gyrase subunit B,DNA gyrase subunit B,DNA gyrase subunit A





- Molecule 2: DNA (5'-5UA*D(P*GP*CP*CP*GP*TP*AP*GP*GP*GP*CP*CP*CP*TP*AP*CP*GP*GP*CP*T)-3')



- Molecule 3: DNA (5'-5UA*D(P*GP*CP*CP*GP*TP*AP*GP*GP*GP*CP*CP*CP*TP*AP*CP*GP*GP*C)-3')



4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	92.95Å 92.95Å 407.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.05 – 2.70 39.43 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.05-2.70) 99.8 (39.43-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.23 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.225 , 0.300 0.226 , 0.297	Depositor DCC
R_{free} test set	2750 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 3.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.31$, $\langle L^2 \rangle = 0.14$	Xtrriage
Estimated twinning fraction	0.379 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11536	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, 5UA, DU5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.69	0/5395	0.89	5/7267 (0.1%)
1	D	0.69	0/5362	0.89	5/7227 (0.1%)
2	E	0.60	0/435	0.93	0/669
3	F	0.58	0/413	0.90	1/635 (0.2%)
All	All	0.68	0/11605	0.89	11/15798 (0.1%)

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1484	ARG	NE-CZ-NH2	-6.05	117.28	120.30
1	B	1470	ARG	NE-CZ-NH2	-6.00	117.30	120.30
1	B	1485	ARG	NE-CZ-NH1	5.95	123.27	120.30
3	F	3	DC	C1'-O4'-C4'	-5.66	104.44	110.10
1	D	1122	ARG	NE-CZ-NH1	5.48	123.04	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	5324	0	5356	110	0
1	D	5291	0	5308	93	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	389	0	213	7	0
3	F	369	0	201	6	0
4	B	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	E	21	0	11	0	1
5	F	21	0	11	0	0
6	E	56	0	0	6	0
7	B	21	0	0	8	0
7	D	37	0	0	9	0
7	E	3	0	0	1	0
7	F	1	0	0	2	0
All	All	11536	0	11100	204	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 204 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:435:GLU:OE2	1:B:508:ASP:OD2	1.81	0.96
6:E:103[A]:DU5:CAR	6:E:103[A]:DU5:CAK	2.50	0.89
1:D:427:GLU:HA	7:D:1619:HOH:O	1.77	0.83
6:E:103[B]:DU5:CAK	6:E:103[B]:DU5:CAR	2.56	0.82
1:D:1297:SER:HG	1:D:1300:THR:HG1	1.22	0.80

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1393:GLU:OE2	5:E:102:5UA:N1[1_655]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	667/692 (96%)	610 (92%)	51 (8%)	6 (1%)	17	40
1	D	664/692 (96%)	612 (92%)	46 (7%)	6 (1%)	17	40
All	All	1331/1384 (96%)	1222 (92%)	97 (7%)	12 (1%)	17	40

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1033	ARG
1	B	1057	GLY
1	B	1119	ALA
1	B	1327	LEU
1	D	1142	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	573/591 (97%)	566 (99%)	7 (1%)	71	88
1	D	568/591 (96%)	555 (98%)	13 (2%)	50	78
All	All	1141/1182 (96%)	1121 (98%)	20 (2%)	59	83

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	1329	THR
1	D	1414	ARG
1	D	1465	LEU
1	D	1450	ASN
1	B	1401	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	1334	ASN
1	D	1390	HIS
1	B	1368	GLN
1	D	476	ASN
1	D	480	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	DU5	E	103[A]	-	31,31,31	1.37	3 (9%)	40,41,41	2.03	10 (25%)
5	5UA	F	1502	3	21,23,23	2.76	3 (14%)	20,33,33	2.52	6 (30%)
5	5UA	E	102	-	21,23,23	2.70	3 (14%)	20,33,33	2.79	6 (30%)
6	DU5	E	103[B]	-	31,31,31	1.36	3 (9%)	40,41,41	2.00	11 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	DU5	E	103[A]	-	-	5/15/25/25	0/4/4/4
5	5UA	F	1502	3	-	1/4/21/21	0/3/3/3
5	5UA	E	102	-	-	3/4/21/21	0/3/3/3
6	DU5	E	103[B]	-	-	6/15/25/25	0/4/4/4

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	1502	5UA	C2-N3	8.79	1.46	1.32
5	E	102	5UA	C2-N3	8.26	1.45	1.32
5	F	1502	5UA	C2-N1	6.61	1.46	1.33
5	E	102	5UA	C2-N1	6.49	1.46	1.33
5	E	102	5UA	O5'-C6'	5.40	1.48	1.33

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	102	5UA	N3-C2-N1	-8.76	114.98	128.68
5	F	1502	5UA	N3-C2-N1	-8.68	115.10	128.68
6	E	103[A]	DU5	CAO-NBA-CAQ	-5.90	96.14	111.23
5	E	102	5UA	O5'-C5'-C4'	5.84	120.80	108.43
6	E	103[B]	DU5	CAO-NBA-CAQ	-5.61	96.89	111.23

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	E	102	5UA	O4'-C4'-C5'-O5'
5	E	102	5UA	C3'-C4'-C5'-O5'
6	E	103[A]	DU5	CAZ-CAX-NBB-CAR
6	E	103[A]	DU5	NAT-CAX-NBB-CAR
6	E	103[B]	DU5	CAZ-CAX-NBB-CAR


There are no ring outliers.


3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	103[A]	DU5	3	0
5	E	102	5UA	0	1
6	E	103[B]	DU5	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	1
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	545:GLY	C	580:TYR	N	3.21
1	B	545:GLY	C	580:TYR	N	3.09

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 5UA E 102:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 5UA F 1502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DU5 E 103 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.