



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 04:36 pm BST

PDB ID : 1FPN
Title : HUMAN RHINOVIRUS SEROTYPE 2 (HRV2)
Authors : Verdaguer, N.; Blaas, D.; Fita, I.
Deposited on : 2000-08-31
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	1	289	68% (green), 21% (yellow), 7% (orange), 4% (red), 2% (grey)
2	2	261	76% (green), 15% (yellow), 7% (orange), 2% (red), 2% (grey)
3	3	237	76% (green), 19% (yellow), 5% (orange), 2% (red), 2% (grey)
4	4	68	28% (green), 7% (yellow), 63% (grey)

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6316 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called COAT PROTEIN VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	1	269	2154	1356	378	409	11	0	0	0

- Molecule 2 is a protein called COAT PROTEIN VP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	2	250	1961	1245	339	369	8	8	0	0


- Molecule 3 is a protein called COAT PROTEIN VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	3	237	1834	1172	304	346	12	0	0	0

- Molecule 4 is a protein called COAT PROTEIN VP4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	4	25	194	122	35	37	0	0	0

- Molecule 5 is LAURIC ACID (three-letter code: DAO) (formula: C₁₂H₂₄O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	1	1	Total	C O	0	0
			14	12 2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	1	51	Total	O	0	0
			51	51		
6	2	49	Total	O	0	0
			49	49		
6	3	56	Total	O	0	0
			56	56		
6	4	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: COAT PROTEIN VP1

Chain 1: 




- Molecule 2: COAT PROTEIN VP2

Chain 2: 



- Molecule 3: COAT PROTEIN VP3

Chain 3: 



- Molecule 4: COAT PROTEIN VP4

Chain 4: 

GLY	A2	Q3	V4	S5	R6	Q7	ASN	VAL	GLY	THR	HIS	SER	THR	GLN	ASN	SER	VAL	SER	ASN	GLY	SER	SER	SER	LEU	N25	Y31	F32	K33	L43	GLU	PHE	THR	GLN	ASP	PRO	SER	SER	LYS	PHE	THR	ASP	PRO	VAL	VAL	LYS	ASP	VAL	LEU	GLU	LYS	GLY	ILE	PRO	THR	LEU	GLN
-----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	308.68Å 352.98Å 380.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.60)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.180 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6316	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
DAO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1	0.69	1/2210 (0.0%)	0.80	1/3010 (0.0%)
2	2	0.63	2/2016 (0.1%)	0.94	4/2752 (0.1%)
3	3	0.67	1/1884 (0.1%)	0.85	3/2579 (0.1%)
4	4	0.80	0/196	0.89	0/261
All	All	0.67	4/6306 (0.1%)	0.87	8/8602 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	2	261	GLN	CG-CD	5.87	1.64	1.51
1	1	199	ASP	CB-CG	-5.52	1.40	1.51
2	2	14	ILE	CA-CB	5.51	1.67	1.54
3	3	217	ASP	CB-CG	-5.39	1.40	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	2	261	GLN	N-CA-CB	-15.18	83.27	110.60
2	2	29	ALA	N-CA-C	-9.48	85.40	111.00
3	3	220	LEU	CA-CB-CG	6.51	130.28	115.30
2	2	181	LEU	C-N-CA	-6.24	109.20	122.30
1	1	199	ASP	CB-CA-C	-6.00	98.39	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2154	0	2077	63	0
2	2	1961	0	1902	35	0
3	3	1834	0	1817	47	0
4	4	194	0	180	2	0
5	1	14	0	23	1	0
6	1	51	0	0	0	0
6	2	49	0	0	0	0
6	3	56	0	0	1	0
6	4	3	0	0	0	0
All	All	6316	0	5999	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:1:268:ARG:HB3	1:1:268:ARG:HH11	1.10	1.09
1:1:268:ARG:CB	1:1:268:ARG:HH11	1.69	1.04
3:3:87:THR:HG22	3:3:185:SER:HB2	1.47	0.97
1:1:199:ASP:HB3	1:1:201:GLN:H	1.32	0.94
1:1:268:ARG:HB3	1:1:268:ARG:NH1	1.87	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	267/289 (92%)	256 (96%)	11 (4%)	0	100	100
2	2	248/261 (95%)	228 (92%)	17 (7%)	3 (1%)	13	27
3	3	235/237 (99%)	222 (94%)	13 (6%)	0	100	100
4	4	21/68 (31%)	21 (100%)	0	0	100	100
All	All	771/855 (90%)	727 (94%)	41 (5%)	3 (0%)	34	57

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	2	160	THR
2	2	31	ALA
2	2	164	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	237/257 (92%)	213 (90%)	24 (10%)	7	14
2	2	216/226 (96%)	198 (92%)	18 (8%)	11	22
3	3	210/210 (100%)	183 (87%)	27 (13%)	4	7
4	4	19/59 (32%)	14 (74%)	5 (26%)	0	1
All	All	682/752 (91%)	608 (89%)	74 (11%)	6	11

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	2	161	ARG
2	2	257	ARG
3	3	230	LEU
2	2	163	ASN
2	2	211	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	2	197	ASN
2	2	220	ASN
3	3	193	GLN
2	2	193	ASN
3	3	227	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DAO	1	6001	-	10,13,13	1.45	0	9,13,13	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	DAO	1	6001	-	-	3/9/11/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	1	6001	DAO	C1-C2-C3-C4
5	1	6001	DAO	C6-C7-C8-C9
5	1	6001	DAO	C5-C6-C7-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	1	6001	DAO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.