



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 25, 2022 – 07:34 PM EDT

PDB ID : 7FPD
Title : PanDDA analysis group deposition – Aar2/RNaseH in complex with fragment P09E01 from the F2X-Universal Library
Authors : Barthel, T.; Wollenhaupt, J.; Lima, G.M.A.; Wahl, M.C.; Weiss, M.S.
Deposited on : 2022-08-26
Resolution : 1.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.31.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4693 (1.64-1.60)
Clashscore	141614	5002 (1.64-1.60)
Ramachandran outliers	138981	4888 (1.64-1.60)
Sidechain outliers	138945	4887 (1.64-1.60)
RSRZ outliers	127900	4609 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	 26% 79% 11% 8%
2	B	308	 20% 89% 8% ..

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9225 atoms, of which 4524 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	237	4068	1287	2060	336	373	12	18	21	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	300	5044	1654	2464	421	485	20	33	17	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	LEU	conflict	UNP P32357
B	167	SER	LYS	conflict	UNP P32357
B	?	-	LEU	deletion	UNP P32357
B	?	-	GLN	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	GLY	deletion	UNP P32357
B	?	-	SER	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	MET	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	?	-	ALA	deletion	UNP P32357
B	?	-	LYS	deletion	UNP P32357
B	?	-	ASN	deletion	UNP P32357
B	?	-	GLU	deletion	UNP P32357
B	170	SER	ASP	conflict	UNP P32357

- Molecule 3 is 1-[(2S)-2-(furan-2-yl)azepan-1-yl]ethan-1-one (three-letter code: VF9) (formula: C₁₂H₁₇NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	15	12	1	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	57	Total	O	0	0
			57	57		
4	B	41	Total	O	0	0
			41	41		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.55Å 81.77Å 93.21Å 90.00° 108.27° 90.00°	Depositor
Resolution (Å)	42.04 – 1.61 44.59 – 1.61	Depositor EDS
% Data completeness (in resolution range)	99.1 (42.04-1.61) 99.2 (44.59-1.61)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.98 (at 1.62Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.283 , 0.321 0.289 , 0.345	Depositor DCC
R_{free} test set	2100 reflections (2.61%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtrriage
Anisotropy	0.309	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9225	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VF9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	3/2149 (0.1%)	0.87	4/2911 (0.1%)
2	B	0.70	1/2739 (0.0%)	0.80	4/3699 (0.1%)
All	All	0.73	4/4888 (0.1%)	0.84	8/6610 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	248	GLU	CB-CG	5.70	1.62	1.52
1	A	1992	TYR	CD1-CE1	5.23	1.47	1.39
1	A	2002[A]	TYR	CD2-CE2	-5.21	1.31	1.39
1	A	2002[B]	TYR	CD2-CE2	-5.21	1.31	1.39

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2008	LEU	CB-CG-CD1	-7.46	98.33	111.00
1	A	1847	ASP	CB-CG-OD2	-6.48	112.47	118.30
1	A	1924	LEU	CB-CG-CD2	-6.01	100.77	111.00
1	A	1994	ASP	CB-CG-OD1	5.69	123.42	118.30
2	B	251	CYS	CA-CB-SG	-5.35	104.38	114.00
2	B	18	ASP	CB-CG-OD2	-5.19	113.63	118.30
2	B	265	ASP	CB-CG-OD2	5.16	122.95	118.30
2	B	80	MET	CA-CB-CG	-5.08	104.66	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2008	2060	1974	15	0
2	B	2580	2464	2398	18	0
3	A	15	0	0	0	0
4	A	57	0	0	1	0
4	B	41	0	0	2	0
All	All	4701	4524	4372	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1895:HIS:O	1:A:1898[A]:VAL:HG22	1.85	0.77
1:A:1976:ASP:HB3	4:A:2204:HOH:O	1.87	0.74
2:B:80:MET:SD	2:B:137:PHE:CG	2.80	0.74
1:A:1862:VAL:HG22	1:A:1872:THR:HG22	1.75	0.68
1:A:1962:ARG:O	1:A:2013:ARG:NH1	2.29	0.66
2:B:80:MET:SD	2:B:137:PHE:CD2	2.89	0.66
1:A:2064:GLY:O	1:A:2068:ASN:N	2.25	0.64
1:A:2061:THR:O	1:A:2064:GLY:N	2.32	0.63
2:B:287:ARG:O	2:B:291:ILE:HD13	2.02	0.59
2:B:1:MET:HB3	2:B:35:ASP:HA	1.87	0.56
2:B:77:LEU:HD21	2:B:79:LYS:HE3	1.89	0.55
2:B:1:MET:N	4:B:401:HOH:O	2.40	0.54
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.57	0.53
1:A:1911:TRP:CD2	2:B:195:MET:HB2	2.45	0.51
1:A:1946:VAL:O	1:A:1949:LEU:HG	2.14	0.48
2:B:50:ASP:OD1	2:B:51:ASN:N	2.48	0.47
2:B:70:GLN:HB3	2:B:81:MET:HE1	1.97	0.47
2:B:53:SER:HB2	4:B:433:HOH:O	2.15	0.46
1:A:1998:ARG:NH1	1:A:2045:ASP:OD2	2.45	0.45
2:B:190:ARG:HG3	2:B:203[B]:TYR:CE2	2.52	0.44
1:A:1859:ARG:HH12	1:A:1979[A]:MET:CE	2.30	0.44
2:B:43:VAL:HG13	2:B:43:VAL:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1844:PHE:O	1:A:1885:LYS:NZ	2.26	0.43
1:A:1863:HIS:CE1	1:A:1871:ALA:HB3	2.53	0.43
2:B:190:ARG:HG3	2:B:203[B]:TYR:CZ	2.53	0.43
2:B:110:GLU:O	2:B:110:GLU:HG2	2.20	0.42
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	2.02	0.42
2:B:251:CYS:HG	2:B:292:CYS:HB3	1.84	0.42
1:A:1862:VAL:HG13	1:A:1870:VAL:CG1	2.49	0.41
2:B:43:VAL:HA	2:B:58:TYR:O	2.20	0.41
2:B:80:MET:CG	2:B:81:MET:N	2.83	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/258 (100%)	252 (98%)	6 (2%)	0	100	100
2	B	315/308 (102%)	303 (96%)	12 (4%)	0	100	100
All	All	573/566 (101%)	555 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/233 (102%)	231 (98%)	6 (2%)	47	21
2	B	294/284 (104%)	292 (99%)	2 (1%)	84	72
All	All	531/517 (103%)	523 (98%)	8 (2%)	76	43

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1931[A]	LYS
1	A	1931[B]	LYS
1	A	1962	ARG
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
2	B	258	LYS
2	B	281	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1863	HIS
2	B	135	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	VF9	A	2101	-	11,16,16	3.23	2 (18%)	13,21,21	1.57	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	VF9	A	2101	-	-	1/4/20/20	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2101	VF9	C9-C8	-9.01	1.27	1.39
3	A	2101	VF9	C2-N	-5.11	1.39	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2101	VF9	C9-C8-C7	2.84	132.51	129.05
3	A	2101	VF9	C3-C2-N	-2.25	109.34	113.09

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2101	VF9	C-C1-N-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/258 (91%)	1.59	68 (28%) 0 0	37, 68, 136, 195	0
2	B	300/308 (97%)	1.35	63 (21%) 1 0	36, 66, 129, 241	0
All	All	537/566 (94%)	1.46	131 (24%) 0 0	36, 67, 132, 241	0

All (131) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	53	SER	14.3
1	A	2068	ASN	11.0
2	B	52	SER	9.4
2	B	1	MET	8.6
1	A	2037	TYR	7.9
1	A	2039	LEU	7.5
1	A	2027	LEU	7.2
1	A	2040	TRP	6.9
1	A	2042	SER	6.8
1	A	2065	ARG	6.6
1	A	2064	GLY	6.4
1	A	2063	TYR	6.2
1	A	1833	GLY	5.9
1	A	2069	VAL	5.9
1	A	2048	TRP	5.9
1	A	1979[A]	MET	5.5
1	A	2034	ILE	5.5
2	B	54[A]	MET	5.5
2	B	108	ILE	4.9
2	B	109	ASP	4.8
1	A	2060	LEU	4.6
2	B	313	TYR	4.6
1	A	2036	SER	4.5
2	B	317	LEU	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2032	ILE	4.4
2	B	277	GLU	4.3
2	B	110	GLU	4.2
2	B	173	ALA	4.1
2	B	50	ASP	4.0
1	A	2038	HIS	4.0
1	A	2028	SER	4.0
2	B	80	MET	4.0
2	B	172	PRO	3.9
1	A	2041	PRO	3.9
1	A	1865	THR	3.8
1	A	1931[A]	LYS	3.8
2	B	51	ASN	3.8
2	B	246	MET	3.8
2	B	283	LEU	3.7
1	A	2022	ALA	3.7
2	B	84	ARG	3.6
2	B	111	ASP	3.6
2	B	174	HIS	3.6
1	A	1927	GLU	3.6
2	B	275	LEU	3.5
2	B	295	SER	3.5
1	A	1870	VAL	3.5
1	A	1886	THR	3.4
2	B	103	VAL	3.4
1	A	1840	TYR	3.4
2	B	106	PRO	3.4
2	B	181	ILE	3.3
1	A	1997	ASP	3.3
1	A	1834	ALA	3.3
2	B	294	TYR	3.3
1	A	1838	SER	3.3
2	B	292	CYS	3.3
2	B	107	LYS	3.2
1	A	1989	PHE	3.2
2	B	22	PHE	3.2
2	B	101	MET	3.2
2	B	91	ASN	3.1
1	A	1995	TRP	3.1
2	B	122[A]	GLN	3.1
1	A	1962	ARG	3.1
2	B	147	VAL	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	152	LEU	3.0
2	B	301	SER	3.0
2	B	254	ALA	3.0
1	A	2001[A]	SER	3.0
2	B	171	ASP	2.9
1	A	2035	LYS	2.9
1	A	2056[A]	ARG	2.9
1	A	1837	SER	2.9
1	A	1866	PHE	2.9
1	A	1878	CYS	2.9
1	A	2061	THR	2.9
2	B	293	LEU	2.9
1	A	2053[A]	SER	2.9
2	B	203[A]	TYR	2.8
1	A	2043	PHE	2.8
1	A	1862	VAL	2.8
2	B	102	MET	2.8
1	A	1872	THR	2.8
2	B	6	PHE	2.8
2	B	274	THR	2.8
1	A	2015	LEU	2.7
2	B	281	ASP	2.7
1	A	2051	ILE	2.7
1	A	1969	MET	2.7
2	B	279	TYR	2.6
2	B	196	GLU	2.6
2	B	193	HIS	2.6
1	A	1924	LEU	2.5
2	B	312	LYS	2.5
1	A	2033	THR	2.5
2	B	96	PHE	2.5
1	A	1964	PRO	2.5
1	A	2059	ILE	2.5
2	B	189	ILE	2.5
2	B	303	HIS	2.5
2	B	308	ILE	2.5
1	A	2067	TYR	2.5
2	B	87	ALA	2.5
1	A	2017[A]	THR	2.5
1	A	1934	ILE	2.4
2	B	316	LEU	2.4
1	A	1880	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2002[A]	TYR	2.4
1	A	2044	THR	2.4
2	B	121	VAL	2.3
2	B	146	THR	2.3
2	B	150	ASN	2.3
1	A	2024[A]	MET	2.3
2	B	89	PHE	2.3
1	A	1869	ASN	2.3
1	A	1998	ARG	2.2
1	A	1952	PRO	2.2
2	B	92	ILE	2.2
2	B	228	PHE	2.2
1	A	1963	LEU	2.2
1	A	2026	LEU	2.1
1	A	2058	LEU	2.1
2	B	95	ASN	2.1
1	A	1852	VAL	2.1
2	B	276	PRO	2.1
2	B	289	TRP	2.1
2	B	23	ASN	2.1
1	A	1954	ILE	2.1
1	A	1959	THR	2.0
1	A	1839	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

3	VF9	A	2101	15/15	0.82	0.26	20,20,20,20	0
---	-----	---	------	-------	------	------	-------------	---

6.5 Other polymers [i](#)

There are no such residues in this entry.