



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 6, 2023 – 04:16 PM EDT

PDB ID : 4FUK  
Title : Aminopeptidase from Trypanosoma brucei  
Authors : El Bakkouri, M.; Tempel, W.; Osman, K.T.; Loppnau, P.; Graslund, S.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Hui, R.; Lin, Y.H.; Structural Genomics Consortium (SGC)  
Deposited on : 2012-06-28  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	B	337	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UNX	B	405	-	-	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Methionine aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2497	1568	434	479	16	0	4	0
1	B	328	2597	1638	457	486	16	0	7	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	GLY	-	expression tag	UNP Q4FKC0
A	86	LEU	PRO	conflict	UNP Q4FKC0
A	93	ARG	CYS	conflict	UNP Q4FKC0
A	395	VAL	-	expression tag	UNP Q4FKC0
B	59	GLY	-	expression tag	UNP Q4FKC0
B	86	LEU	PRO	conflict	UNP Q4FKC0
B	93	ARG	CYS	conflict	UNP Q4FKC0
B	395	VAL	-	expression tag	UNP Q4FKC0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	X	0	0
			2	2		
3	B	15	Total	X	0	0
			15	15		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	57	Total O 57 57	0	0
5	B	303	Total O 303 303	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Methionine aminopeptidase



- Molecule 1: Methionine aminopeptidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.38Å 75.55Å 181.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.19 – 1.75 47.14 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.19-1.75) 100.0 (47.14-1.75)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.36 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.7.0027	Depositor
R, $R_{free}$	0.194 , 0.229 0.197 , 0.233	Depositor DCC
$R_{free}$ test set	3476 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.2	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5493	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, UNX, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/2566	0.59	0/3497
1	B	0.67	0/2679	0.77	1/3639 (0.0%)
All	All	0.57	0/5245	0.68	1/7136 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	78	ARG	NE-CZ-NH2	-5.26	117.67	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2497	0	2326	28	0
1	B	2597	0	2554	31	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	2	0	0	0	0
3	B	15	0	0	0	0
4	B	18	0	24	2	0
5	A	57	0	0	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	303	0	0	6	0
All	All	5493	0	4904	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:133:VAL:HG23	1:B:170[B]:MET:HE3	1.38	1.05
1:B:133:VAL:HG23	1:B:170[B]:MET:CE	1.96	0.95
1:A:281:THR:O	1:B:63[B]:THR:HG22	1.84	0.77
1:B:78:ARG:HD2	5:B:766:HOH:O	1.89	0.72
1:B:87[A]:ARG:HG2	1:B:91:ALA:CB	2.23	0.69
1:B:170[B]:MET:HE1	1:B:214:SER:CB	2.27	0.63
1:A:76[B]:THR:HG23	1:A:77:PRO:HD2	1.81	0.61
1:B:238:ARG:HA	1:B:384:PRO:HG3	1.83	0.59
1:B:343[A]:ARG:NH1	5:B:513:HOH:O	2.35	0.59
1:B:70:LEU:HD21	4:B:413:GOL:H11	1.85	0.58
1:B:319:PRO:HG2	5:B:756:HOH:O	2.04	0.57
1:B:170[B]:MET:HE1	1:B:214:SER:HB2	1.87	0.56
1:B:170[B]:MET:HE1	1:B:214:SER:HB3	1.87	0.55
1:B:133:VAL:HG23	1:B:170[B]:MET:HE1	1.85	0.54
1:B:170[A]:MET:HG2	1:B:216:LEU:HB2	1.91	0.53
1:B:343[B]:ARG:HH11	1:B:343[B]:ARG:HG3	1.72	0.53
1:B:343[A]:ARG:NH2	5:B:628:HOH:O	2.42	0.52
1:B:170[B]:MET:HE3	1:B:216:LEU:HB2	1.90	0.52
1:A:110:LYS:HG2	5:A:544:HOH:O	2.10	0.51
1:B:343[B]:ARG:NH2	5:B:628:HOH:O	2.44	0.50
4:B:413:GOL:H12	5:B:562:HOH:O	2.12	0.50
1:B:87[A]:ARG:HG2	1:B:91:ALA:HB3	1.94	0.50
1:A:211:ASP:HA	1:A:224:ASN:HB3	1.93	0.50
1:A:81:VAL:HG13	1:A:87:ARG:HH21	1.78	0.48
1:A:273:GLN:NE2	1:A:386:ARG:HD3	2.28	0.48
1:B:133:VAL:CG2	1:B:170[B]:MET:CE	2.83	0.48
1:B:214:SER:O	1:B:220:HIS:HA	2.13	0.48
1:A:194:HIS:HE2	1:B:60:ALA:N	2.12	0.47
1:A:285:HIS:CE1	1:A:292:HIS:HD2	2.32	0.47
1:A:325:THR:HG22	1:A:341:ASP:CG	2.34	0.47
1:A:143:ILE:HD11	1:A:168:ARG:HH22	1.80	0.47
1:B:343[B]:ARG:HG3	1:B:343[B]:ARG:NH1	2.29	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:SER:HB3	1:A:223:LEU:HD21	1.98	0.46
1:A:211:ASP:HA	1:A:224:ASN:CB	2.47	0.45
1:A:317:ILE:HG22	1:A:319:PRO:HD3	1.98	0.45
1:A:239:LEU:HD13	1:A:323:LEU:HG	1.99	0.45
1:A:98:SER:O	1:A:102:GLU:HG2	2.17	0.44
1:A:365:TRP:CE2	1:A:370:PRO:HA	2.52	0.44
1:A:62:LYS:HE3	5:A:505:HOH:O	2.17	0.44
1:A:236:SER:OG	1:A:344:ARG:NH1	2.51	0.44
1:A:137:SER:OG	1:A:214:SER:HB2	2.18	0.44
1:B:170[B]:MET:CE	1:B:216:LEU:HB2	2.48	0.43
1:A:228:PHE:CD1	1:A:232:PRO:HB3	2.53	0.43
1:B:78:ARG:HB2	1:B:175:LEU:HD11	2.01	0.43
1:A:78:ARG:NH2	1:A:176:ASN:OD1	2.51	0.43
1:A:60:ALA:N	1:B:194:HIS:HE2	2.17	0.42
1:A:335:TRP:CZ2	1:B:334:LYS:HE3	2.54	0.42
1:A:138:ARG:HA	1:A:223:LEU:HD11	2.02	0.42
1:A:325:THR:HG22	1:A:341:ASP:OD2	2.20	0.41
1:A:88:PRO:HD3	1:A:215:TYR:CD2	2.56	0.41
1:A:323:LEU:HD12	1:A:344:ARG:HB2	2.02	0.41
1:B:367:ASP:OD2	1:B:374:LYS:NZ	2.48	0.41
1:B:365:TRP:CE2	1:B:370:PRO:HA	2.56	0.41
1:A:76[B]:THR:CG2	1:A:77:PRO:HD2	2.50	0.41
1:B:274:TYR:CZ	1:B:386:ARG:HG2	2.56	0.40
1:B:170[B]:MET:CE	1:B:214:SER:HB3	2.50	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	325/337 (96%)	312 (96%)	12 (4%)	1 (0%)	41 22

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	333/337 (99%)	326 (98%)	7 (2%)	0	100	100
All	All	658/674 (98%)	638 (97%)	19 (3%)	1 (0%)	47	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	386	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/287 (89%)	255 (100%)	0	100	100
1	B	283/287 (99%)	279 (99%)	4 (1%)	67	52
All	All	538/574 (94%)	534 (99%)	4 (1%)	86	75

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	78	ARG
1	B	87[A]	ARG
1	B	87[B]	ARG
1	B	387	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 4 are monoatomic and 17 are unknown - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	B	408	-	5,5,5	0.72	0	5,5,5	0.64	0
4	GOL	B	413	-	5,5,5	0.51	0	5,5,5	1.86	1 (20%)
4	GOL	B	409	-	5,5,5	0.61	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	408	-	-	0/4/4/4	-
4	GOL	B	413	-	-	3/4/4/4	-
4	GOL	B	409	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	413	GOL	C3-C2-C1	-3.74	97.18	111.70

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	413	GOL	C1-C2-C3-O3
4	B	413	GOL	O2-C2-C3-O3
4	B	413	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	413	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	329/337 (97%)	0.95	58 (17%) <b>1</b> <b>2</b>	22, 46, 68, 79	0
1	B	328/337 (97%)	-0.09	0 <b>100</b> <b>100</b>	13, 20, 35, 56	0
All	All	657/674 (97%)	0.43	58 (8%) <b>10</b> <b>13</b>	13, 31, 64, 79	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	379	TRP	5.0
1	A	366	VAL	4.7
1	A	382	MET	4.5
1	A	230	GLY	4.2
1	A	369	VAL	4.1
1	A	376	LEU	4.0
1	A	259	LEU	3.9
1	A	140	VAL	3.9
1	A	392	ALA	3.8
1	A	306	LEU	3.7
1	A	121	SER	3.7
1	A	308	MET	3.4
1	A	125	ALA	3.4
1	A	169	ASN	3.4
1	A	151	GLY	3.3
1	A	105	ARG	3.3
1	A	118	HIS	3.3
1	A	254	VAL	3.2
1	A	251	ILE	3.2
1	A	357	GLY	3.1
1	A	367	ASP	3.1
1	A	111	VAL	3.0
1	A	119	ASP	2.9
1	A	241	HIS	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	114	ILE	2.8
1	A	117	LEU	2.7
1	A	311	PRO	2.7
1	A	244	TYR	2.6
1	A	83	SER	2.5
1	A	144	ALA	2.5
1	A	147	ALA	2.5
1	A	312	GLY	2.5
1	A	362	PHE	2.5
1	A	356	ASN	2.5
1	A	371	THR	2.5
1	A	237	VAL	2.5
1	A	104	ASP	2.4
1	A	383	LEU	2.3
1	A	112	TYR	2.3
1	A	247	LEU	2.3
1	A	115	GLN	2.3
1	A	239	LEU	2.3
1	A	258	ALA	2.3
1	A	120	ASP	2.2
1	A	375	GLN	2.2
1	A	234	ASP	2.2
1	A	166	VAL	2.2
1	A	353	VAL	2.2
1	A	267	ILE	2.2
1	A	365	TRP	2.2
1	A	252	GLY	2.1
1	A	358	GLY	2.1
1	A	274	TYR	2.1
1	A	82	PRO	2.1
1	A	359	VAL	2.1
1	A	307	GLY	2.1
1	A	81	VAL	2.0
1	A	275	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	UNX	B	405	1/1	0.59	0.48	50,50,50,50	0
3	UNX	A	403	1/1	0.64	0.30	32,32,32,32	0
3	UNX	B	420	1/1	0.65	0.37	34,34,34,34	0
3	UNX	B	404	1/1	0.69	0.26	25,25,25,25	0
3	UNX	B	419	1/1	0.73	0.33	34,34,34,34	0
3	UNX	B	415	1/1	0.80	0.32	37,37,37,37	0
3	UNX	B	407	1/1	0.80	0.20	33,33,33,33	0
3	UNX	B	414	1/1	0.80	0.26	25,25,25,25	0
3	UNX	B	406	1/1	0.83	0.22	29,29,29,29	0
3	UNX	B	416	1/1	0.84	0.14	36,36,36,36	0
3	UNX	B	412	1/1	0.85	0.16	35,35,35,35	0
3	UNX	B	410	1/1	0.86	0.31	38,38,38,38	0
4	GOL	B	413	6/6	0.87	0.20	18,25,28,39	0
3	UNX	B	411	1/1	0.88	0.27	43,43,43,43	0
3	UNX	A	404	1/1	0.88	0.48	41,41,41,41	0
4	GOL	B	409	6/6	0.90	0.14	30,36,36,51	0
3	UNX	B	418	1/1	0.90	0.31	26,26,26,26	0
4	GOL	B	408	6/6	0.91	0.11	24,28,29,30	0
2	ZN	A	401	1/1	0.98	0.09	30,30,30,30	0
3	UNX	B	417	1/1	0.98	0.20	28,28,28,28	0
2	ZN	A	402	1/1	0.98	0.06	32,32,32,32	0
3	UNX	B	403	1/1	0.98	0.15	28,28,28,28	0
2	ZN	B	401	1/1	0.99	0.08	17,17,17,17	0
2	ZN	B	402	1/1	1.00	0.10	17,17,17,17	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.