



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2023 – 08:04 AM EDT

PDB ID : 8FUQ
Title : Crystal structure of rabies virus (strain CVS-11) P3 dimerization domain
Authors : Donnelly, C.M.; Cross, E.M.; Forwood, J.K.
Deposited on : 2023-01-18
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	2% 38% 5% 57%
1	B	119	% 32% 11% 57%
1	C	119	% 39% .. 58%
1	D	119	3% 33% 6% • 61%
1	E	119	3% 34% 5% • 59%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	119	 35% . 61%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4970 atoms, of which 2370 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	51	Total 821	C 272	H 398	N 70	O 80	S 1	0	1	0
1	B	51	Total 831	C 271	H 410	N 72	O 77	S 1	0	1	0
1	C	50	Total 794	C 261	H 388	N 66	O 78	S 1	0	0	0
1	D	47	Total 778	C 252	H 389	N 65	O 71	S 1	0	1	0
1	E	49	Total 795	C 259	H 394	N 67	O 74	S 1	0	0	0
1	F	47	Total 782	C 256	H 391	N 62	O 72	S 1	0	2	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	SER	-	expression tag	UNP P15198
B	53	SER	-	expression tag	UNP P15198
C	53	SER	-	expression tag	UNP P15198
D	53	SER	-	expression tag	UNP P15198
E	53	SER	-	expression tag	UNP P15198
F	53	SER	-	expression tag	UNP P15198

- Molecule 2 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	30	Total 30	O 30	0	0
2	B	46	Total 46	O 46	0	0
2	C	30	Total 30	O 30	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	29	Total O 29 29	0	0
2	E	12	Total O 12 12	0	0
2	F	22	Total O 22 22	0	0

- Molecule 1: Phosphoprotein



- Molecule 1: Phosphoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	67.05Å 44.10Å 111.79Å 90.00° 100.44° 90.00°	Depositor
Resolution (Å)	52.51 – 1.80 52.51 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (52.51-1.80) 99.7 (52.51-1.80)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.80Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.208 , 0.237 0.207 , 0.238	Depositor DCC
R_{free} test set	1499 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtrriage
Anisotropy	0.109	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4970	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	0/436	0.73	0/594
1	B	0.67	0/434	0.67	0/592
1	C	0.75	0/414	0.74	0/562
1	D	0.69	0/400	0.70	0/542
1	E	0.81	1/410 (0.2%)	0.83	0/558
1	F	0.69	0/402	0.65	0/547
All	All	0.73	1/2496 (0.0%)	0.72	0/3395

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	1
1	D	0	2
1	F	0	1
All	All	0	7

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	124	GLU	CB-CG	5.92	1.63	1.52

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	109	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	113	ARG	Sidechain
1	B	113	ARG	Sidechain
1	C	109	ARG	Sidechain
1	D	109	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	423	398	416	6	0
1	B	421	410	411	10	0
1	C	406	388	393	1	0
1	D	389	389	389	4	0
1	E	401	394	394	9	0
1	F	391	391	384	6	0
2	A	30	0	0	0	0
2	B	46	0	0	0	1
2	C	30	0	0	0	0
2	D	29	0	0	0	1
2	E	12	0	0	0	0
2	F	22	0	0	0	0
All	All	2600	2370	2387	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 25 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:115:LEU:HD23	1:F:105:VAL:CG1	2.22	0.69
1:A:136:PRO:HG3	1:B:133:PHE:CD2	2.29	0.67
1:E:115:LEU:HD23	1:F:105:VAL:HG13	1.76	0.66
1:B:122:VAL:O	1:B:126:VAL:HG23	2.06	0.55
1:B:98:ASP:HB3	1:D:102:VAL:HG13	1.92	0.52

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:216:HOH:O	2:D:218:HOH:O[4_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	50/119 (42%)	50 (100%)	0	0	100	100
1	B	50/119 (42%)	50 (100%)	0	0	100	100
1	C	48/119 (40%)	47 (98%)	1 (2%)	0	100	100
1	D	46/119 (39%)	46 (100%)	0	0	100	100
1	E	47/119 (40%)	47 (100%)	0	0	100	100
1	F	47/119 (40%)	47 (100%)	0	0	100	100
All	All	288/714 (40%)	287 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	50/110 (46%)	50 (100%)	0	100	100
1	B	48/110 (44%)	46 (96%)	2 (4%)	30	15
1	C	46/110 (42%)	43 (94%)	3 (6%)	17	6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	45/110 (41%)	44 (98%)	1 (2%)	52	39
1	E	46/110 (42%)	45 (98%)	1 (2%)	52	39
1	F	44/110 (40%)	44 (100%)	0	100	100
All	All	279/660 (42%)	272 (98%)	7 (2%)	46	34

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	116	LYS
1	C	135	ASN
1	E	115	LEU
1	D	113	ARG
1	C	109	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	120	GLN
1	B	90	ASN
1	C	107	GLN
1	C	132	ASN
1	D	107	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	51/119 (42%)	0.16	2 (3%) 39 33	17, 28, 46, 76	0
1	B	51/119 (42%)	0.27	1 (1%) 65 61	14, 24, 49, 60	0
1	C	50/119 (42%)	-0.00	1 (2%) 65 61	16, 24, 48, 50	0
1	D	47/119 (39%)	0.07	3 (6%) 19 15	14, 24, 46, 60	0
1	E	49/119 (41%)	0.46	4 (8%) 11 9	23, 36, 53, 92	0
1	F	47/119 (39%)	0.18	0 100 100	19, 34, 54, 57	0
All	All	295/714 (41%)	0.19	11 (3%) 41 36	14, 29, 53, 92	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	136	PRO	6.0
1	E	134	PRO	4.9
1	D	135	ASN	4.8
1	E	91	LEU	4.5
1	C	135	ASN	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.