



Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:45 pm BST

PDB ID : 3FVZ
Title : Structure of Peptidyl-alpha-hydroxyglycine alpha-Amidating Lyase (PAL)
Authors : Chufan, E.E.; De, M.; Eipper, B.A.; Mains, R.E.; Amzel, L.M.
Deposited on : 2009-01-16
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	329	 78% 18%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2891 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidyl-glycine alpha-amidating monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2724	1728	477	511	8	0	13	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	492	HIS	-	EXPRESSION TAG	UNP P14925
A	493	HIS	-	EXPRESSION TAG	UNP P14925
A	494	HIS	-	EXPRESSION TAG	UNP P14925
A	495	HIS	-	EXPRESSION TAG	UNP P14925
A	496	HIS	-	EXPRESSION TAG	UNP P14925
A	497	HIS	-	EXPRESSION TAG	UNP P14925

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	148	Total 148	O 148	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptidyl-glycine alpha-amidating monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.18Å 75.06Å 97.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.22 – 2.35 39.22 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.6 (39.22-2.35) 99.6 (39.22-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.82 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.263 0.267 , 0.306	Depositor DCC
R_{free} test set	847 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	52.8	Xtrriage
Anisotropy	0.439	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2891	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, ACT, CA, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.41	0/2807	0.59	1/3814 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	811	LEU	CA-CB-CG	5.21	127.29	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	653	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2724	0	2556	75	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	4	0	3	0	0
6	A	12	0	16	2	0
7	A	148	0	0	1	0
All	All	2891	0	2575	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:502[B]:GLU:OE1	1:A:818[B]:ARG:HD2	1.37	1.25
1:A:819[B]:SER:O	1:A:820[B]:VAL:HG22	1.43	1.12
1:A:500[B]:HIS:HB2	1:A:819[B]:SER:HB3	1.29	1.08
1:A:500[B]:HIS:HB2	1:A:819[B]:SER:CB	1.88	1.04
1:A:496[B]:HIS:NE2	1:A:817[B]:HIS:CD2	2.32	0.96
1:A:819[B]:SER:O	1:A:820[B]:VAL:CG2	2.17	0.93
1:A:564:VAL:HG13	1:A:574:GLN:HG2	1.53	0.90
1:A:819[B]:SER:O	1:A:820[B]:VAL:HG13	1.78	0.83
1:A:500[B]:HIS:CG	1:A:819[B]:SER:HA	2.15	0.82
1:A:500[B]:HIS:CB	1:A:819[B]:SER:HB3	2.13	0.78
1:A:500[B]:HIS:CD2	1:A:819[B]:SER:HA	2.20	0.77
1:A:502[B]:GLU:OE1	1:A:818[B]:ARG:CD	2.28	0.76
1:A:792[A]:GLU:OE2	7:A:148:HOH:O	2.04	0.76
1:A:819[B]:SER:C	1:A:820[B]:VAL:HG22	2.06	0.75
1:A:500[B]:HIS:CB	1:A:819[B]:SER:CB	2.66	0.73
1:A:638:ASP:CG	6:A:824:GOL:H12	2.09	0.72
1:A:558:GLU:O	1:A:578:LYS:HG3	1.93	0.69
1:A:819[B]:SER:O	1:A:820[B]:VAL:CG1	2.40	0.69
1:A:784:MET:CE	1:A:801:ALA:HB3	2.22	0.69
1:A:496[B]:HIS:CE1	1:A:817[B]:HIS:CD2	2.81	0.68
1:A:750:LYS:HA	1:A:759:VAL:HG22	1.76	0.67
1:A:527:ASN:HD22	1:A:567:PRO:HD3	1.60	0.67
1:A:784:MET:HE2	1:A:801:ALA:HB3	1.76	0.67
1:A:819[B]:SER:O	1:A:820[B]:VAL:CB	2.44	0.65
1:A:564:VAL:CG1	1:A:574:GLN:HG2	2.24	0.65
1:A:561:THR:HG22	1:A:577:GLY:HA3	1.78	0.64
1:A:564:VAL:HG13	1:A:574:GLN:CG	2.25	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:496[B]:HIS:CG	1:A:497[B]:HIS:H	2.15	0.62
1:A:497[B]:HIS:CE1	1:A:772:ILE:HD12	2.35	0.61
1:A:500[B]:HIS:HB2	1:A:819[B]:SER:OG	2.00	0.59
1:A:502[B]:GLU:OE1	1:A:818[B]:ARG:NH1	2.35	0.59
1:A:812:THR:HG21	1:A:814:LYS:HE3	1.85	0.58
1:A:496[B]:HIS:NE2	1:A:817[B]:HIS:CG	2.72	0.57
1:A:496[B]:HIS:CG	1:A:497[B]:HIS:N	2.73	0.56
1:A:497[B]:HIS:CE1	1:A:499:PHE:HD1	2.25	0.55
1:A:500[B]:HIS:CD2	1:A:502[B]:GLU:HG3	2.42	0.55
1:A:497[B]:HIS:CD2	1:A:772:ILE:O	2.60	0.54
1:A:659:ILE:HD11	1:A:703:VAL:HG21	1.89	0.54
1:A:638:ASP:OD1	6:A:824:GOL:H12	2.08	0.53
1:A:564:VAL:HG22	1:A:573:LEU:HB2	1.91	0.52
1:A:642:GLU:OE2	1:A:716:THR:HG21	2.09	0.52
1:A:497[B]:HIS:CE1	1:A:772:ILE:HB	2.44	0.52
1:A:784:MET:HE1	1:A:801:ALA:HB3	1.92	0.52
1:A:817[B]:HIS:HB3	1:A:820[B]:VAL:H	1.75	0.52
1:A:497[B]:HIS:O	1:A:817[B]:HIS:CD2	2.64	0.51
1:A:671:GLN:HG2	1:A:672:TRP:N	2.24	0.51
1:A:497[B]:HIS:HE1	1:A:499:PHE:HD1	1.60	0.49
1:A:561:THR:HG21	1:A:581:PHE:CD1	2.48	0.48
1:A:692:LEU:HD22	1:A:701:LEU:HD11	1.95	0.48
1:A:733:ASN:HD22	1:A:749:GLY:HA3	1.79	0.48
1:A:723:ARG:HG2	1:A:724:GLU:N	2.29	0.47
1:A:676:SER:HA	1:A:685:GLN:HB3	1.95	0.47
1:A:500[B]:HIS:CB	1:A:819[B]:SER:OG	2.61	0.46
1:A:500[B]:HIS:CG	1:A:819[B]:SER:CA	2.93	0.46
1:A:496[A]:HIS:HB3	1:A:497[A]:HIS:H	1.27	0.46
1:A:792[A]:GLU:CD	1:A:792[A]:GLU:H	2.19	0.46
1:A:718:THR:O	1:A:719:LYS:HB2	2.15	0.45
1:A:639:VAL:HG22	1:A:650:VAL:HG22	1.98	0.45
1:A:598:THR:HG22	1:A:605:VAL:HG22	1.97	0.45
1:A:496[B]:HIS:CD2	1:A:497[B]:HIS:H	2.35	0.44
1:A:653:GLY:O	1:A:688[A]:VAL:HA	2.19	0.43
1:A:500[B]:HIS:CG	1:A:819[B]:SER:HB3	2.53	0.43
1:A:691:SER:HB3	1:A:704:ALA:HB3	1.99	0.43
1:A:507:TRP:HA	1:A:508:PRO:HD3	1.86	0.43
1:A:819[B]:SER:C	1:A:820[B]:VAL:CG2	2.76	0.42
1:A:531:PHE:CE2	1:A:584:PRO:HB3	2.55	0.41
1:A:561:THR:HG21	1:A:581:PHE:HD1	1.85	0.41
1:A:600:VAL:HG13	1:A:635:GLN:HG2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:706:ARG:O	1:A:733:ASN:HA	2.20	0.41
1:A:500[A]:HIS:CE1	1:A:814:LYS:HB2	2.55	0.41
1:A:531:PHE:CE1	1:A:561:THR:HG23	2.56	0.41
1:A:497[B]:HIS:NE2	1:A:499:PHE:HB2	2.37	0.40
1:A:494:HIS:HB2	1:A:817[B]:HIS:CE1	2.56	0.40
1:A:500[B]:HIS:CG	1:A:819[B]:SER:CB	3.05	0.40
1:A:687[A]:SER:O	1:A:688[A]:VAL:HB	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/329 (103%)	320 (94%)	17 (5%)	2 (1%)	25 27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	495	HIS
1	A	653	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/289 (104%)	279 (92%)	23 (8%)	13 13

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	496[A]	HIS
1	A	496[B]	HIS
1	A	497[A]	HIS
1	A	497[B]	HIS
1	A	500[A]	HIS
1	A	500[B]	HIS
1	A	548	VAL
1	A	554	LEU
1	A	561	THR
1	A	654	TYR
1	A	655	CYS
1	A	658	ARG
1	A	663	SER
1	A	671	GLN
1	A	679	SER
1	A	706	ARG
1	A	707	GLU
1	A	716	THR
1	A	718	THR
1	A	723	ARG
1	A	744	LEU
1	A	759	VAL
1	A	811	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	495	HIS
1	A	527	ASN
1	A	579	ASN
1	A	708	ASN
1	A	733	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	2	-	5,5,5	0.46	0	5,5,5	0.23	0
5	ACT	A	1	2	1,3,3	1.40	0	0,3,3	0.00	-
6	GOL	A	824	4	5,5,5	0.48	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	2	-	-	4/4/4/4	-
6	GOL	A	824	4	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	2	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	2	GOL	C1-C2-C3-O3
6	A	824	GOL	O1-C1-C2-C3
6	A	824	GOL	O1-C1-C2-O2
6	A	2	GOL	O1-C1-C2-O2
6	A	2	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	824	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.