



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 06:57 pm BST

PDB ID : 4G4M  
Title : Crystal structure of the de novo designed fluorinated peptide alpha4F3(6-13)  
Authors : Buer, B.C.; Meagher, J.L.; Stuckey, J.A.; Marsh, E.N.G.  
Deposited on : 2012-07-16  
Resolution : 1.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	27	
1	B	27	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	102	-	X	-	-
3	EDO	A	103	-	X	-	-
3	EDO	B	102	-	X	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha4F3(6-13).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	F	N	O			
1	A	25	Total 232	C 132	F 18	N 42	O 40	0	0	0
1	B	25	Total 236	C 135	F 18	N 43	O 40	0	0	0

- Molecule 2 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 7	C 4	O 3	0	0
2	B	1	Total 7	C 4	O 3	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	12	Total O 12 12	0	0
4	B	16	Total O 16 16	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha4F3(6-13)

Chain A:  70% 22% 7%



- Molecule 1: alpha4F3(6-13)

Chain B:  81% 11% 7%



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.58Å 49.58Å 41.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.48 9.96 – 1.48	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.48) 99.9 (9.96-1.48)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.57 (at 1.48Å)	Xtrriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 1.6.0	Depositor
R, $R_{free}$	0.252 , 0.318 0.192 , 0.211	Depositor DCC
$R_{free}$ test set	394 reflections (4.66%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.9	Xtrriage
Anisotropy	0.285	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 62.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.22$	Xtrriage
Estimated twinning fraction	0.468 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	522	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, 6FL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/187	0.72	0/240
1	B	0.46	0/191	0.70	0/244
All	All	0.49	0/378	0.71	0/484

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	232	0	203	4	0
1	B	236	0	214	1	0
2	A	7	0	9	0	0
2	B	7	0	9	0	0
3	A	8	0	10	2	0
3	B	4	0	5	0	0
4	A	12	0	0	1	0
4	B	16	0	0	0	0
All	All	522	0	450	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:ALA:H	3:A:103:EDO:H21	1.67	0.59
1:A:6:6FL:O	1:A:10:6FL:HG	2.07	0.54
1:A:3:ALA:HB2	3:A:103:EDO:H12	1.95	0.49
1:B:6:6FL:O	1:B:10:6FL:HG	2.18	0.44
1:A:18:ARG:NE	4:A:205:HOH:O	2.53	0.42

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	20/27 (74%)	19 (95%)	1 (5%)	0	100	100
1	B	20/27 (74%)	19 (95%)	1 (5%)	0	100	100
All	All	40/54 (74%)	38 (95%)	2 (5%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	19/21 (90%)	17 (90%)	2 (10%)	7	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	20/21 (95%)	19 (95%)	1 (5%)	24	3
All	All	39/42 (93%)	36 (92%)	3 (8%)	13	1

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	12	ASP
1	B	2	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	6FL	A	6	1	10,13,14	0.83	0	15,20,22	1.25	1 (6%)
1	6FL	A	10	1	10,13,14	0.94	0	15,20,22	1.40	3 (20%)
1	6FL	B	10	1	10,13,14	0.84	1 (10%)	15,20,22	1.67	3 (20%)
1	6FL	B	6	1	10,13,14	0.61	0	15,20,22	1.50	3 (20%)
1	6FL	B	13	1	10,13,14	0.73	0	15,20,22	1.58	3 (20%)
1	6FL	A	13	1	10,13,14	0.88	1 (10%)	15,20,22	1.27	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	6FL	A	6	1	-	4/17/18/20	-
1	6FL	A	10	1	-	1/17/18/20	-
1	6FL	B	10	1	-	1/17/18/20	-
1	6FL	B	6	1	-	7/17/18/20	-
1	6FL	B	13	1	-	1/17/18/20	-
1	6FL	A	13	1	-	1/17/18/20	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	6FL	CD2-CG	2.19	1.55	1.50
1	B	10	6FL	CD2-CG	2.01	1.55	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	6FL	CB-CG-CD2	3.22	115.53	110.95
1	B	10	6FL	FAG-CD2-CG	-3.14	106.34	112.88
1	B	10	6FL	FAD-CD1-CG	-2.98	106.68	112.88
1	A	13	6FL	CB-CG-CD2	2.89	115.06	110.95
1	B	13	6FL	CB-CG-CD1	-2.84	106.93	110.95
1	B	10	6FL	FAH-CD2-FAF	2.81	113.14	106.87
1	B	6	6FL	FAC-CD1-CG	-2.69	107.28	112.88
1	B	13	6FL	FAF-CD2-CG	-2.54	107.59	112.88
1	B	13	6FL	FAC-CD1-CG	-2.43	107.82	112.88
1	B	6	6FL	FAH-CD2-CG	-2.21	108.27	112.88
1	B	6	6FL	FAF-CD2-CG	-2.18	108.35	112.88
1	A	10	6FL	FAH-CD2-CG	-2.16	108.39	112.88
1	A	10	6FL	FAD-CD1-CG	-2.03	108.66	112.88
1	A	6	6FL	CB-CG-CD2	2.01	113.81	110.95

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	10	6FL	CA-CB-CG-CD2
1	B	6	6FL	C-CA-CB-CG
1	B	6	6FL	CA-CB-CG-CD1
1	A	6	6FL	FAC-CD1-CG-CB

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	6	6FL	FAF-CD2-CG-CB
1	B	6	6FL	FAG-CD2-CG-CB
1	B	6	6FL	FAH-CD2-CG-CB
1	A	6	6FL	CA-CB-CG-CD2
1	A	10	6FL	CA-CB-CG-CD2
1	B	6	6FL	CA-CB-CG-CD2
1	B	13	6FL	CA-CB-CG-CD2
1	A	13	6FL	CA-CB-CG-CD2
1	A	6	6FL	FAD-CD1-CG-CB
1	A	6	6FL	FAE-CD1-CG-CB
1	B	6	6FL	N-CA-CB-CG

There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	6	6FL	1	0
1	A	10	6FL	1	0
1	B	10	6FL	1	0
1	B	6	6FL	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PG4	B	101	-	6,6,12	1.98	2 (33%)	5,5,11	2.37	2 (40%)
3	EDO	A	103	-	3,3,3	1.98	2 (66%)	2,2,2	2.17	1 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	A	102	-	3,3,3	2.26	2 (66%)	2,2,2	2.61	2 (100%)
2	PG4	A	101	-	6,6,12	1.77	1 (16%)	5,5,11	2.45	2 (40%)
3	EDO	B	102	-	3,3,3	2.06	2 (66%)	2,2,2	2.30	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PG4	B	101	-	-	1/4/4/10	-
3	EDO	A	103	-	-	1/1/1/1	-
3	EDO	A	102	-	-	1/1/1/1	-
2	PG4	A	101	-	-	1/4/4/10	-
3	EDO	B	102	-	-	1/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	101	PG4	O5-C8	3.96	1.62	1.42
2	A	101	PG4	O5-C8	3.63	1.60	1.42
3	A	102	EDO	O1-C1	2.73	1.56	1.42
3	B	102	EDO	O1-C1	2.45	1.54	1.42
3	A	103	EDO	O1-C1	2.41	1.54	1.42
3	A	102	EDO	C2-C1	2.40	1.64	1.48
2	B	101	PG4	C6-C5	2.29	1.61	1.49
3	B	102	EDO	C2-C1	2.19	1.63	1.48
3	A	103	EDO	C2-C1	2.14	1.63	1.48

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	101	PG4	C7-O4-C6	-4.43	94.09	113.29
2	A	101	PG4	C7-O4-C6	-4.31	94.59	113.29
2	A	101	PG4	O4-C7-C8	3.10	123.71	110.07
3	A	102	EDO	O2-C2-C1	3.07	134.03	111.91
3	A	103	EDO	O2-C2-C1	2.76	131.77	111.91
3	B	102	EDO	O2-C2-C1	2.67	131.13	111.91
2	B	101	PG4	O4-C7-C8	2.65	121.69	110.07
3	A	102	EDO	O1-C1-C2	2.05	126.65	111.91

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	101	PG4	O3-C5-C6-O4
3	A	103	EDO	O1-C1-C2-O2
3	A	102	EDO	O1-C1-C2-O2
2	A	101	PG4	C5-C6-O4-C7
3	B	102	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	103	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.