



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 3, 2023 – 10:22 pm GMT

PDB ID : 1GK9
Title : Crystal structures of penicillin acylase enzyme-substrate complexes: Structural insights into the catalytic mechanism
Authors : McVey, C.E.; Walsh, M.A.; Dodson, G.G.; Wilson, K.S.; Brannigan, J.A.
Deposited on : 2001-08-10
Resolution : 1.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PENICILLIN G ACYLASE ALPHA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	208	1750	1114	293	334	9	5	10	0

- Molecule 2 is a protein called PENICILLIN G ACYLASE BETA SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	557	4751	3016	825	899	11	8	40	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	4	2	2	0	0
3	A	1	8	4	4	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 8	C 4	O 4	0	1
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0
3	B	1	Total 4	C 2	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	1
			8	4	4		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	254	Total	O	0	5
			254	254		
5	B	669	Total	O	0	10
			669	669		

MolProbity failed to run properly - this section is therefore empty.

3 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.30Å 131.60Å 63.90Å 90.00° 105.90° 90.00°	Depositor
Resolution (Å)	20.00 – 1.30 19.99 – 1.30	Depositor EDS
% Data completeness (in resolution range)	95.5 (20.00-1.30) 95.3 (19.99-1.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 1.30Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.148 , 0.169 0.154 , 0.174	Depositor DCC
R_{free} test set	5756 reflections (3.03%)	wwPDB-VP
Wilson B-factor (Å ²)	10.8	Xtrriage
Anisotropy	0.118	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7533	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 1 is monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	B	1578[A]	-	3,3,3	0.67	0	2,2,2	0.54	0
3	EDO	B	1572	-	3,3,3	0.44	0	2,2,2	0.47	0
3	EDO	B	1560	-	3,3,3	0.80	0	2,2,2	0.22	0
3	EDO	A	1211[B]	-	3,3,3	0.49	0	2,2,2	0.13	0
3	EDO	B	1569	-	3,3,3	0.53	0	2,2,2	0.21	0
3	EDO	B	1561	-	3,3,3	0.99	0	2,2,2	0.61	0
3	EDO	B	1562	-	3,3,3	0.59	0	2,2,2	0.74	0
3	EDO	A	1210	-	3,3,3	0.59	0	2,2,2	0.24	0
3	EDO	B	1577	-	3,3,3	0.56	0	2,2,2	0.53	0
3	EDO	B	1571	-	3,3,3	0.58	0	2,2,2	0.36	0
3	EDO	B	1564	-	3,3,3	0.42	0	2,2,2	1.04	0
3	EDO	A	1211[A]	-	3,3,3	0.52	0	2,2,2	0.25	0
3	EDO	B	1559	-	3,3,3	0.53	0	2,2,2	0.25	0
3	EDO	B	1574	-	3,3,3	0.45	0	2,2,2	0.70	0
3	EDO	B	1573	-	3,3,3	0.55	0	2,2,2	0.72	0
3	EDO	B	1563	-	3,3,3	1.04	0	2,2,2	1.04	0
3	EDO	B	1558	-	3,3,3	0.68	0	2,2,2	1.16	0
3	EDO	B	1566[B]	-	3,3,3	0.36	0	2,2,2	0.54	0
3	EDO	A	1212	-	3,3,3	0.57	0	2,2,2	0.57	0
3	EDO	B	1567	-	3,3,3	0.41	0	2,2,2	0.39	0
3	EDO	B	1568	-	3,3,3	0.57	0	2,2,2	0.02	0
3	EDO	B	1575	-	3,3,3	1.11	0	2,2,2	1.15	0
3	EDO	B	1578[B]	-	3,3,3	0.56	0	2,2,2	0.28	0
3	EDO	B	1570	-	3,3,3	0.54	0	2,2,2	0.60	0
3	EDO	B	1565	-	3,3,3	0.28	0	2,2,2	0.13	0
3	EDO	B	1566[A]	-	3,3,3	0.43	0	2,2,2	0.71	0
3	EDO	B	1576	-	3,3,3	0.44	0	2,2,2	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	1578[A]	-	-	0/1/1/1	-
3	EDO	B	1572	-	-	0/1/1/1	-
3	EDO	B	1560	-	-	0/1/1/1	-
3	EDO	A	1211[B]	-	-	1/1/1/1	-
3	EDO	B	1569	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	1561	-	-	0/1/1/1	-
3	EDO	B	1562	-	-	0/1/1/1	-
3	EDO	A	1210	-	-	1/1/1/1	-
3	EDO	B	1577	-	-	0/1/1/1	-
3	EDO	B	1571	-	-	0/1/1/1	-
3	EDO	B	1564	-	-	0/1/1/1	-
3	EDO	A	1211[A]	-	-	0/1/1/1	-
3	EDO	B	1559	-	-	0/1/1/1	-
3	EDO	B	1574	-	-	0/1/1/1	-
3	EDO	B	1573	-	-	0/1/1/1	-
3	EDO	B	1563	-	-	0/1/1/1	-
3	EDO	B	1558	-	-	0/1/1/1	-
3	EDO	B	1566[B]	-	-	1/1/1/1	-
3	EDO	A	1212	-	-	0/1/1/1	-
3	EDO	B	1567	-	-	1/1/1/1	-
3	EDO	B	1568	-	-	0/1/1/1	-
3	EDO	B	1575	-	-	0/1/1/1	-
3	EDO	B	1578[B]	-	-	1/1/1/1	-
3	EDO	B	1570	-	-	0/1/1/1	-
3	EDO	B	1565	-	-	0/1/1/1	-
3	EDO	B	1566[A]	-	-	0/1/1/1	-
3	EDO	B	1576	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1566[B]	EDO	O1-C1-C2-O2
3	A	1210	EDO	O1-C1-C2-O2
3	B	1578[B]	EDO	O1-C1-C2-O2
3	A	1211[B]	EDO	O1-C1-C2-O2
3	B	1567	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/260 (80%)	-0.22	2 (0%) 82 83	8, 12, 20, 68	2 (0%)
2	B	557/557 (100%)	-0.15	15 (2%) 54 52	6, 12, 26, 62	4 (0%)
All	All	765/817 (93%)	-0.17	17 (2%) 62 61	6, 12, 24, 68	6 (0%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLN	10.0
2	B	132	THR	7.3
2	B	131	GLN	4.7
2	B	133	THR	4.1
2	B	380[A]	GLN	3.9

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	EDO	A	1210	4/4	0.81	0.12	20,26,30,33	0
3	EDO	B	1567	4/4	0.84	0.11	26,29,31,35	0
3	EDO	A	1212	4/4	0.88	0.14	22,25,27,28	0
3	EDO	B	1577	4/4	0.92	0.25	18,22,24,25	0
3	EDO	B	1575	4/4	0.93	0.13	15,20,21,23	0
3	EDO	B	1565	4/4	0.93	0.07	18,19,24,25	0
3	EDO	B	1573	4/4	0.94	0.19	19,21,21,22	0
3	EDO	B	1558	4/4	0.95	0.09	12,14,15,18	0
3	EDO	B	1563	4/4	0.95	0.09	12,12,16,16	0
3	EDO	B	1570	4/4	0.95	0.12	17,23,23,23	0
3	EDO	A	1211[A]	4/4	0.96	0.14	14,17,19,22	4
3	EDO	B	1574	4/4	0.96	0.09	10,15,19,21	0
3	EDO	B	1569	4/4	0.96	0.08	13,15,17,18	0
3	EDO	B	1576	4/4	0.96	0.19	16,21,28,29	0
3	EDO	A	1211[B]	4/4	0.96	0.14	12,16,18,20	4
3	EDO	B	1578[A]	4/4	0.96	0.19	13,19,21,21	4
3	EDO	B	1578[B]	4/4	0.96	0.19	20,21,25,29	4
3	EDO	B	1572	4/4	0.97	0.16	10,22,24,29	0
3	EDO	B	1561	4/4	0.97	0.08	16,18,20,23	0
3	EDO	B	1564	4/4	0.97	0.07	15,16,17,18	0
3	EDO	B	1562	4/4	0.97	0.06	11,11,13,16	0
3	EDO	B	1559	4/4	0.98	0.06	9,10,11,13	0
3	EDO	B	1566[A]	4/4	0.98	0.07	8,10,10,10	4
3	EDO	B	1566[B]	4/4	0.98	0.07	13,13,16,17	4
3	EDO	B	1571	4/4	0.98	0.08	11,12,13,13	0
3	EDO	B	1568	4/4	0.99	0.05	13,13,14,14	0
3	EDO	B	1560	4/4	0.99	0.05	8,8,9,9	0
4	CA	B	1579	1/1	1.00	0.02	7,7,7,7	0

5.5 Other polymers [\(i\)](#)

There are no such residues in this entry.