



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 26, 2023 – 02:24 PM EDT

PDB ID : 3GLS  
Title : Crystal Structure of Human SIRT3  
Authors : Jin, L.; Wei, W.; Jiang, Y.; Peng, H.; Cai, J.; Mao, C.; Dai, H.; Bemis, J.E.;  
Jirousek, M.R.; Milne, J.C.; Westphal, C.H.; Perni, R.B.  
Deposited on : 2009-03-12  
Resolution : 2.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	285	
1	B	285	
1	C	285	
1	D	285	
1	E	285	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	285	 64% 29% • 5%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent deacetylase sirtuin-3, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	273	2144	1382	369	384	9	4	0	0
1	B	272	2135	1376	367	383	9	3	0	0
1	C	273	2144	1382	369	384	9	4	0	0
1	D	274	2148	1384	370	385	9	4	0	0
1	E	274	2148	1384	370	385	9	5	0	0
1	F	272	2135	1376	367	383	9	3	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	115	SER	-	expression tag	UNP Q9NTG7
A	116	ASN	-	expression tag	UNP Q9NTG7
A	117	ALA	-	expression tag	UNP Q9NTG7
B	115	SER	-	expression tag	UNP Q9NTG7
B	116	ASN	-	expression tag	UNP Q9NTG7
B	117	ALA	-	expression tag	UNP Q9NTG7
C	115	SER	-	expression tag	UNP Q9NTG7
C	116	ASN	-	expression tag	UNP Q9NTG7
C	117	ALA	-	expression tag	UNP Q9NTG7
D	115	SER	-	expression tag	UNP Q9NTG7
D	116	ASN	-	expression tag	UNP Q9NTG7
D	117	ALA	-	expression tag	UNP Q9NTG7
E	115	SER	-	expression tag	UNP Q9NTG7
E	116	ASN	-	expression tag	UNP Q9NTG7
E	117	ALA	-	expression tag	UNP Q9NTG7
F	115	SER	-	expression tag	UNP Q9NTG7
F	116	ASN	-	expression tag	UNP Q9NTG7

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	117	ALA	-	expression tag	UNP Q9NTG7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	E	1	Total Zn 1 1	0	0
2	F	1	Total Zn 1 1	0	0

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0
3	F	1	Total C O 7 4 3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	16	Total	O	0	0
			16	16		
5	B	17	Total	O	0	0
			17	17		
5	C	22	Total	O	0	0
			22	22		
5	D	19	Total	O	0	0
			19	19		
5	E	19	Total	O	0	0
			19	19		
5	F	26	Total	O	0	0
			26	26		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD-dependent deacetylase sirtuin-3, mitochondrial




- Molecule 1: NAD-dependent deacetylase sirtuin-3, mitochondrial




- Molecule 1: NAD-dependent deacetylase sirtuin-3, mitochondrial






- Molecule 1: NAD-dependent deacetylase sirtuin-3, mitochondrial

Chain D: 55% (green), 36% (yellow), 5% (orange)




- Molecule 1: NAD-dependent deacetylase sirtuin-3, mitochondrial

Chain E: 63% (green), 31% (yellow), 6% (orange)



- Molecule 1: NAD-dependent deacetylase sirtuin-3, mitochondrial

Chain F: 64% (green), 29% (yellow), 5% (orange)





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.37Å 143.94Å 91.66Å 90.00° 117.10° 90.00°	Depositor
Resolution (Å)	65.00 – 2.70 53.97 – 2.70	Depositor EDS
% Data completeness (in resolution range)	84.7 (65.00-2.70) 84.7 (53.97-2.70)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.69 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.182 , 0.244 0.183 , 0.245	Depositor DCC
$R_{free}$ test set	2313 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.1	Xtrriage
Anisotropy	0.242	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	12998	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.83	0/2199	0.84	4/2996 (0.1%)
1	B	0.87	4/2190 (0.2%)	0.77	0/2985
1	C	0.92	3/2199 (0.1%)	0.80	3/2996 (0.1%)
1	D	0.82	5/2203 (0.2%)	0.76	4/3001 (0.1%)
1	E	0.88	3/2203 (0.1%)	0.77	3/3001 (0.1%)
1	F	0.83	1/2190 (0.0%)	0.79	1/2985 (0.0%)
All	All	0.86	16/13184 (0.1%)	0.79	15/17964 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2
1	D	0	1
All	All	0	3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	332	GLU	CD-OE2	15.46	1.42	1.25
1	C	332	GLU	CD-OE2	15.36	1.42	1.25
1	C	332	GLU	CG-CD	-12.35	1.33	1.51
1	F	385	ASP	CG-OD1	12.12	1.53	1.25
1	C	235	ARG	CZ-NH2	-11.29	1.18	1.33

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	235	ARG	NE-CZ-NH2	13.48	127.04	120.30
1	A	235	ARG	NE-CZ-NH2	-12.34	114.13	120.30
1	A	235	ARG	NE-CZ-NH1	12.22	126.41	120.30
1	D	385	ASP	CB-CG-OD2	8.99	126.39	118.30
1	A	235	ARG	CD-NE-CZ	7.84	134.57	123.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	126	GLN	Sidechain
1	C	389	ARG	Peptide
1	D	380	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2144	0	2160	80	0
1	B	2135	0	2147	77	0
1	C	2144	0	2160	95	0
1	D	2148	0	2163	106	0
1	E	2148	0	2163	80	0
1	F	2135	0	2147	82	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	7	0	9	0	0
3	F	7	0	9	1	0
4	E	5	0	0	1	0
5	A	16	0	0	5	0
5	B	17	0	0	2	0
5	C	22	0	0	3	0
5	D	19	0	0	4	0
5	E	19	0	0	4	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	26	0	0	5	0
All	All	12998	0	12958	507	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 507 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:135:ARG:NH2	4:E:400:SO4:O2	1.75	1.19
1:B:384:ARG:O	1:B:388:GLN:HG2	1.46	1.16
1:C:358:ARG:HG3	1:C:358:ARG:HH11	1.15	1.11
1:F:385:ASP:O	1:F:389:ARG:HG3	1.51	1.09
1:B:125:LEU:HD21	1:B:362:GLN:HG2	1.28	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/285 (95%)	257 (95%)	13 (5%)	1 (0%)	34	60
1	B	270/285 (95%)	256 (95%)	14 (5%)	0	100	100
1	C	271/285 (95%)	256 (94%)	15 (6%)	0	100	100
1	D	272/285 (95%)	252 (93%)	16 (6%)	4 (2%)	10	26
1	E	272/285 (95%)	262 (96%)	9 (3%)	1 (0%)	34	60
1	F	270/285 (95%)	259 (96%)	11 (4%)	0	100	100
All	All	1626/1710 (95%)	1542 (95%)	78 (5%)	6 (0%)	34	60

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	322	LEU
1	D	324	VAL
1	D	290	ASP
1	E	290	ASP
1	D	147	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/246 (96%)	227 (96%)	10 (4%)	30	58
1	B	236/246 (96%)	222 (94%)	14 (6%)	19	43
1	C	237/246 (96%)	223 (94%)	14 (6%)	19	43
1	D	237/246 (96%)	222 (94%)	15 (6%)	18	40
1	E	237/246 (96%)	229 (97%)	8 (3%)	37	66
1	F	236/246 (96%)	222 (94%)	14 (6%)	19	43
All	All	1420/1476 (96%)	1345 (95%)	75 (5%)	22	48

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	279	ARG
1	F	384	ARG
1	E	320	THR
1	F	225	LEU
1	C	126	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	210	HIS
1	E	210	HIS
1	E	169	GLN
1	F	169	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	210	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	E	400	-	4,4,4	0.35	0	6,6,6	0.60	0
3	PGE	F	400	-	6,6,9	0.71	0	5,5,8	0.58	0
3	PGE	A	400	-	6,6,9	0.98	0	5,5,8	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGE	F	400	-	-	3/4/4/7	-
3	PGE	A	400	-	-	3/4/4/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	400	PGE	O1-C1-C2-O2
3	F	400	PGE	O3-C5-C6-O4
3	A	400	PGE	O2-C3-C4-O3
3	F	400	PGE	O2-C3-C4-O3
3	F	400	PGE	C3-C4-O3-C5

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	400	SO4	1	0
3	F	400	PGE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	273/285 (95%)	-0.40	0 <a href="#">100</a>   <a href="#">100</a>	10, 23, 40, 49	5 (1%)
1	B	272/285 (95%)	-0.36	1 (0%) <a href="#">92</a>   <a href="#">93</a>	11, 26, 49, 57	10 (3%)
1	C	273/285 (95%)	-0.38	0 <a href="#">100</a>   <a href="#">100</a>	9, 23, 39, 53	9 (3%)
1	D	274/285 (96%)	-0.24	1 (0%) <a href="#">92</a>   <a href="#">93</a>	17, 31, 52, 61	10 (3%)
1	E	274/285 (96%)	-0.33	0 <a href="#">100</a>   <a href="#">100</a>	12, 25, 37, 44	5 (1%)
1	F	272/285 (95%)	-0.34	1 (0%) <a href="#">92</a>   <a href="#">93</a>	11, 25, 47, 69	7 (2%)
All	All	1638/1710 (95%)	-0.34	3 (0%) <a href="#">95</a>   <a href="#">96</a>	9, 26, 46, 69	46 (2%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	123	LEU	2.9
1	D	383	MET	2.1
1	F	385	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PGE	F	400	7/10	0.91	0.16	19,21,22,22	0
4	SO4	E	400	5/5	0.94	0.21	50,51,53,54	5
3	PGE	A	400	7/10	0.95	0.10	23,24,24,25	0
2	ZN	A	2	1/1	0.98	0.14	38,38,38,38	0
2	ZN	B	4	1/1	0.99	0.11	22,22,22,22	0
2	ZN	D	6	1/1	0.99	0.10	36,36,36,36	0
2	ZN	E	1	1/1	0.99	0.06	32,32,32,32	0
2	ZN	F	3	1/1	1.00	0.10	20,20,20,20	0
2	ZN	C	5	1/1	1.00	0.11	15,15,15,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.