



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 02:37 am BST

PDB ID : 3GME
Title : Crystal Structure of Polynucleotide Phosphorylase in complex with RNase E and manganese
Authors : Nurmohamed, S.; Luisi, B.L.
Deposited on : 2009-03-13
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	
2	D	41	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MN	A	550	-	-	-	X
3	MN	A	551	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyribonucleotide nucleotidyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	482	3560	2239	617	684	20	0	0	0

- Molecule 2 is a protein called Ribonuclease E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	21	134	80	30	24	0	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
3	A	2	2	2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	150	150	150	0	0
4	D	8	8	8	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	158.57Å 158.57Å 156.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.91 – 2.40 25.78 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (25.91-2.40) 99.3 (25.78-2.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.286 , 0.298 0.281 , 0.295	Depositor DCC
R_{free} test set	1490 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtrriage
Anisotropy	0.106	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3854	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.54	2/3615 (0.1%)	0.47	0/4909
2	D	0.32	0/139	0.42	0/191
All	All	0.53	2/3754 (0.1%)	0.47	0/5100

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	63	TYR	CD1-CE1	-5.57	1.30	1.39
1	A	31	VAL	CB-CG2	-5.49	1.41	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3560	0	3508	59	0
2	D	134	0	119	2	0
3	A	2	0	0	0	0
4	A	150	0	0	2	0
4	D	8	0	0	1	0
All	All	3854	0	3627	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:LEU:HD11	1:A:349:GLN:HE21	1.05	1.10
1:A:406:LEU:HD11	1:A:506:GLN:OE1	1.67	0.93
1:A:342:LEU:HD11	1:A:349:GLN:NE2	1.84	0.92
1:A:202:LEU:O	1:A:202:LEU:HD12	1.72	0.89
1:A:342:LEU:CD1	1:A:349:GLN:HE21	1.89	0.83
1:A:493:PHE:HZ	1:A:523:LEU:CD2	1.97	0.78
1:A:375:THR:HG22	4:A:628:HOH:O	1.82	0.78
1:A:406:LEU:CD1	1:A:506:GLN:OE1	2.32	0.77
1:A:396:PRO:HA	1:A:400:GLU:OE1	1.84	0.77
1:A:519:MET:O	1:A:523:LEU:HD23	1.84	0.76
1:A:470:LYS:HE2	1:A:472:GLY:O	1.90	0.70
1:A:308:ARG:HG2	1:A:313:GLU:OE1	1.90	0.70
1:A:544:PRO:O	1:A:545:ARG:C	2.30	0.69
1:A:493:PHE:CZ	1:A:523:LEU:CD2	2.76	0.69
1:A:493:PHE:CZ	1:A:523:LEU:HD22	2.28	0.68
1:A:188:MET:O	1:A:189:VAL:HG23	1.93	0.68
1:A:493:PHE:HZ	1:A:523:LEU:HD21	1.57	0.68
1:A:16:VAL:HG22	1:A:35:MET:HG3	1.79	0.63
1:A:456:PRO:HA	1:A:544:PRO:HG2	1.82	0.60
1:A:375:THR:CG2	4:A:628:HOH:O	2.45	0.59
1:A:466:MET:HB2	1:A:493:PHE:HB3	1.84	0.58
1:A:181:GLY:HA2	1:A:187:LEU:HD12	1.87	0.56
1:A:202:LEU:C	1:A:202:LEU:HD12	2.17	0.55
1:A:144:PRO:HD3	1:A:231:TRP:HB2	1.88	0.55
1:A:325:ARG:HD2	1:A:346:GLY:HA3	1.87	0.55
1:A:477:VAL:HB	1:A:525:GLN:OE1	2.06	0.55
1:A:188:MET:O	1:A:189:VAL:CG2	2.55	0.54
2:D:1056:ARG:HD2	4:D:118:HOH:O	2.09	0.53
1:A:397:LYS:O	1:A:398:ARG:C	2.46	0.52
1:A:33:VAL:HG21	1:A:133:GLY:CA	2.40	0.51
1:A:493:PHE:CZ	1:A:523:LEU:HD21	2.43	0.51
1:A:478:LEU:N	1:A:478:LEU:HD12	2.26	0.51
1:A:188:MET:CG	1:A:189:VAL:N	2.73	0.51
1:A:188:MET:HG2	1:A:189:VAL:N	2.25	0.51
1:A:534:LEU:O	1:A:538:GLU:HG3	2.11	0.51
1:A:397:LYS:O	1:A:399:ARG:N	2.45	0.50
1:A:64:GLN:HE21	1:A:116:THR:HB	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:THR:HB	2:D:1048:HIS:HB2	1.94	0.49
1:A:322:ASP:HA	1:A:525:GLN:HE21	1.78	0.48
1:A:342:LEU:HD11	1:A:349:GLN:CG	2.43	0.48
1:A:33:VAL:HG21	1:A:133:GLY:HA2	1.96	0.47
1:A:2:LEU:HD12	1:A:2:LEU:N	2.32	0.45
1:A:478:LEU:N	1:A:478:LEU:CD1	2.80	0.45
1:A:20:THR:OG1	1:A:21:GLY:N	2.50	0.43
1:A:351:LEU:HB3	1:A:430:GLU:HB2	2.00	0.43
1:A:519:MET:O	1:A:523:LEU:CD2	2.63	0.43
1:A:28:THR:HG21	1:A:143:ILE:HD11	2.01	0.43
1:A:44:VAL:HG22	1:A:113:VAL:HG22	2.01	0.43
1:A:2:LEU:HB3	1:A:22:MET:HG3	2.01	0.42
1:A:189:VAL:HB	1:A:507:MET:HB2	2.01	0.42
1:A:91:ILE:HA	1:A:94:LEU:HD12	2.00	0.42
1:A:316:ILE:HG13	1:A:316:ILE:H	1.72	0.42
1:A:502:ILE:HG12	1:A:530:ARG:HH22	1.85	0.42
1:A:543:ALA:HA	1:A:544:PRO:HD3	1.88	0.42
1:A:322:ASP:HB3	1:A:525:GLN:NE2	2.35	0.42
1:A:406:LEU:CG	1:A:506:GLN:OE1	2.67	0.42
1:A:4:PRO:HB3	1:A:19:GLU:OE2	2.20	0.41
1:A:509:ILE:HG21	1:A:514:ILE:HG21	2.02	0.41
1:A:3:ASN:HA	1:A:4:PRO:HD3	1.93	0.41
1:A:67:THR:HG21	1:A:74:PRO:HG3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	472/549 (86%)	447 (95%)	24 (5%)	1 (0%)	47 62
2	D	19/41 (46%)	18 (95%)	1 (5%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	491/590 (83%)	465 (95%)	25 (5%)	1 (0%)	47 62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	369/446 (83%)	366 (99%)	3 (1%)	81 91
2	D	10/28 (36%)	10 (100%)	0	100 100
All	All	379/474 (80%)	376 (99%)	3 (1%)	81 91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	202	LEU
1	A	342	LEU
1	A	406	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	64	GLN
1	A	213	GLN
1	A	349	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	482/549 (87%)	0.87	67 (13%) 2 2	24, 45, 90, 112	1 (0%)
2	D	21/41 (51%)	0.84	4 (19%) 1 1	36, 52, 71, 80	0
All	All	503/590 (85%)	0.87	71 (14%) 2 2	24, 46, 90, 112	1 (0%)

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	241	ALA	8.4
1	A	238	VAL	7.6
1	A	53	GLY	7.2
1	A	55	ASP	6.7
1	A	267	TYR	6.0
1	A	314	PRO	5.5
1	A	398	ARG	5.4
1	A	263	LYS	5.2
1	A	395	SER	5.1
1	A	394	GLY	5.1
1	A	266	ARG	5.1
1	A	244	ALA	5.0
1	A	396	PRO	4.8
1	A	77	PHE	4.8
1	A	52	PRO	4.8
1	A	543	ALA	4.7
1	A	76	SER	4.6
1	A	261	THR	4.5
1	A	237	PRO	4.4
1	A	234	GLN	4.2
1	A	265	GLU	4.2
1	A	1	MET	4.0
1	A	297	LEU	3.9
1	A	301	GLU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	243	ASN	3.9
2	D	1059	PRO	3.9
1	A	246	VAL	3.8
1	A	298	HIS	3.8
1	A	232	ASP	3.7
1	A	262	ASP	3.7
1	A	56	PHE	3.5
1	A	239	ASN	3.4
1	A	300	ILE	3.4
1	A	242	LEU	3.1
1	A	304	VAL	3.1
1	A	264	GLN	3.1
1	A	474	ASN	3.0
1	A	236	GLU	2.9
1	A	83	ARG	2.9
1	A	49	LYS	2.9
1	A	493	PHE	2.9
1	A	224	LYS	2.9
1	A	542	ASN	2.8
1	A	517	GLU	2.8
1	A	514	ILE	2.8
1	A	544	PRO	2.8
1	A	75	GLY	2.7
1	A	464	ILE	2.7
1	A	513	GLY	2.6
1	A	188	MET	2.5
1	A	44	VAL	2.5
1	A	3	ASN	2.5
1	A	473	ASP	2.4
1	A	133	GLY	2.4
1	A	142	GLY	2.4
2	D	1039	GLY	2.4
2	D	1040	ALA	2.4
1	A	132	ILE	2.3
1	A	545	ARG	2.3
1	A	487	HIS	2.3
1	A	13	GLN	2.2
1	A	316	ILE	2.2
1	A	496	ALA	2.2
2	D	1055	ALA	2.2
1	A	321	LYS	2.2
1	A	235	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	466	MET	2.2
1	A	31	VAL	2.1
1	A	499	ARG	2.1
1	A	138	LEU	2.1
1	A	33	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	A	550	1/1	0.43	0.62	228,228,228,228	0
3	MN	A	551	1/1	0.47	0.49	82,82,82,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.