



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 23, 2024 – 10:02 PM EDT

PDB ID : 1GPO
Title : CRYSTAL STRUCTURE OF THE RATIONALLY DESIGNED ANTIBODY
M41 AS A FAB FRAGMENT
Authors : Schiweck, W.; Skerra, A.
Deposited on : 1997-03-27
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	219	
1	M	219	
2	H	221	
2	I	221	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7281 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANTIBODY M41.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	216	1676	1046	279	346	5	2	0	0
1	M	218	1693	1055	282	351	5	78	0	0

- Molecule 2 is a protein called ANTIBODY M41.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	214	1636	1033	266	331	6	2	0	0
2	I	212	1617	1023	261	327	6	3	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	3	LYS	GLN	conflict	EMBL AJ303449
H	32	ASP	GLY	conflict	EMBL AJ303449
H	33	PHE	TYR	conflict	EMBL AJ303449
H	35	SER	ASN	conflict	EMBL AJ303449
H	39	GLN	LYS	conflict	EMBL AJ303449
H	44	ARG	LYS	conflict	EMBL AJ303449
H	50	PHE	TYR	conflict	EMBL AJ303449
H	51	VAL	ILE	conflict	EMBL AJ303449
H	52	GLN	SER	conflict	EMBL AJ303449
H	54	SER	GLY	conflict	EMBL AJ303449
H	56	GLU	SER	conflict	EMBL AJ303449
H	58	ALA	TYR	conflict	EMBL AJ303449
H	64	LYS	GLU	conflict	EMBL AJ303449
H	81	ASP	GLN	conflict	EMBL AJ303449
H	92	VAL	THR	conflict	EMBL AJ303449
H	94	TYR	PHE	conflict	EMBL AJ303449

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	97	ASN	ARG	conflict	EMBL AJ303449
H	98	TRP	LEU	conflict	EMBL AJ303449
H	99	HIS	PHE	conflict	EMBL AJ303449
H	?	-	SER	deletion	EMBL AJ303449
H	?	-	TYR	deletion	EMBL AJ303449
H	?	-	TYR	deletion	EMBL AJ303449
H	?	-	PHE	deletion	EMBL AJ303449
H	109	VAL	LEU	conflict	EMBL AJ303449
I	3	LYS	GLN	conflict	EMBL AJ303449
I	32	ASP	GLY	conflict	EMBL AJ303449
I	33	PHE	TYR	conflict	EMBL AJ303449
I	35	SER	ASN	conflict	EMBL AJ303449
I	39	GLN	LYS	conflict	EMBL AJ303449
I	44	ARG	LYS	conflict	EMBL AJ303449
I	50	PHE	TYR	conflict	EMBL AJ303449
I	51	VAL	ILE	conflict	EMBL AJ303449
I	52	GLN	SER	conflict	EMBL AJ303449
I	54	SER	GLY	conflict	EMBL AJ303449
I	56	GLU	SER	conflict	EMBL AJ303449
I	58	ALA	TYR	conflict	EMBL AJ303449
I	64	LYS	GLU	conflict	EMBL AJ303449
I	81	ASP	GLN	conflict	EMBL AJ303449
I	92	VAL	THR	conflict	EMBL AJ303449
I	94	TYR	PHE	conflict	EMBL AJ303449
I	97	ASN	ARG	conflict	EMBL AJ303449
I	98	TRP	LEU	conflict	EMBL AJ303449
I	99	HIS	PHE	conflict	EMBL AJ303449
I	?	-	SER	deletion	EMBL AJ303449
I	?	-	TYR	deletion	EMBL AJ303449
I	?	-	TYR	deletion	EMBL AJ303449
I	?	-	PHE	deletion	EMBL AJ303449
I	109	VAL	LEU	conflict	EMBL AJ303449

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	H	1	Total O S 5 4 1	0	0
3	I	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	174	Total O 174 174	0	0
4	H	185	Total O 185 185	0	0
4	M	134	Total O 134 134	0	0
4	I	156	Total O 156 156	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ANTIBODY M41




- Molecule 1: ANTIBODY M41



- Molecule 2: ANTIBODY M41



- Molecule 2: ANTIBODY M41



W154	W155	S156	V169	L170	D173	L174	Y175	T176	L177	W188	T192	C195	W196	H199	S202	S203	T204	V211	P212	ARG	ASP	CYS	HIS	HIS	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.47Å 103.51Å 113.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.95 8.00 – 1.96	Depositor EDS
% Data completeness (in resolution range)	81.3 (8.00-1.95) 83.8 (8.00-1.96)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 1.96Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.214 , 0.283 0.206 , 0.266	Depositor DCC
R_{free} test set	6871 reflections (10.14%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtrriage
Anisotropy	0.381	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.49 , 133.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7281	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.07 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.3991e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.53	0/1714	0.77	0/2330
1	M	0.56	0/1731	0.76	0/2353
2	H	0.58	0/1681	0.84	1/2307 (0.0%)
2	I	0.57	0/1662	0.81	1/2282 (0.0%)
All	All	0.56	0/6788	0.80	2/9272 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	131	GLN	N-CA-C	7.63	131.60	111.00
2	I	177	LEU	CA-CB-CG	5.66	128.32	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1676	0	1600	25	0
1	M	1693	0	1612	33	0
2	H	1636	0	1578	23	0
2	I	1617	0	1561	24	0
3	H	5	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	I	5	0	0	1	0
4	H	185	0	0	2	0
4	I	156	0	0	2	0
4	L	174	0	0	2	0
4	M	134	0	0	2	0
All	All	7281	0	6351	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 103 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:153:THR:HG22	2:I:196:ASN:HD22	1.34	0.91
1:M:195:ASN:HA	1:M:216:ARG:HG3	1.56	0.88
2:H:156:SER:H	2:H:196:ASN:HD21	1.30	0.79
1:L:95:GLN:HE22	1:L:98:ASP:H	1.33	0.77
1:M:95:GLN:HE22	1:M:98:ASP:H	1.35	0.72

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/219 (98%)	202 (94%)	12 (6%)	0	100	100
1	M	216/219 (99%)	201 (93%)	14 (6%)	1 (0%)	29	17
2	H	212/221 (96%)	197 (93%)	11 (5%)	4 (2%)	8	2
2	I	210/221 (95%)	195 (93%)	13 (6%)	2 (1%)	15	6
All	All	852/880 (97%)	795 (93%)	50 (6%)	7 (1%)	19	9

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	64	LYS
2	H	130	ALA
2	H	128	SER
1	M	195	ASN
2	I	128	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	194/197 (98%)	182 (94%)	12 (6%)	18	7
1	M	196/197 (100%)	181 (92%)	15 (8%)	13	4
2	H	191/198 (96%)	176 (92%)	15 (8%)	12	3
2	I	189/198 (96%)	178 (94%)	11 (6%)	20	8
All	All	770/790 (98%)	717 (93%)	53 (7%)	15	5

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	14	THR
1	M	138	VAL
2	I	153	THR
1	M	18	SER
1	M	65	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	17	ASN
1	M	129	GLN
2	I	196	ASN
1	M	95	GLN
1	M	142	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	I	651	-	4,4,4	0.85	0	6,6,6	0.32	0
3	SO4	H	650	-	4,4,4	0.68	0	6,6,6	0.31	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	651	SO4	1	0
3	H	650	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	216/219 (98%)	-0.22	1 (0%) 91 94	9, 24, 42, 64	1 (0%)
1	M	208/219 (94%)	0.28	19 (9%) 9 15	10, 29, 67, 87	2 (0%)
2	H	214/221 (96%)	-0.25	7 (3%) 46 56	8, 20, 42, 75	2 (0%)
2	I	212/221 (95%)	-0.08	9 (4%) 36 45	9, 24, 46, 81	3 (1%)
All	All	850/880 (96%)	-0.07	36 (4%) 36 45	8, 24, 57, 87	8 (0%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	1	GLU	6.2
2	H	132	THR	5.6
2	H	130	ALA	5.4
1	M	32	GLU	5.3
1	M	217	ASN	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	H	650	5/5	0.99	0.08	27,31,35,36	0
3	SO4	I	651	5/5	0.99	0.05	21,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.