



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 3, 2023 – 11:40 pm GMT

PDB ID : 1GPW  
Title : Structural evidence for ammonia tunneling across the (beta/alpha)<sub>8</sub> barrel of the imidazole glycerol phosphate synthase bienzyme complex.  
Authors : Walker, M.; Beismann-Driemeyer, S.; Sterner, R.; Wilmanns, M.  
Deposited on : 2001-11-12  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	253	69% 26% 5%
1	C	253	66% 30% 5%
1	E	253	34% 51% 14% .
2	B	201	51% 45% . .
2	D	201	59% 36% 5%
2	F	201	49% 46% 5%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISF PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	253	1937	1232	330	369	6	0	0	0
1	C	253	1937	1230	331	370	6	0	0	0
1	E	251	1927	1225	328	368	6	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	ASN	ASP	engineered mutation	UNP Q9X0C6
C	11	ASN	ASP	engineered mutation	UNP Q9X0C6
E	11	ASN	ASP	engineered mutation	UNP Q9X0C6

- Molecule 2 is a protein called AMIDOTRANSFERASE HISH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	201	1616	1029	285	294	8	0	0	1
2	D	201	1612	1028	281	295	8	0	0	0
2	F	201	1614	1029	282	295	8	0	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	117	Total O 117 117	0	0
4	B	49	Total O 49 49	0	0
4	C	98	Total O 98 98	0	0
4	D	84	Total O 84 84	0	0
4	E	40	Total O 40 40	0	0
4	F	71	Total O 71 71	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: HISF PROTEIN

Chain A: 




- Molecule 1: HISF PROTEIN


Chain C: 



- Molecule 1: HISF PROTEIN

Chain E: 






- Molecule 2: AMIDOTRANSFERASE HISH

Chain B: 51% 45%



- Molecule 2: AMIDOTRANSFERASE HISH

Chain D: 59% 36% 5%



- Molecule 2: AMIDOTRANSFERASE HISH

Chain F: 49% 46% 5%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.00Å 91.90Å 131.00Å 90.00° 105.90° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40	Depositor
% Data completeness (in resolution range)	97.8 (30.00-2.40)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.226 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	11132	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1964	0.67	0/2648
1	C	0.43	0/1963	0.67	0/2647
1	E	0.46	0/1952	0.69	0/2630
2	B	0.45	0/1650	0.67	0/2220
2	D	0.44	0/1646	0.68	0/2214
2	F	0.43	0/1648	0.67	0/2217
All	All	0.44	0/10823	0.68	0/14576

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1937	0	1967	69	0
1	C	1937	0	1973	71	1
1	E	1927	0	1962	207	1
2	B	1616	0	1611	90	1
2	D	1612	0	1598	72	0
2	F	1614	0	1599	95	1
3	A	10	0	0	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	15	0	0	0	0
3	E	5	0	0	0	0
4	A	117	0	0	4	0
4	B	49	0	0	6	0
4	C	98	0	0	4	0
4	D	84	0	0	6	0
4	E	40	0	0	4	0
4	F	71	0	0	6	0
All	All	11132	0	10710	590	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 590 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:53:THR:HG21	1:E:57:GLU:HB2	1.34	1.07
1:E:9:CYS:HB2	1:E:222:LEU:HD11	1.34	1.06
1:E:53:THR:HG22	1:E:54:ALA:H	1.18	1.04
1:E:123:GLN:HE21	1:E:123:GLN:N	1.60	1.00
2:F:1:MET:HE2	2:F:194:ILE:HG23	1.41	0.99

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:55:SER:N	2:F:200:ARG:CB[1_456]	1.73	0.47
2:B:114:ARG:NH2	1:E:208:GLU:OE2[1_455]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/253 (99%)	231 (92%)	14 (6%)	6 (2%)	6	6
1	C	251/253 (99%)	228 (91%)	18 (7%)	5 (2%)	7	9
1	E	247/253 (98%)	184 (74%)	35 (14%)	28 (11%)	0	0
2	B	199/201 (99%)	167 (84%)	25 (13%)	7 (4%)	3	3
2	D	199/201 (99%)	171 (86%)	20 (10%)	8 (4%)	3	2
2	F	199/201 (99%)	177 (89%)	13 (6%)	9 (4%)	2	2
All	All	1346/1362 (99%)	1158 (86%)	125 (9%)	63 (5%)	2	1

5 of 63 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	GLU
1	A	55	SER
1	A	251	GLU
2	B	200	ARG
1	C	19	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/208 (98%)	193 (95%)	11 (5%)	22	36
1	C	205/208 (99%)	192 (94%)	13 (6%)	18	28
1	E	204/208 (98%)	184 (90%)	20 (10%)	8	11
2	B	176/177 (99%)	165 (94%)	11 (6%)	18	28
2	D	174/177 (98%)	166 (95%)	8 (5%)	27	43
2	F	174/177 (98%)	162 (93%)	12 (7%)	15	25
All	All	1137/1155 (98%)	1062 (93%)	75 (7%)	16	26

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	148	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	F	167	VAL
1	E	229	PHE
2	F	29	ASP
1	C	31	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	123	GLN
2	D	88	GLN
2	F	88	GLN
2	D	15	ASN
1	E	103	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	E	301	-	4,4,4	1.49	0	6,6,6	0.41	0
3	PO4	C	302	-	4,4,4	1.40	0	6,6,6	0.43	0
3	PO4	A	301	-	4,4,4	1.59	0	6,6,6	0.41	0
3	PO4	C	303	-	4,4,4	1.29	0	6,6,6	0.44	0
3	PO4	C	301	-	4,4,4	1.43	0	6,6,6	0.46	0
3	PO4	A	302	-	4,4,4	1.23	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.