



Full wwPDB X-ray Structure Validation Report i

Mar 5, 2024 – 10:08 PM JST

PDB ID : 8GPY

Title : Crystal structure of Omicron BA.4/5 RBD in complex with a neutralizing antibody scFv

Authors : Gao, Y.X.; Song, Z.D.; Wang, W.M.; Guo, Y.

Deposited on : 2022-08-27

Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Xtriage (Phenix) : 1.13

EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 6558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	191	1524	982	258	276	8	0	0	0
1	B	180	1458	940	245	265	8	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	339	ASP	GLY	variant	UNP P0DTC2
A	371	PHE	SER	variant	UNP P0DTC2
A	373	PRO	SER	variant	UNP P0DTC2
A	375	PHE	SER	variant	UNP P0DTC2
A	376	ALA	THR	variant	UNP P0DTC2
A	405	ASN	ASP	variant	UNP P0DTC2
A	408	SER	ARG	variant	UNP P0DTC2
A	417	ASN	LYS	variant	UNP P0DTC2
A	440	LYS	ASN	variant	UNP P0DTC2
A	452	ARG	LEU	variant	UNP P0DTC2
A	477	ASN	SER	variant	UNP P0DTC2
A	478	LYS	THR	variant	UNP P0DTC2
A	484	ALA	GLU	variant	UNP P0DTC2
A	486	VAL	PHE	variant	UNP P0DTC2
A	498	ARG	GLN	variant	UNP P0DTC2
A	501	TYR	ASN	variant	UNP P0DTC2
A	505	HIS	TYR	variant	UNP P0DTC2
A	531	ALA	-	expression tag	UNP P0DTC2
A	532	ALA	-	expression tag	UNP P0DTC2
A	533	ALA	-	expression tag	UNP P0DTC2
B	339	ASP	GLY	variant	UNP P0DTC2
B	371	PHE	SER	variant	UNP P0DTC2
B	373	PRO	SER	variant	UNP P0DTC2
B	375	PHE	SER	variant	UNP P0DTC2
B	376	ALA	THR	variant	UNP P0DTC2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	405	ASN	ASP	variant	UNP P0DTC2
B	408	SER	ARG	variant	UNP P0DTC2
B	417	ASN	LYS	variant	UNP P0DTC2
B	440	LYS	ASN	variant	UNP P0DTC2
B	452	ARG	LEU	variant	UNP P0DTC2
B	477	ASN	SER	variant	UNP P0DTC2
B	478	LYS	THR	variant	UNP P0DTC2
B	484	ALA	GLU	variant	UNP P0DTC2
B	486	VAL	PHE	variant	UNP P0DTC2
B	498	ARG	GLN	variant	UNP P0DTC2
B	501	TYR	ASN	variant	UNP P0DTC2
B	505	HIS	TYR	variant	UNP P0DTC2
B	531	ALA	-	expression tag	UNP P0DTC2
B	532	ALA	-	expression tag	UNP P0DTC2
B	533	ALA	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called scFv.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	227	Total	C	N	O	S	27	0	0
			1704	1068	286	341	9			
2	E	230	Total	C	N	O	S	24	0	0
			1723	1079	290	345	9			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	16	Total	O	0	0
			16	16		
3	B	11	Total	O	0	0
			11	11		
3	C	53	Total	O	0	0
			53	53		
3	E	69	Total	O	0	0
			69	69		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 1: Spike protein S1



- Molecule 2: scFv



- Molecule 2: scFv





4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.86 Å 92.09 Å 89.27 Å 90.00° 92.40° 90.00°	Depositor
Resolution (Å)	49.60 – 2.51 49.60 – 2.51	Depositor EDS
% Data completeness (in resolution range)	71.8 (49.60-2.51) 81.6 (49.60-2.51)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	0.88 (at 2.51 Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R , R_{free}	0.215 , 0.251 0.217 , 0.255	Depositor DCC
R_{free} test set	2003 reflections (6.69%)	wwPDB-VP
Wilson B-factor (Å ²)	36.1	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6558	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.30	0/1570	0.58	0/2136
1	B	0.32	0/1499	0.59	0/2033
2	C	0.31	0/1744	0.58	0/2370
2	E	0.37	0/1763	0.61	0/2396
All	All	0.33	0/6576	0.59	0/8935

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1524	0	1448	42	0
1	B	1458	0	1376	23	0
2	C	1704	0	1615	18	0
2	E	1723	0	1633	15	0
3	A	16	0	0	0	0
3	B	11	0	0	1	0
3	C	53	0	0	2	0
3	E	69	0	0	0	0
All	All	6558	0	6072	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:LEU:HA	1:A:390:LEU:HD21	1.22	1.16
1:A:387:LEU:HA	1:A:390:LEU:CD2	1.82	1.10
1:A:387:LEU:CD2	1:A:390:LEU:HD21	1.87	1.04
1:A:387:LEU:HD23	1:A:390:LEU:CD2	1.87	1.03
1:A:390:LEU:HB3	1:A:392:PHE:CD2	1.95	1.02
1:A:387:LEU:HD23	1:A:390:LEU:HD21	1.41	0.99
1:A:387:LEU:CA	1:A:390:LEU:HD21	1.97	0.94
1:A:387:LEU:CD2	1:A:390:LEU:CD2	2.49	0.89
1:A:390:LEU:HB3	1:A:392:PHE:CE2	2.08	0.88
1:A:390:LEU:HD22	1:A:392:PHE:HE2	1.38	0.86
2:C:34:MET:SD	2:C:98:ASN:ND2	2.53	0.82
1:A:384:PRO:HA	1:A:387:LEU:HG	1.67	0.77
1:A:360:ASN:H	1:A:523:THR:HG22	1.51	0.76
1:A:390:LEU:HB3	1:A:392:PHE:HD2	1.52	0.73
1:A:393:THR:CG2	1:A:520:ALA:O	2.38	0.72
1:A:390:LEU:HD22	1:A:392:PHE:CE2	2.25	0.69
2:C:156:SER:HB3	2:C:211:CYS:HB3	1.75	0.69
1:A:390:LEU:CD2	1:A:392:PHE:HE2	2.09	0.66
2:C:157:TRP:HB2	2:C:170:ILE:HB	1.77	0.65
1:B:474:GLN:HG3	1:B:480:CYS:HB2	1.81	0.63
2:E:39:GLN:O	2:E:92:ALA:HB1	2.00	0.62
1:A:393:THR:HG21	1:A:520:ALA:O	2.00	0.61
1:A:387:LEU:HD22	1:A:390:LEU:HD21	1.81	0.61
2:E:34:MET:HB3	2:E:79:LEU:HD22	1.83	0.61
1:B:369:TYR:CD2	1:B:370:ASN:N	2.70	0.60
1:A:387:LEU:HD22	1:A:390:LEU:CD2	2.33	0.59
1:A:393:THR:HG22	1:A:520:ALA:O	2.01	0.59
1:B:354:ASN:ND2	3:B:601:HOH:O	2.36	0.59
1:A:389:ASP:OD1	1:A:389:ASP:N	2.36	0.59
2:C:142:CYS:O	2:C:192:THR:HA	2.03	0.58
2:C:87:ARG:NH1	3:C:305:HOH:O	2.36	0.58
1:B:409:GLN:HA	1:B:414:GLN:HG2	1.87	0.55
1:A:387:LEU:CD2	1:A:390:LEU:HD22	2.35	0.55
2:C:13:GLN:NE2	3:C:303:HOH:O	2.31	0.55
1:A:354:ASN:OD1	1:B:360:ASN:ND2	2.40	0.55
1:A:387:LEU:HA	1:A:390:LEU:HD23	1.83	0.54
2:C:83:MET:HB3	2:C:86:LEU:HD21	1.89	0.54
2:C:91:THR:HG23	2:C:117:THR:HA	1.89	0.54
2:E:157:TRP:HB2	2:E:170:ILE:HB	1.90	0.54
2:C:34:MET:HB3	2:C:79:LEU:HD22	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:LEU:CB	1:A:390:LEU:HD21	2.37	0.53
1:A:387:LEU:HD23	1:A:390:LEU:HD22	1.87	0.52
1:B:384:PRO:O	1:B:387:LEU:HB2	2.09	0.52
1:A:447:GLY:HA2	1:A:498:ARG:HG2	1.93	0.51
1:B:358:ILE:HB	1:B:395:VAL:HB	1.93	0.50
1:B:425:LEU:HD21	1:B:512:VAL:HG11	1.92	0.50
2:C:22:CYS:HB3	2:C:79:LEU:HB3	1.94	0.49
1:A:368:LEU:HA	1:A:371:PHE:HB2	1.94	0.49
1:A:390:LEU:CB	1:A:392:PHE:CE2	2.90	0.49
1:A:398:ASP:HB2	1:A:512:VAL:HB	1.94	0.49
1:A:387:LEU:CA	1:A:390:LEU:CD2	2.70	0.49
1:B:461:LEU:HD22	1:B:465:GLU:HB3	1.96	0.48
2:E:51:ILE:HG12	2:E:72:ARG:HD2	1.95	0.48
1:B:349:SER:O	1:B:352:ALA:O	2.31	0.48
1:A:347:PHE:CE2	1:A:399:SER:HB2	2.49	0.47
1:A:349:SER:O	1:A:352:ALA:O	2.33	0.46
1:A:365:TYR:OH	1:A:390:LEU:HD11	2.16	0.46
1:B:359:SER:HA	1:B:524:VAL:HG22	1.97	0.46
1:B:372:ALA:HB1	1:B:373:PRO:HD2	1.98	0.46
1:B:334:ASN:O	1:B:334:ASN:OD1	2.34	0.46
1:B:472:ILE:H	1:B:472:ILE:HG12	1.54	0.45
1:A:387:LEU:O	1:A:390:LEU:HG	2.16	0.45
1:B:401:VAL:HG22	1:B:509:ARG:HG2	1.98	0.45
1:B:362:VAL:HA	1:B:525:CYS:O	2.17	0.45
2:C:64:VAL:HG13	2:C:68:PHE:HB2	1.98	0.45
2:E:22:CYS:HB3	2:E:79:LEU:HB3	1.97	0.44
1:B:478:LYS:O	1:B:479:PRO:C	2.55	0.44
2:C:168:LEU:O	2:C:177:PRO:HG3	2.17	0.44
2:E:60:TYR:HB2	2:E:65:LYS:HE3	2.00	0.43
2:E:144:GLY:O	2:E:191:ASN:HB3	2.18	0.43
2:C:102:THR:OG1	2:C:103:ALA:N	2.51	0.43
2:E:143:THR:HG23	2:E:192:THR:OG1	2.18	0.43
2:E:18:LEU:HB2	2:E:86:LEU:HD11	2.01	0.43
1:B:350:VAL:HG22	1:B:422:ASN:HB3	2.00	0.43
2:C:125:THR:OG1	2:C:143:THR:HB	2.18	0.43
1:A:360:ASN:N	1:A:523:THR:HG22	2.27	0.43
1:B:383:SER:HA	1:B:384:PRO:HD3	1.79	0.42
2:C:107:PHE:O	2:C:168:LEU:HD13	2.20	0.42
2:E:91:THR:HG23	2:E:117:THR:HA	2.00	0.42
2:C:107:PHE:O	2:C:168:LEU:CD1	2.68	0.42
2:E:142:CYS:O	2:E:192:THR:HA	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:387:LEU:C	1:A:390:LEU:HG	2.39	0.42
1:B:379:CYS:SG	1:B:384:PRO:HG3	2.60	0.41
1:A:374:PHE:HD1	1:A:436:TRP:HB3	1.84	0.41
1:B:369:TYR:CG	1:B:370:ASN:N	2.82	0.41
1:A:439:ASN:HA	1:A:507:PRO:HG2	2.02	0.41
2:E:99:GLY:HA3	2:E:103:ALA:O	2.20	0.41
2:E:51:ILE:HD12	2:E:58:THR:HG22	2.03	0.41
1:A:421:TYR:O	1:A:461:LEU:HD11	2.21	0.41
1:A:517:LEU:HD23	1:A:517:LEU:HA	1.87	0.40
1:B:426:PRO:HG3	1:B:463:PRO:HB3	2.03	0.40
1:B:384:PRO:HA	1:B:387:LEU:HD12	2.03	0.40
2:C:86:LEU:HD23	2:C:86:LEU:HA	1.85	0.40
2:C:156:SER:O	2:C:211:CYS:N	2.50	0.40
2:E:12:VAL:HG11	2:E:18:LEU:HG	2.03	0.40
1:A:370:ASN:HD22	1:A:370:ASN:HA	1.69	0.40
2:E:52:SER:HB3	2:E:57:SER:HB2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	189/201 (94%)	183 (97%)	6 (3%)	0	100 100
1	B	172/201 (86%)	163 (95%)	8 (5%)	1 (1%)	25 43
2	C	223/232 (96%)	214 (96%)	9 (4%)	0	100 100
2	E	226/232 (97%)	217 (96%)	9 (4%)	0	100 100
All	All	810/866 (94%)	777 (96%)	32 (4%)	1 (0%)	51 73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	372	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	163/170 (96%)	156 (96%)	7 (4%)	29 53
1	B	158/170 (93%)	153 (97%)	5 (3%)	39 65
2	C	182/184 (99%)	179 (98%)	3 (2%)	62 84
2	E	183/184 (100%)	171 (93%)	12 (7%)	16 32
All	All	686/708 (97%)	659 (96%)	27 (4%)	32 57

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	359	SER
1	A	367	VAL
1	A	370	ASN
1	A	371	PHE
1	A	394	ASN
1	A	487	ASN
1	A	523	THR
1	B	343	ASN
1	B	375	PHE
1	B	383	SER
1	B	471	GLU
1	B	514	SER
2	C	64	VAL
2	C	147	SER
2	C	148	ASP
2	E	19	ARG
2	E	63	SER
2	E	64	VAL
2	E	65	LYS
2	E	67	ARG
2	E	119	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	148	ASP
2	E	156	SER
2	E	194	SER
2	E	212	SER
2	E	216	THR
2	E	229	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	354	ASN
1	A	370	ASN
1	B	360	ASN
2	C	160	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/201 (95%)	0.98	32 (16%) 1 1	28, 49, 98, 105	0
1	B	180/201 (89%)	0.73	26 (14%) 2 2	31, 53, 91, 101	0
2	C	227/232 (97%)	-0.08	2 (0%) 84 86	24, 33, 52, 70	9 (3%)
2	E	230/232 (99%)	-0.10	0 100 100	20, 32, 53, 63	8 (3%)
All	All	828/866 (95%)	0.34	60 (7%) 15 16	20, 40, 88, 105	17 (2%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	369	TYR	9.2
1	A	391	CYS	8.2
1	A	525	CYS	7.5
1	A	390	LEU	6.0
1	B	389	ASP	6.0
1	B	369	TYR	6.0
1	A	392	PHE	5.8
1	A	521	PRO	5.8
1	A	524	VAL	5.6
1	A	478	LYS	5.4
1	A	366	SER	5.3
1	A	518	LEU	5.3
1	A	522	ALA	4.8
1	A	520	ALA	4.8
1	A	387	LEU	4.6
1	A	481	ASN	4.4
1	A	479	PRO	4.1
1	A	362	VAL	4.0
1	B	456	PHE	3.7
1	A	517	LEU	3.7
1	B	474	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	480	CYS	3.4
1	B	374	PHE	3.4
1	B	370	ASN	3.4
1	B	382	VAL	3.3
1	B	371	PHE	3.3
1	B	358	ILE	3.2
2	C	120	SER	3.2
1	A	370	ASN	3.2
1	A	371	PHE	3.1
1	B	396	TYR	3.1
1	A	368	LEU	3.1
1	A	480	CYS	2.9
1	A	394	ASN	2.9
2	C	119	SER	2.9
1	B	390	LEU	2.8
1	A	357	ARG	2.7
1	A	340	GLU	2.6
1	B	334	ASN	2.5
1	B	391	CYS	2.5
1	A	339	ASP	2.5
1	B	526	GLY	2.4
1	A	367	VAL	2.4
1	B	473	TYR	2.4
1	B	475	ALA	2.4
1	A	514	SER	2.3
1	A	519	HIS	2.3
1	B	523	THR	2.3
1	B	373	PRO	2.3
1	B	388	ASN	2.3
1	A	337	PRO	2.3
1	A	483	VAL	2.3
1	A	388	ASN	2.3
1	B	478	LYS	2.2
1	B	525	CYS	2.2
1	B	479	PRO	2.1
1	B	487	ASN	2.1
1	A	375	PHE	2.1
1	B	367	VAL	2.0
1	B	335	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.