



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2023 – 06:02 PM EST

PDB ID : 1GRA  
Title : SUBSTRATE BINDING AND CATALYSIS BY GLUTATHIONE REDUCTASE AS DERIVED FROM REFINED ENZYME: SUBSTRATE CRYSTAL STRUCTURES AT 2 ANGSTROMS RESOLUTION  
Authors : Karplus, P.A.; Schulz, G.E.  
Deposited on : 1992-12-15  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.32.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.32.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTATHIONE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	461	3499	2212	603	660	24	0	0	0

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	27	10	5	10	2	0	0

- Molecule 4 is GLUTATHIONE (three-letter code: GSH) (formula:  $C_{10}H_{17}N_3O_6S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	20	10	3	6	1	0	0
4	A	1	20	10	3	6	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	530	Total 530	O 530	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: GLUTATHIONE REDUCTASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	B 1 1 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	119.80Å 84.50Å 63.20Å 90.00° 90.00° 58.70°	Depositor
Resolution (Å)	10.00 – 2.00 72.20 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 97.1 (72.20-1.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	TNT	Depositor
R, $R_{free}$	0.157 , (Not available) 0.147 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.1	Xtrriage
Anisotropy	0.186	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 90.0	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4149	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, FAD, GSH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.87	23/3566 (0.6%)	1.22	41/4824 (0.8%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	185	GLU	CD-OE1	6.47	1.32	1.25
1	A	358	GLU	CD-OE2	6.41	1.32	1.25
1	A	124	GLU	CD-OE1	6.34	1.32	1.25
1	A	427	GLU	CD-OE1	6.29	1.32	1.25
1	A	165	GLU	CD-OE2	6.25	1.32	1.25
1	A	236	GLU	CD-OE1	6.24	1.32	1.25
1	A	473	GLU	CD-OE2	6.22	1.32	1.25
1	A	262	GLU	CD-OE1	6.18	1.32	1.25
1	A	77	GLU	CD-OE1	6.17	1.32	1.25
1	A	394	GLU	CD-OE1	6.09	1.32	1.25
1	A	253	GLU	CD-OE2	6.06	1.32	1.25
1	A	239	GLU	CD-OE1	5.81	1.32	1.25
1	A	472	GLU	CD-OE2	5.78	1.32	1.25
1	A	317	GLU	CD-OE1	5.68	1.31	1.25
1	A	91	GLU	CD-OE2	5.63	1.31	1.25
1	A	101	GLU	CD-OE2	5.57	1.31	1.25
1	A	141	GLU	CD-OE2	5.54	1.31	1.25
1	A	384	GLU	CD-OE1	5.48	1.31	1.25
1	A	41	GLU	CD-OE2	5.41	1.31	1.25
1	A	428	GLU	CD-OE1	5.34	1.31	1.25
1	A	386	GLU	CD-OE1	5.30	1.31	1.25
1	A	184	GLU	CD-OE2	5.24	1.31	1.25
1	A	237	GLU	CD-OE1	5.23	1.31	1.25

All (41) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	GLU	C-N-CA	-10.78	99.66	122.30
1	A	461	ASP	CB-CG-OD2	-8.90	110.29	118.30
1	A	309	ASP	CB-CG-OD2	-7.85	111.23	118.30
1	A	104	ASP	CB-CG-OD2	-7.65	111.42	118.30
1	A	104	ASP	CB-CG-OD1	7.36	124.92	118.30
1	A	283	ASP	CB-CG-OD1	-7.26	111.77	118.30
1	A	297	ASP	CB-CG-OD2	-7.21	111.81	118.30
1	A	227	ASP	CB-CG-OD2	6.85	124.46	118.30
1	A	308	ASP	CB-CG-OD2	-6.82	112.16	118.30
1	A	308	ASP	CB-CG-OD1	6.80	124.42	118.30
1	A	135	ASP	CB-CG-OD1	-6.73	112.24	118.30
1	A	461	ASP	CB-CG-OD1	6.73	124.36	118.30
1	A	331	ASP	CB-CG-OD1	-6.58	112.38	118.30
1	A	22	ASP	CB-CG-OD1	-6.47	112.48	118.30
1	A	84	ASP	CB-CG-OD1	-6.45	112.49	118.30
1	A	441	ASP	CB-CG-OD1	6.36	124.03	118.30
1	A	385	ASP	CB-CG-OD1	-6.30	112.63	118.30
1	A	178	ASP	CB-CG-OD2	-6.27	112.66	118.30
1	A	159	MET	CB-CA-C	-6.24	97.91	110.40
1	A	283	ASP	CB-CG-OD2	6.18	123.87	118.30
1	A	385	ASP	CB-CG-OD2	6.07	123.76	118.30
1	A	359	ASP	CB-CG-OD1	6.05	123.74	118.30
1	A	331	ASP	CB-CG-OD2	5.92	123.63	118.30
1	A	224	ARG	NE-CZ-NH2	5.89	123.25	120.30
1	A	359	ASP	CB-CG-OD2	-5.78	113.10	118.30
1	A	297	ASP	CB-CG-OD1	5.75	123.47	118.30
1	A	97	ARG	NE-CZ-NH2	5.67	123.13	120.30
1	A	178	ASP	CB-CG-OD1	5.62	123.36	118.30
1	A	363	ASP	CB-CG-OD2	-5.55	113.30	118.30
1	A	91	GLU	CA-C-N	5.48	127.16	116.20
1	A	231	SER	N-CA-CB	5.44	118.66	110.50
1	A	92	GLY	N-CA-C	5.39	126.59	113.10
1	A	441	ASP	CB-CG-OD2	-5.39	113.45	118.30
1	A	281	ASP	CB-CG-OD1	-5.25	113.57	118.30
1	A	459	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	A	459	ASP	CB-CG-OD1	5.17	122.95	118.30
1	A	135	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	316	ASP	CB-CG-OD2	5.11	122.90	118.30
1	A	291	ARG	NE-CZ-NH2	5.08	122.84	120.30
1	A	227	ASP	CB-CG-OD1	-5.08	113.73	118.30
1	A	81	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3537	45	1
2	A	53	0	31	1	0
3	A	27	0	11	0	0
4	A	40	0	28	4	0
5	A	530	0	0	8	2
All	All	4149	0	3607	47	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ILE:HG21	4:A:482:GSH:HA32	1.51	0.90
1:A:296:LYS:HB2	5:A:583:HOH:O	1.83	0.78
1:A:293:PRO:HG3	1:A:312:HIS:CD2	2.23	0.73
1:A:185:GLU:HG3	5:A:790:HOH:O	1.88	0.72
1:A:113:ILE:HG21	4:A:482:GSH:CA3	2.22	0.69
1:A:77:GLU:OE2	1:A:374:HIS:ND1	2.26	0.68
1:A:93:LYS:O	1:A:93:LYS:HG2	1.94	0.67
1:A:373:SER:HB2	5:A:961:HOH:O	1.95	0.67
1:A:114:TYR:OH	4:A:481:GSH:HB23	1.95	0.66
1:A:164:HIS:HB2	5:A:899:HOH:O	2.08	0.54
1:A:115:GLN:O	1:A:119:THR:HG23	2.07	0.54
1:A:198:ILE:O	1:A:202:MET:HG2	2.07	0.54
1:A:224:ARG:HD2	5:A:954:HOH:O	2.06	0.54
1:A:266:VAL:HG22	1:A:276:MET:HG2	1.90	0.53
1:A:78:PHE:HB3	5:A:944:HOH:O	2.09	0.53
1:A:388:ILE:HG23	1:A:393:ILE:HG12	1.91	0.53
1:A:292:VAL:HG23	1:A:293:PRO:HD2	1.91	0.52
1:A:167:GLN:HG2	5:A:643:HOH:O	2.11	0.51
1:A:315:VAL:HA	1:A:320:ASN:O	2.11	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:MET:HB3	1:A:160:PRO:CD	2.43	0.49
1:A:202:MET:HA	1:A:202:MET:CE	2.44	0.48
1:A:426:LYS:HB3	1:A:426:LYS:HE3	1.63	0.48
1:A:65:PRO:HB2	1:A:181:PHE:CE1	2.49	0.47
1:A:339:THR:HB	1:A:340:PRO:HD3	1.97	0.47
4:A:481:GSH:HG12	5:A:984:HOH:O	2.14	0.46
1:A:232:THR:O	1:A:236:GLU:HG3	2.16	0.46
1:A:91:GLU:O	1:A:91:GLU:HG3	2.12	0.46
1:A:403:PHE:CE1	1:A:473:GLU:HG3	2.51	0.45
1:A:45:ARG:NH2	1:A:124:GLU:OE1	2.49	0.45
1:A:334:GLY:HA2	1:A:337:LEU:HD21	2.00	0.44
1:A:159:MET:HB3	1:A:160:PRO:HD2	2.00	0.44
1:A:421:MET:SD	1:A:457:LYS:HD3	2.57	0.43
1:A:371:VAL:HB	1:A:377:ILE:HB	2.01	0.43
1:A:202:MET:HA	1:A:202:MET:HE2	2.00	0.43
1:A:91:GLU:HG2	1:A:92:GLY:N	2.33	0.42
1:A:138:PRO:HB3	1:A:326:ILE:HD11	2.02	0.42
1:A:411:THR:O	1:A:414:LYS:HE2	2.19	0.42
1:A:100:LYS:O	1:A:100:LYS:HG2	2.18	0.42
1:A:339:THR:CB	1:A:340:PRO:HD3	2.50	0.41
1:A:153:LEU:HD11	1:A:329:VAL:HG13	2.02	0.41
1:A:222:VAL:O	1:A:231:SER:HB2	2.21	0.41
1:A:23:TYR:O	1:A:46:ALA:HA	2.20	0.41
1:A:192:ILE:HD13	1:A:203:ALA:HA	2.02	0.41
2:A:479:FAD:H9	2:A:479:FAD:HI'1	1.86	0.41
1:A:331:ASP:HA	1:A:337:LEU:HD22	2.03	0.40
1:A:262:GLU:HA	1:A:279:ILE:O	2.21	0.40
1:A:258:LEU:HD23	1:A:258:LEU:HA	1.89	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:579:HOH:O	5:A:579:HOH:O[2_665]	0.47	1.73
1:A:90:CYS:SG	1:A:90:CYS:SG[2_665]	1.29	0.91
5:A:593:HOH:O	5:A:641:HOH:O[3_554]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	459/478 (96%)	441 (96%)	17 (4%)	1 (0%)	47 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	382/393 (97%)	363 (95%)	19 (5%)	24 20

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	CYS
1	A	75	HIS
1	A	89	SER
1	A	90	CYS
1	A	91	GLU
1	A	93	LYS
1	A	134	SER
1	A	167	GLN
1	A	185	GLU
1	A	212	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	252	LYS
1	A	253	GLU
1	A	256	LYS
1	A	276	MET
1	A	296	LYS
1	A	317	GLU
1	A	339	THR
1	A	397	LYS
1	A	426	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	HIS
1	A	117	ASN
1	A	182	GLN
1	A	366	ASN
1	A	425	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GSH	A	482	4	18,19,19	0.99	0	23,24,24	1.76	3 (13%)
2	FAD	A	479	-	53,58,58	1.08	3 (5%)	68,89,89	1.15	8 (11%)
3	NDP	A	480	-	25,29,52	1.62	5 (20%)	31,45,80	1.51	7 (22%)
4	GSH	A	481	4	18,19,19	0.78	0	23,24,24	1.13	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GSH	A	482	4	-	5/24/24/24	-
2	FAD	A	479	-	-	5/30/50/50	0/6/6/6
3	NDP	A	480	-	-	1/11/31/77	0/3/3/5
4	GSH	A	481	4	-	0/24/24/24	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	480	NDP	O4B-C1B	5.02	1.48	1.41
2	A	479	FAD	O4B-C1B	4.18	1.46	1.41
3	A	480	NDP	P2B-O2B	3.10	1.65	1.59
3	A	480	NDP	PA-O2A	-2.49	1.45	1.54
2	A	479	FAD	C2A-N1A	2.32	1.38	1.33
2	A	479	FAD	O2-C2	-2.14	1.20	1.24
3	A	480	NDP	C2A-N1A	2.13	1.37	1.33
3	A	480	NDP	PA-O3	2.12	1.63	1.54

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	482	GSH	CA2-CB2-SG2	-6.09	107.35	114.19
3	A	480	NDP	O2A-PA-O1A	3.42	124.08	110.68
2	A	479	FAD	O4B-C1B-C2B	-3.26	102.16	106.93
4	A	481	GSH	CB1-CG1-CD1	-3.11	106.10	113.04
2	A	479	FAD	C5A-C6A-N1A	-2.73	114.17	120.35
3	A	480	NDP	C2A-N1A-C6A	2.68	123.34	118.75

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	479	FAD	O2A-PA-O1A	2.56	124.91	112.24
3	A	480	NDP	C5A-C6A-N1A	-2.49	114.70	120.35
2	A	479	FAD	C5A-C6A-N6A	2.47	124.11	120.35
3	A	480	NDP	C4A-C5A-N7A	2.46	111.96	109.40
3	A	480	NDP	C5A-C6A-N6A	2.42	124.03	120.35
3	A	480	NDP	PA-O5B-C5B	2.41	124.95	118.30
2	A	479	FAD	C4A-C5A-N7A	2.41	111.91	109.40
3	A	480	NDP	O3-PA-O5B	-2.31	100.59	106.73
4	A	482	GSH	CG1-CB1-CA1	-2.17	108.78	113.84
2	A	479	FAD	C5'-C4'-C3'	-2.14	108.07	112.20
2	A	479	FAD	C2A-N1A-C6A	2.10	122.35	118.75
4	A	482	GSH	OE1-CD1-N2	2.02	126.36	122.95
2	A	479	FAD	O2P-P-O1P	2.00	122.15	112.24

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	482	GSH	O31-C3-CA3-N3
4	A	482	GSH	CA1-CB1-CG1-CD1
4	A	482	GSH	O32-C3-CA3-N3
2	A	479	FAD	PA-O3P-P-O5'
2	A	479	FAD	O4B-C4B-C5B-O5B
2	A	479	FAD	P-O3P-PA-O1A
4	A	482	GSH	N3-C2-CA2-N2
4	A	482	GSH	O2-C2-CA2-N2
2	A	479	FAD	P-O3P-PA-O2A
3	A	480	NDP	O4B-C4B-C5B-O5B
2	A	479	FAD	C3B-C4B-C5B-O5B



There are no ring outliers.

3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	482	GSH	2	0
2	A	479	FAD	1	0
4	A	481	GSH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	461/478 (96%)	-0.93	1 (0%) <b>95</b> <b>94</b>	8, 18, 41, 65	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	91	GLU	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GSH	A	482	20/20	0.90	0.14	13,26,56,63	0
3	NDP	A	480	27/48	0.96	0.10	17,26,38,50	0
4	GSH	A	481	20/20	0.97	0.10	17,24,32,33	0
2	FAD	A	479	53/53	0.99	0.06	5,11,18,21	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers


as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around NDP A 480:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.