

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 10, 2021 – 07:04 PM EDT

PDB ID	:	3GSQ
Title	:	Crystal structure of the binary complex between HLA-A2 and HCMV NLV-
		M5S peptide variant
Authors	:	Reiser, JB.; Saulquin, X.; Gras, S.; Debeaupuis, E.; Echasserieau, K.;
		Kissenpfennig, A.; Legoux, F.; Chouquet, A.; Le Gorrec, M.; Machillot, P.;
		Neveu, B.; Thielens, N.; Malissen, B.; Bonneville, M.; Housset, D.
Deposited on	:	2009-03-27
Resolution	:	2.12  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	6778 (2.14-2.10)
Ramachandran outliers	138981	6705 (2.14-2.10)
Sidechain outliers	138945	6706 (2.14-2.10)
RSRZ outliers	127900	6112 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	274	2% 91% 8%	
2	В	100	5% 90% 10%	6
3	Р	9	100%	



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called HLA class I histocompatibility antigen, A-2 alpha chain.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	274	Total 2269	C 1421	N 410	0 429	S 9	0	5	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	245	VAL	ALA	engineered mutation	UNP P01892

• Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	100	Total 870	$\begin{array}{c} \mathrm{C} \\ 555 \end{array}$	N 145	O 166	$\frac{S}{4}$	3	5	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	0	MET	-	initiating methionine	UNP P61769

• Molecule 3 is a protein called HCMV pp65 fragment 495-503, variant M5S (NLVPSVATV).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
3	Р	9	Total 63	C 40	N 10	0 13	0	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	111	Total O 111 111	0	0
4	В	50	Total         O           50         50	0	0

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Р	7	Total O 7 7	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HLA class I histocompatibility antigen, A-2 alpha chain



Chain P: 100%

There are no outlier residues recorded for this chain.



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	54.05Å 80.90Å 57.66Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $114.17^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	14.92 - 2.12	Depositor
Resolution (A)	14.92 - 2.12	EDS
% Data completeness	95.9 (14.92-2.12)	Depositor
(in resolution range)	95.9(14.92-2.12)	EDS
R <sub>merge</sub>	0.04	Depositor
R <sub>sym</sub>	0.04	Depositor
$< I/\sigma(I) > 1$	$4.38 (at 2.12 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
D D.	0.196 , $0.246$	Depositor
$R, R_{free}$	0.203 , (Not available)	DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	24.1	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , $40.0$	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3370	wwPDB-VP
Average B, all atoms $(Å^2)$	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.37% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain Bor		nd lengths	Bo	ond angles
	Ullalli	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.55	0/2349	0.62	1/3187~(0.0%)
2	В	1.03	2/902~(0.2%)	0.74	2/1218~(0.2%)
3	Р	0.68	0/63	0.71	0/86
All	All	0.72	2/3314~(0.1%)	0.66	3/4491~(0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	В	58[A]	LYS	CG-CD	18.48	2.15	1.52
2	В	58[B]	LYS	CG-CD	18.48	2.15	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
2	В	58[A]	LYS	CB-CG-CD	-11.24	82.37	111.60
2	В	58[B]	LYS	CB-CG-CD	-11.24	82.37	111.60
1	А	270	LEU	CA-CB-CG	5.74	128.49	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2269	0	2133	12	0
2	В	870	0	844	7	0

Continued on next page...



001000	continued from proceed as page							
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes		
3	Р	63	0	70	0	0		
4	А	111	0	0	3	0		
4	В	50	0	0	0	0		
4	Р	7	0	0	0	0		
All	All	3370	0	3047	16	0		

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 16 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:74:GLU:OE2	2:B:75[B]:LYS:NZ	1.72	1.21
2:B:1:ILE:CD1	2:B:3:ARG:HH21	2.04	0.70
1:A:49:ALA:O	1:A:52:ILE:HG22	2.06	0.55
1:A:146:LYS:HE2	4:A:381:HOH:O	2.07	0.55
1:A:127:LYS:HD2	1:A:132:SER:OG	2.07	0.55

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	277/274~(101%)	273~(99%)	4 (1%)	0	100	100
2	В	103/100~(103%)	103 (100%)	0	0	100	100
3	Р	7/9~(78%)	7 (100%)	0	0	100	100
All	All	387/383~(101%)	383~(99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	236/231~(102%)	228~(97%)	8 (3%)	37 38
2	В	100/95~(105%)	97~(97%)	3(3%)	41 43
3	Р	8/8 (100%)	8 (100%)	0	100 100
All	All	344/334~(103%)	333~(97%)	11 (3%)	41 40

5 of 11 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	270	LEU
2	В	4[A]	THR
2	В	70	PHE
2	В	4[B]	THR
1	А	197	HIS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	А	260	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



### 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	274/274~(100%)	0.17	6 (2%) 62 66	25, 34, 47, 58	13 (4%)
2	В	100/100~(100%)	0.34	5 (5%) 28 34	26, 36, 49, 59	6 (6%)
3	Р	9/9~(100%)	-0.28	0 100 100	30, 31, 35, 36	0
All	All	383/383~(100%)	0.20	11 (2%) 51 57	25, 34, 48, 59	19 (4%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	194	VAL	4.4
1	А	196	ASP	3.4
2	В	98	ASP	3.4
1	А	195	SER	3.3
2	В	0	MET	3.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.



## 6.5 Other polymers (i)

There are no such residues in this entry.

