



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 4, 2023 – 11:13 pm GMT

PDB ID : 1GTF
Title : The structure of the trp RNA-binding attenuation protein (TRAP) bound to a 53-nucleotide RNA molecule containing GAGUU repeats
Authors : Hopcroft, N.H.; Wendt, A.L.; Gollnick, P.; Antson, A.A.
Deposited on : 2002-01-15
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)
RNA backbone	3102	1079 (2.40-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	74	 5% 85% 7% • 7%
1	B	74	 80% 12% 8%
1	C	74	 12% 80% 12% • 5%
1	D	74	 11% 77% 15% • 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	74	% 81% 9% • 8%
1	F	74	% 81% 7% 5% 7%
1	G	74	3% 77% 16% • 5%
1	H	74	86% 5% 8%
1	I	74	3% 85% 8% 7%
1	J	74	82% 7% • 9%
1	K	74	11% 80% 12% • 7%
1	L	74	4% 84% 8% • 5%
1	M	74	3% 82% 11% • •
1	N	74	3% 84% 11% 5%
1	O	74	% 84% 8% • 5%
1	P	74	% 80% 15% 5%
1	Q	74	3% 84% 9% • 5%
1	R	74	4% 77% 15% • 5%
1	S	74	% 81% 14% 5%
1	T	74	% 80% 15% 5%
1	U	74	% 77% 16% • 5%
1	V	74	3% 85% 8% • 5%
2	W	55	16% 15% 38% 25% • 20%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 14607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRP RNA-BINDING ATTENUATION PROTEIN (TRAP).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	69	Total	C	N	O	15	0	0
			536	336	98	102			
1	B	68	Total	C	N	O	16	0	0
			527	330	96	101			
1	C	70	Total	C	N	O	13	0	0
			544	340	100	104			
1	D	69	Total	C	N	O	26	0	0
			536	336	98	102			
1	E	68	Total	C	N	O	21	0	0
			527	330	96	101			
1	F	69	Total	C	N	O	30	0	0
			536	336	98	102			
1	G	70	Total	C	N	O	26	0	0
			544	340	100	104			
1	H	68	Total	C	N	O	16	0	0
			527	330	96	101			
1	I	69	Total	C	N	O	23	0	0
			536	336	98	102			
1	J	67	Total	C	N	O	13	0	0
			523	328	95	100			
1	K	69	Total	C	N	O	21	0	0
			536	336	98	102			
1	L	70	Total	C	N	O	13	0	0
			542	338	99	105			
1	M	71	Total	C	N	O	14	0	0
			551	344	101	106			
1	N	70	Total	C	N	O	4	0	0
			542	338	99	105			
1	O	70	Total	C	N	O	0	0	0
			542	338	99	105			
1	P	70	Total	C	N	O	6	0	0
			542	338	99	105			

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	Q	70	Total	C	N	O	3	0	0
			542	338	99	105			
1	R	70	Total	C	N	O	17	0	0
			542	338	99	105			
1	S	70	Total	C	N	O	8	0	0
			542	338	99	105			
1	T	70	Total	C	N	O	7	0	0
			542	338	99	105			
1	U	70	Total	C	N	O	10	0	0
			542	338	99	105			
1	V	70	Total	C	N	O	7	0	0
			542	338	99	105			

- Molecule 2 is a RNA chain called (GAGUU)10GAG 53-NUCLEOTIDE RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	W	44	Total	C	N	O	P	0	0	0
			968	429	187	308	44			

- Molecule 3 is TRYPTOPHAN (three-letter code: TRP) (formula: C₁₁H₁₂N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	11	2	2		
3	B	1	Total	C	N	O	0	0
			15	11	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			15	11	2	2		
3	D	1	Total	C	N	O	0	0
			15	11	2	2		
3	E	1	Total	C	N	O	0	0
			15	11	2	2		
3	F	1	Total	C	N	O	0	0
			15	11	2	2		
3	G	1	Total	C	N	O	0	0
			15	11	2	2		
3	H	1	Total	C	N	O	0	0
			15	11	2	2		
3	I	1	Total	C	N	O	0	0
			15	11	2	2		
3	J	1	Total	C	N	O	0	0
			15	11	2	2		
3	K	1	Total	C	N	O	0	0
			15	11	2	2		
3	L	1	Total	C	N	O	0	0
			15	11	2	2		
3	M	1	Total	C	N	O	0	0
			15	11	2	2		
3	N	1	Total	C	N	O	0	0
			15	11	2	2		
3	O	1	Total	C	N	O	0	0
			15	11	2	2		
3	P	1	Total	C	N	O	0	0
			15	11	2	2		
3	Q	1	Total	C	N	O	0	0
			15	11	2	2		
3	R	1	Total	C	N	O	0	0
			15	11	2	2		
3	S	1	Total	C	N	O	0	0
			15	11	2	2		
3	T	1	Total	C	N	O	0	0
			15	11	2	2		
3	U	1	Total	C	N	O	0	0
			15	11	2	2		
3	V	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	94	Total O 94 94	0	0
4	B	69	Total O 69 69	0	0
4	C	89	Total O 89 89	0	0
4	D	70	Total O 70 70	0	0
4	E	63	Total O 63 63	0	0
4	F	60	Total O 60 60	0	0
4	G	64	Total O 64 64	0	0
4	H	62	Total O 62 62	0	0
4	I	68	Total O 68 68	0	0
4	J	71	Total O 71 71	0	0
4	K	82	Total O 82 82	0	0
4	L	44	Total O 44 44	0	0
4	M	51	Total O 51 51	0	0
4	N	46	Total O 46 46	0	0
4	O	49	Total O 49 49	0	0
4	P	54	Total O 54 54	0	0
4	Q	59	Total O 59 59	0	0
4	R	75	Total O 75 75	0	0
4	S	51	Total O 51 51	0	0
4	T	63	Total O 63 63	0	0
4	U	51	Total O 51 51	0	0
4	V	54	Total O 54 54	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	W	77	Total	O	0	0
			77	77		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)





- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)









- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)




- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)



- Molecule 1: TRP RNA-BINDING ATTENUATION PROTEIN (TRAP)



- Molecule 2: (GAGUU)10GAG 53-NUCLEOTIDE RNA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.08Å 111.49Å 138.23Å 90.00° 117.28° 90.00°	Depositor
Resolution (Å)	47.67 – 1.75 47.93 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.67-1.75) 95.9 (47.93-1.75)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 1.75Å)	Xtrriage
Refinement program	REFMAC 5.1.07	Depositor
R, R_{free}	0.194 , 0.242 0.234 , 0.259	Depositor DCC
R_{free} test set	1848 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtrriage
Anisotropy	0.281	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14607	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.16	3/543 (0.6%)	1.07	3/728 (0.4%)
1	B	1.15	4/534 (0.7%)	1.10	4/717 (0.6%)
1	C	1.09	3/551 (0.5%)	1.29	4/739 (0.5%)
1	D	1.00	2/543 (0.4%)	1.08	4/728 (0.5%)
1	E	1.19	3/534 (0.6%)	0.98	2/717 (0.3%)
1	F	1.32	5/543 (0.9%)	1.17	7/728 (1.0%)
1	G	1.03	2/551 (0.4%)	1.05	5/739 (0.7%)
1	H	1.02	1/534 (0.2%)	0.97	1/717 (0.1%)
1	I	0.95	1/543 (0.2%)	1.12	7/728 (1.0%)
1	J	1.32	2/530 (0.4%)	0.98	4/712 (0.6%)
1	K	1.15	3/543 (0.6%)	1.03	4/728 (0.5%)
1	L	0.74	0/549	0.90	1/738 (0.1%)
1	M	0.93	2/558 (0.4%)	0.91	1/749 (0.1%)
1	N	0.82	1/549 (0.2%)	0.84	0/738
1	O	0.76	0/549	0.96	0/738
1	P	0.82	0/549	0.90	0/738
1	Q	0.82	0/549	0.93	2/738 (0.3%)
1	R	0.88	1/549 (0.2%)	1.02	2/738 (0.3%)
1	S	0.85	1/549 (0.2%)	0.96	0/738
1	T	0.82	1/549 (0.2%)	0.93	0/738
1	U	0.84	1/549 (0.2%)	0.90	1/738 (0.1%)
1	V	0.75	1/549 (0.2%)	0.85	0/738
2	W	2.61	11/1078 (1.0%)	1.90	35/1661 (2.1%)
All	All	1.21	48/13075 (0.4%)	1.12	87/17771 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

The worst 5 of 48 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	W	153	G	O3'-P	30.19	1.97	1.61
2	W	103	G	O3'-P	27.15	1.93	1.61
2	W	108	G	O3'-P	26.71	1.93	1.61
2	W	118	G	O3'-P	25.44	1.91	1.61
2	W	123	G	O3'-P	23.34	1.89	1.61

The worst 5 of 87 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	66	ARG	NE-CZ-NH2	15.20	127.90	120.30
1	F	66	ARG	CD-NE-CZ	-14.46	103.35	123.60
2	W	118	G	O4'-C1'-N9	12.22	117.98	108.20
2	W	126	G	P-O3'-C3'	11.70	133.74	119.70
1	C	66	ARG	NE-CZ-NH1	-11.62	114.49	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	29	ASP	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	536	0	541	2	0
1	B	527	0	528	2	0
1	C	544	0	547	7	0
1	D	536	0	541	11	0
1	E	527	0	528	4	0
1	F	536	0	541	3	0
1	G	544	0	547	11	0
1	H	527	0	528	1	0
1	I	536	0	541	1	0
1	J	523	0	525	3	0
1	K	536	0	541	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	542	0	541	8	0
1	M	551	0	554	9	0
1	N	542	0	541	4	0
1	O	542	0	541	8	0
1	P	542	0	541	19	0
1	Q	542	0	541	14	0
1	R	542	0	541	9	0
1	S	542	0	541	6	0
1	T	542	0	541	5	0
1	U	542	0	541	7	0
1	V	542	0	541	6	0
2	W	968	0	484	26	0
3	A	15	0	9	1	0
3	B	15	0	9	1	0
3	C	15	0	9	1	0
3	D	15	0	9	1	0
3	E	15	0	9	1	0
3	F	15	0	9	1	0
3	G	15	0	9	1	0
3	H	15	0	9	1	0
3	I	15	0	9	1	0
3	J	15	0	9	1	0
3	K	15	0	9	1	0
3	L	15	0	9	1	0
3	M	15	0	9	1	0
3	N	15	0	9	1	0
3	O	15	0	9	1	0
3	P	15	0	9	1	0
3	Q	15	0	9	1	0
3	R	15	0	9	1	0
3	S	15	0	9	1	0
3	T	15	0	9	1	0
3	U	15	0	9	1	0
3	V	15	0	9	1	0
4	A	94	0	0	1	0
4	B	69	0	0	0	0
4	C	89	0	0	3	1
4	D	70	0	0	6	0
4	E	63	0	0	3	0
4	F	60	0	0	1	0
4	G	64	0	0	2	0
4	H	62	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	I	68	0	0	0	0
4	J	71	0	0	0	0
4	K	82	0	0	1	1
4	L	44	0	0	3	0
4	M	51	0	0	4	0
4	N	46	0	0	1	0
4	O	49	0	0	2	1
4	P	54	0	0	2	0
4	Q	59	0	0	2	0
4	R	75	0	0	2	0
4	S	51	0	0	1	1
4	T	63	0	0	0	0
4	U	51	0	0	2	0
4	V	54	0	0	3	0
4	W	77	0	0	4	0
All	All	14607	0	12554	138	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:W:153:G:O3'	2:W:154:U:P	1.97	1.23
1:P:66:ARG:CZ	1:Q:66:ARG:HH21	1.62	1.12
1:R:8:ASP:OD1	4:R:2007:HOH:O	1.72	1.07
1:P:66:ARG:NH2	1:Q:66:ARG:HH21	1.53	1.04
1:P:66:ARG:HH12	1:Q:66:ARG:NH2	1.58	1.01

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:2047:HOH:O	4:K:2061:HOH:O[4_555]	1.95	0.25
4:O:2005:HOH:O	4:S:2049:HOH:O[2_656]	1.98	0.22

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/74 (90%)	67 (100%)	0	0	100	100
1	B	66/74 (89%)	66 (100%)	0	0	100	100
1	C	68/74 (92%)	68 (100%)	0	0	100	100
1	D	67/74 (90%)	67 (100%)	0	0	100	100
1	E	66/74 (89%)	66 (100%)	0	0	100	100
1	F	67/74 (90%)	67 (100%)	0	0	100	100
1	G	68/74 (92%)	68 (100%)	0	0	100	100
1	H	66/74 (89%)	66 (100%)	0	0	100	100
1	I	67/74 (90%)	67 (100%)	0	0	100	100
1	J	65/74 (88%)	65 (100%)	0	0	100	100
1	K	67/74 (90%)	67 (100%)	0	0	100	100
1	L	68/74 (92%)	68 (100%)	0	0	100	100
1	M	69/74 (93%)	69 (100%)	0	0	100	100
1	N	68/74 (92%)	68 (100%)	0	0	100	100
1	O	68/74 (92%)	68 (100%)	0	0	100	100
1	P	68/74 (92%)	68 (100%)	0	0	100	100
1	Q	68/74 (92%)	68 (100%)	0	0	100	100
1	R	68/74 (92%)	68 (100%)	0	0	100	100
1	S	68/74 (92%)	68 (100%)	0	0	100	100
1	T	68/74 (92%)	68 (100%)	0	0	100	100
1	U	68/74 (92%)	68 (100%)	0	0	100	100
1	V	68/74 (92%)	68 (100%)	0	0	100	100
All	All	1483/1628 (91%)	1483 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	57/62 (92%)	55 (96%)	2 (4%)	36 13
1	B	56/62 (90%)	54 (96%)	2 (4%)	35 13
1	C	58/62 (94%)	52 (90%)	6 (10%)	7 1
1	D	57/62 (92%)	54 (95%)	3 (5%)	22 5
1	E	56/62 (90%)	54 (96%)	2 (4%)	35 13
1	F	57/62 (92%)	53 (93%)	4 (7%)	15 2
1	G	58/62 (94%)	57 (98%)	1 (2%)	60 42
1	H	56/62 (90%)	55 (98%)	1 (2%)	59 40
1	I	57/62 (92%)	57 (100%)	0	100 100
1	J	56/62 (90%)	53 (95%)	3 (5%)	22 5
1	K	57/62 (92%)	54 (95%)	3 (5%)	22 5
1	L	58/62 (94%)	56 (97%)	2 (3%)	37 14
1	M	59/62 (95%)	56 (95%)	3 (5%)	24 6
1	N	58/62 (94%)	56 (97%)	2 (3%)	37 14
1	O	58/62 (94%)	55 (95%)	3 (5%)	23 6
1	P	58/62 (94%)	56 (97%)	2 (3%)	37 14
1	Q	58/62 (94%)	55 (95%)	3 (5%)	23 6
1	R	58/62 (94%)	55 (95%)	3 (5%)	23 6
1	S	58/62 (94%)	57 (98%)	1 (2%)	60 42
1	T	58/62 (94%)	56 (97%)	2 (3%)	37 14
1	U	58/62 (94%)	55 (95%)	3 (5%)	23 6
1	V	58/62 (94%)	56 (97%)	2 (3%)	37 14
All	All	1264/1364 (93%)	1211 (96%)	53 (4%)	30 9

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	35	SER
1	O	8	ASP
1	U	8	ASP
1	M	6	ASN
1	N	8	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	Q	6	ASN
1	P	6	ASN
1	O	6	ASN
1	L	6	ASN
1	O	33	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	W	34/55 (61%)	7 (20%)	1 (2%)

5 of 7 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	W	102	A
2	W	107	A
2	W	112	A
2	W	122	A
2	W	127	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	W	126	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TRP	J	81	-	14,16,16	1.04	0	16,22,22	1.28	2 (12%)
3	TRP	T	81	-	14,16,16	1.15	1 (7%)	16,22,22	0.85	0
3	TRP	K	81	-	14,16,16	1.08	1 (7%)	16,22,22	0.99	1 (6%)
3	TRP	H	81	-	14,16,16	1.18	1 (7%)	16,22,22	0.76	0
3	TRP	V	81	-	14,16,16	1.05	1 (7%)	16,22,22	1.30	2 (12%)
3	TRP	A	81	-	14,16,16	0.94	0	16,22,22	1.10	2 (12%)
3	TRP	L	81	-	14,16,16	0.83	0	16,22,22	1.22	2 (12%)
3	TRP	G	81	-	14,16,16	0.99	0	16,22,22	1.02	0
3	TRP	P	81	-	14,16,16	1.02	1 (7%)	16,22,22	1.24	2 (12%)
3	TRP	U	81	-	14,16,16	0.97	1 (7%)	16,22,22	1.18	2 (12%)
3	TRP	D	81	-	14,16,16	1.01	1 (7%)	16,22,22	0.83	0
3	TRP	N	81	-	14,16,16	1.02	2 (14%)	16,22,22	1.08	1 (6%)
3	TRP	M	81	-	14,16,16	0.91	0	16,22,22	1.35	2 (12%)
3	TRP	B	81	-	14,16,16	1.14	1 (7%)	16,22,22	1.16	2 (12%)
3	TRP	R	81	-	14,16,16	1.08	2 (14%)	16,22,22	1.04	2 (12%)
3	TRP	Q	81	-	14,16,16	1.13	2 (14%)	16,22,22	1.01	0
3	TRP	I	81	-	14,16,16	1.03	1 (7%)	16,22,22	0.78	0
3	TRP	E	81	-	14,16,16	0.96	0	16,22,22	1.09	1 (6%)
3	TRP	F	81	-	14,16,16	1.07	1 (7%)	16,22,22	1.01	0
3	TRP	C	81	-	14,16,16	0.96	0	16,22,22	1.18	3 (18%)
3	TRP	S	81	-	14,16,16	0.87	0	16,22,22	1.25	3 (18%)
3	TRP	O	81	-	14,16,16	0.91	0	16,22,22	1.01	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRP	J	81	-	-	1/7/8/8	0/2/2/2
3	TRP	T	81	-	-	1/7/8/8	0/2/2/2
3	TRP	K	81	-	-	1/7/8/8	0/2/2/2
3	TRP	H	81	-	-	1/7/8/8	0/2/2/2
3	TRP	V	81	-	-	1/7/8/8	0/2/2/2
3	TRP	A	81	-	-	1/7/8/8	0/2/2/2
3	TRP	L	81	-	-	1/7/8/8	0/2/2/2
3	TRP	G	81	-	-	1/7/8/8	0/2/2/2
3	TRP	P	81	-	-	1/7/8/8	0/2/2/2
3	TRP	U	81	-	-	1/7/8/8	0/2/2/2
3	TRP	D	81	-	-	1/7/8/8	0/2/2/2
3	TRP	N	81	-	-	1/7/8/8	0/2/2/2
3	TRP	M	81	-	-	0/7/8/8	0/2/2/2
3	TRP	B	81	-	-	1/7/8/8	0/2/2/2
3	TRP	R	81	-	-	1/7/8/8	0/2/2/2
3	TRP	Q	81	-	-	0/7/8/8	0/2/2/2
3	TRP	I	81	-	-	1/7/8/8	0/2/2/2
3	TRP	E	81	-	-	0/7/8/8	0/2/2/2
3	TRP	F	81	-	-	0/7/8/8	0/2/2/2
3	TRP	C	81	-	-	0/7/8/8	0/2/2/2
3	TRP	S	81	-	-	1/7/8/8	0/2/2/2
3	TRP	O	81	-	-	1/7/8/8	0/2/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	81	TRP	OXT-C	-2.79	1.21	1.30
3	B	81	TRP	CZ3-CE3	2.70	1.42	1.36
3	H	81	TRP	CZ3-CE3	2.57	1.42	1.36
3	Q	81	TRP	OXT-C	-2.53	1.22	1.30
3	F	81	TRP	OXT-C	-2.48	1.22	1.30

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	81	TRP	OXT-C-CA	3.40	124.97	113.38
3	V	81	TRP	OXT-C-O	-2.88	117.56	124.09
3	L	81	TRP	OXT-C-CA	2.87	123.15	113.38
3	V	81	TRP	OXT-C-CA	2.86	123.12	113.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	81	TRP	OXT-C-CA	2.85	123.10	113.38

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	I	81	TRP	OXT-C-CA-N
3	U	81	TRP	OXT-C-CA-N
3	D	81	TRP	OXT-C-CA-N
3	T	81	TRP	OXT-C-CA-N
3	L	81	TRP	OXT-C-CA-N

There are no ring outliers.

22 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	81	TRP	1	0
3	T	81	TRP	1	0
3	K	81	TRP	1	0
3	H	81	TRP	1	0
3	V	81	TRP	1	0
3	A	81	TRP	1	0
3	L	81	TRP	1	0
3	G	81	TRP	1	0
3	P	81	TRP	1	0
3	U	81	TRP	1	0
3	D	81	TRP	1	0
3	N	81	TRP	1	0
3	M	81	TRP	1	0
3	B	81	TRP	1	0
3	R	81	TRP	1	0
3	Q	81	TRP	1	0
3	I	81	TRP	1	0
3	E	81	TRP	1	0
3	F	81	TRP	1	0
3	C	81	TRP	1	0
3	S	81	TRP	1	0
3	O	81	TRP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	W	11

The worst 5 of 11 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	W	139:U	O3'	141:G	P	6.01
1	W	129:U	O3'	131:G	P	5.77
1	W	144:U	O3'	146:G	P	5.74
1	W	104:U	O3'	106:G	P	5.72
1	W	119:U	O3'	121:G	P	5.71

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	69/74 (93%)	0.31	4 (5%) 23 28	8, 11, 20, 26	5 (7%)
1	B	68/74 (91%)	0.09	0 100 100	8, 11, 18, 21	7 (10%)
1	C	70/74 (94%)	0.77	9 (12%) 3 5	7, 11, 19, 26	5 (7%)
1	D	69/74 (93%)	0.46	8 (11%) 4 6	8, 11, 18, 29	8 (11%)
1	E	68/74 (91%)	0.05	1 (1%) 73 80	7, 11, 16, 22	8 (11%)
1	F	69/74 (93%)	-0.11	1 (1%) 75 82	8, 11, 16, 23	8 (11%)
1	G	70/74 (94%)	0.08	2 (2%) 51 57	9, 11, 20, 24	10 (14%)
1	H	68/74 (91%)	-0.17	0 100 100	9, 11, 17, 20	5 (7%)
1	I	69/74 (93%)	0.07	2 (2%) 51 57	7, 11, 17, 27	7 (10%)
1	J	67/74 (90%)	0.26	0 100 100	8, 11, 17, 21	4 (5%)
1	K	69/74 (93%)	0.60	8 (11%) 4 6	8, 11, 15, 26	6 (8%)
1	L	70/74 (94%)	0.19	3 (4%) 35 41	7, 10, 16, 23	3 (4%)
1	M	71/74 (95%)	0.17	2 (2%) 53 58	7, 11, 17, 24	5 (7%)
1	N	70/74 (94%)	0.06	2 (2%) 51 57	7, 10, 17, 23	1 (1%)
1	O	70/74 (94%)	0.12	1 (1%) 75 82	7, 11, 18, 20	0
1	P	70/74 (94%)	0.17	1 (1%) 75 82	7, 11, 18, 25	2 (2%)
1	Q	70/74 (94%)	0.18	2 (2%) 51 57	7, 11, 16, 23	1 (1%)
1	R	70/74 (94%)	0.32	3 (4%) 35 41	7, 11, 16, 19	5 (7%)
1	S	70/74 (94%)	0.11	1 (1%) 75 82	8, 11, 16, 21	2 (2%)
1	T	70/74 (94%)	0.13	1 (1%) 75 82	7, 11, 17, 23	2 (2%)
1	U	70/74 (94%)	-0.03	1 (1%) 75 82	7, 10, 17, 21	3 (4%)
1	V	70/74 (94%)	0.17	2 (2%) 51 57	7, 11, 17, 20	2 (2%)
2	W	44/55 (80%)	1.24	9 (20%) 1 1	11, 13, 24, 24	0
All	All	1571/1683 (93%)	0.21	63 (4%) 38 45	7, 11, 19, 29	99 (6%)

The worst 5 of 63 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	75	LYS	4.6
1	D	74	GLY	4.3
1	R	74	GLY	4.2
1	L	32	PHE	4.1
1	V	74	GLY	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TRP	A	81	15/15	0.93	0.11	8,9,11,12	0
3	TRP	B	81	15/15	0.93	0.08	5,9,13,13	0
3	TRP	C	81	15/15	0.94	0.10	7,9,11,12	0
3	TRP	L	81	15/15	0.94	0.07	7,9,11,12	0
3	TRP	R	81	15/15	0.94	0.10	9,10,11,11	0
3	TRP	F	81	15/15	0.95	0.08	7,9,11,12	0
3	TRP	J	81	15/15	0.95	0.09	6,9,12,12	0
3	TRP	K	81	15/15	0.95	0.12	7,10,12,12	0
3	TRP	D	81	15/15	0.95	0.10	7,9,12,13	0
3	TRP	M	81	15/15	0.95	0.06	6,8,11,12	0
3	TRP	P	81	15/15	0.95	0.10	6,8,11,11	0
3	TRP	Q	81	15/15	0.95	0.09	7,8,10,10	0
3	TRP	E	81	15/15	0.95	0.08	6,9,12,12	0
3	TRP	S	81	15/15	0.95	0.08	8,9,12,13	0
3	TRP	V	81	15/15	0.95	0.07	6,8,11,11	0
3	TRP	H	81	15/15	0.96	0.08	7,9,10,11	0
3	TRP	I	81	15/15	0.96	0.07	7,10,12,13	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	TRP	G	81	15/15	0.96	0.07	7,9,11,12	0
3	TRP	N	81	15/15	0.96	0.07	7,8,11,12	0
3	TRP	U	81	15/15	0.96	0.09	7,9,10,12	0
3	TRP	O	81	15/15	0.96	0.07	7,9,10,10	0
3	TRP	T	81	15/15	0.97	0.07	7,8,10,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.