



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 05:49 PM BST

PDB ID : 2GUP
Title : Structural Genomics, the crystal structure of a ROK family protein from *Streptococcus pneumoniae* TIGR4 in complex with sucrose
Authors : Tan, K.; Li, H.; Abdullah, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2006-05-01
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	
2	B	2	
2	C	2	
2	D	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	D	1	-	-	-	X
3	TRS	A	293	-	X	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2372 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ROK family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	289	2211	1397	372	432	4	6	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	cloning artifact	GB 14973654
A	-1	ASN	-	cloning artifact	GB 14973654
A	0	ALA	-	cloning artifact	GB 14973654
A	1	MSE	MET	modified residue	GB 14973654
A	57	MSE	MET	modified residue	GB 14973654
A	131	MSE	MET	modified residue	GB 14973654
A	151	MSE	MET	modified residue	GB 14973654
A	172	MSE	MET	modified residue	GB 14973654
A	208	MSE	MET	modified residue	GB 14973654

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	0	0
2	C	2	23	12	11	0	0	0
2	D	2	23	12	11	0	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	8	4	1	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	84	84	84	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ROK family protein



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.68Å 114.67Å 117.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.51 – 2.01 19.51 – 2.01	Depositor EDS
% Data completeness (in resolution range)	91.8 (19.51-2.01) 91.8 (19.51-2.01)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.185 , 0.228 0.189 , 0.231	Depositor DCC
R_{free} test set	1178 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	38.5	Xtrriage
Anisotropy	0.101	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2372	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, GLC, FRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.40	18/2252 (0.8%)	1.10	13/3055 (0.4%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269	TYR	CD1-CE1	8.98	1.52	1.39
1	A	69	VAL	CB-CG1	6.69	1.67	1.52
1	A	122	VAL	CB-CG1	6.47	1.66	1.52
1	A	84	TRP	CB-CG	6.21	1.61	1.50
1	A	276	TYR	CD1-CE1	6.20	1.48	1.39
1	A	178	GLU	CG-CD	6.17	1.61	1.51
1	A	14	LYS	CB-CG	6.08	1.69	1.52
1	A	276	TYR	CE1-CZ	5.87	1.46	1.38
1	A	66	GLU	CG-CD	5.82	1.60	1.51
1	A	82	PHE	CE2-CZ	5.41	1.47	1.37
1	A	273	ALA	C-O	5.37	1.33	1.23
1	A	149	GLY	N-CA	5.16	1.53	1.46
1	A	37	GLU	CG-CD	5.11	1.59	1.51
1	A	273	ALA	CA-CB	5.10	1.63	1.52
1	A	256	GLU	CG-CD	5.09	1.59	1.51
1	A	66	GLU	CB-CG	5.09	1.61	1.52
1	A	191	TYR	CB-CG	5.08	1.59	1.51
1	A	269	TYR	CD2-CE2	5.06	1.47	1.39

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	110	LEU	CB-CG-CD2	-8.19	97.08	111.00
1	A	97	LEU	CB-CG-CD2	-7.50	98.25	111.00
1	A	137	LEU	CB-CG-CD2	-7.13	98.88	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	253	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	97	LEU	CB-CG-CD1	-6.17	100.50	111.00
1	A	171	ASN	CB-CA-C	5.56	121.53	110.40
1	A	97	LEU	CA-CB-CG	5.27	127.42	115.30
1	A	253	ASP	CB-CG-OD2	-5.14	113.67	118.30
1	A	276	TYR	CZ-CE2-CD2	5.02	124.32	119.80
1	A	250	ASP	CB-CG-OD2	-5.02	113.78	118.30
1	A	139	ARG	NE-CZ-NH1	-5.01	117.80	120.30
1	A	276	TYR	CB-CG-CD1	5.00	124.00	121.00
1	A	14	LYS	CD-CE-NZ	-5.00	100.20	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2211	0	2148	27	0
2	B	23	0	21	1	0
2	C	23	0	21	1	0
2	D	23	0	21	0	0
3	A	8	0	12	0	0
4	A	84	0	0	2	0
All	All	2372	0	2223	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:PRO:CB	1:A:77:PRO:CG	1.74	1.44
1:A:227:VAL:HG23	1:A:263:VAL:HG22	1.60	0.84
1:A:259:THR:HG23	1:A:260:VAL:H	1.49	0.75
1:A:30:ILE:HD13	1:A:42:TRP:CD2	2.22	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:HIS:HD2	4:A:321:HOH:O	1.70	0.73
1:A:182:HIS:HD2	1:A:184:ASP:H	1.45	0.64
1:A:115:LEU:HD22	1:A:118:ALA:HB2	1.79	0.63
1:A:223:ILE:HG13	1:A:225:PRO:HD3	1.79	0.63
1:A:206:GLU:OE2	1:A:210:ARG:NH1	2.34	0.60
1:A:30:ILE:HD13	1:A:42:TRP:CG	2.38	0.57
1:A:115:LEU:HD21	1:A:227:VAL:CG1	2.36	0.56
1:A:14:LYS:HE2	1:A:275:LEU:HD11	1.87	0.55
2:B:1:GLC:HO6	2:C:2:FRU:HO3	1.52	0.55
1:A:115:LEU:HD21	1:A:227:VAL:HG12	1.93	0.51
1:A:139:ARG:HG3	1:A:139:ARG:HH11	1.75	0.51
1:A:76:VAL:O	1:A:79:ILE:HG12	2.11	0.51
1:A:30:ILE:CD1	1:A:42:TRP:CD2	2.94	0.49
1:A:110:LEU:CD2	1:A:276:TYR:HB3	2.44	0.47
1:A:30:ILE:HD13	1:A:42:TRP:CE2	2.51	0.45
1:A:76:VAL:HG12	1:A:79:ILE:HG23	1.99	0.44
1:A:259:THR:HG23	1:A:260:VAL:N	2.25	0.44
1:A:46:ARG:NH1	1:A:49:GLU:OE2	2.53	0.41
1:A:70:ILE:HG22	1:A:80:HIS:CE1	2.55	0.41
1:A:136:ARG:NH2	4:A:355:HOH:O	2.53	0.41
1:A:259:THR:HG23	1:A:260:VAL:HG13	2.02	0.41
1:A:115:LEU:HD21	1:A:227:VAL:HG11	2.01	0.41
1:A:207:ARG:O	1:A:211:ASN:ND2	2.52	0.41
1:A:160:LEU:O	1:A:161:ASN:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	287/292 (98%)	279 (97%)	7 (2%)	1 (0%)	41 37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	232/228 (102%)	220 (95%)	12 (5%)	23 19

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	31	SER
1	A	46	ARG
1	A	114	GLU
1	A	136	ARG
1	A	154	LEU
1	A	163	TRP
1	A	174	ARG
1	A	188	ARG
1	A	207	ARG
1	A	263	VAL
1	A	279	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	HIS
1	A	209	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	11,11,12	0.95	0	15,15,17	1.60	2 (13%)
2	FRU	B	2	2	11,12,12	1.16	1 (9%)	10,18,18	1.29	2 (20%)
2	GLC	C	1	2	11,11,12	1.23	0	15,15,17	2.13	4 (26%)
2	FRU	C	2	2	11,12,12	1.13	1 (9%)	10,18,18	1.65	2 (20%)
2	GLC	D	1	2	11,11,12	0.88	0	15,15,17	1.54	5 (33%)
2	FRU	D	2	2	11,12,12	0.63	0	10,18,18	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	2/2/19/22	0/1/1/1
2	FRU	B	2	2	-	0/5/24/24	0/1/1/1
2	GLC	C	1	2	-	2/2/19/22	0/1/1/1
2	FRU	C	2	2	-	2/5/24/24	0/1/1/1
2	GLC	D	1	2	-	0/2/19/22	0/1/1/1
2	FRU	D	2	2	-	2/5/24/24	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	FRU	O2-C2	2.72	1.45	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	FRU	O2-C2	2.53	1.45	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GLC	C1-C2-C3	4.30	114.95	109.67
2	C	1	GLC	O4-C4-C5	4.10	119.48	109.30
2	C	1	GLC	O5-C1-C2	-3.40	105.53	110.77
2	B	1	GLC	O5-C1-C2	-3.21	105.82	110.77
2	C	2	FRU	O5-C5-C6	3.18	117.69	108.85
2	D	1	GLC	C1-O5-C5	2.99	116.24	112.19
2	B	2	FRU	O4-C4-C5	2.95	119.59	111.05
2	C	2	FRU	O2-C2-O5	2.90	115.11	109.50
2	B	1	GLC	O2-C2-C3	2.64	115.43	110.14
2	C	1	GLC	C1-O5-C5	2.61	115.73	112.19
2	D	1	GLC	O4-C4-C5	2.38	115.21	109.30
2	D	1	GLC	C1-C2-C3	-2.21	106.95	109.67
2	B	2	FRU	O1-C1-C2	-2.16	107.28	111.86
2	D	1	GLC	O2-C2-C1	2.14	113.53	109.15
2	D	1	GLC	C2-C3-C4	-2.04	107.37	110.89

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	FRU	O1-C1-C2-O2
2	B	1	GLC	O5-C5-C6-O6
2	C	1	GLC	C4-C5-C6-O6
2	B	1	GLC	C4-C5-C6-O6
2	C	2	FRU	O5-C5-C6-O6
2	C	2	FRU	C4-C5-C6-O6
2	C	1	GLC	O5-C5-C6-O6
2	D	2	FRU	O1-C1-C2-O5


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	GLC	1	0
2	C	2	FRU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	TRS	A	293	-	7,7,7	0.63	0	9,9,9	1.56	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	293	-	-	9/9/9/9	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	293	TRS	C3-C-C2	2.87	119.70	110.81
3	A	293	TRS	O2-C2-C	2.11	117.68	111.00
3	A	293	TRS	O3-C3-C	2.04	117.46	111.00

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	293	TRS	C1-C-C2-O2
3	A	293	TRS	C3-C-C2-O2
3	A	293	TRS	N-C-C2-O2
3	A	293	TRS	C1-C-C3-O3
3	A	293	TRS	C2-C-C3-O3
3	A	293	TRS	N-C-C3-O3
3	A	293	TRS	N-C-C1-O1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	293	TRS	C2-C-C1-O1
3	A	293	TRS	C3-C-C1-O1

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/292 (96%)	0.09	8 (2%) 53 51	26, 39, 58, 63	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	PHE	4.2
1	A	196	ALA	3.0
1	A	66	GLU	3.0
1	A	158	GLU	3.0
1	A	256	GLU	2.4
1	A	195	ALA	2.4
1	A	114	GLU	2.3
1	A	230	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	1	11/12	0.76	0.41	70,72,74,75	0
2	GLC	C	1	11/12	0.79	0.29	42,49,55,56	11
2	FRU	C	2	12/12	0.82	0.34	56,63,65,65	12
2	FRU	D	2	12/12	0.85	0.24	51,60,64,67	0


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FRU	B	2	12/12	0.92	0.19	37,47,55,62	0
2	GLC	B	1	11/12	0.94	0.19	40,49,61,65	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TRS	A	293	8/8	0.74	0.22	68,69,70,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.