

wwPDB X-ray Structure Validation Summary Report (i)

Oct 4, 2023 – 02:07 PM JST

PDB ID	:	8GU7
Title	:	Selective targeting of the Beclin 2-Atg14L coiled coil complex by stapled pep-
		tides promotes autophagy and endolysosomal trafficking of GPCRs
Authors	:	Qiu, X.; Zhao, Y.
Deposited on	:	2022-09-11
Resolution	:	2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

Clashscore

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.60 Å.

141614

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

3518(2.60-2.60)

Metric	Percen	tile Ranks	Value
Clashscore			4
W	orse		Better
I I	Percentile relative to all X-ray str	ructures	
0 1	Percentile relative to X-ray struc	tures of similar resolution	
Metric	Whole archive	Similar	resolution
metric	(# Entries)	(# Entries, rese	range(Å)



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	37	Total	С	Ν	Ο	S	0	0	0
	A	51	308	192	57	56	3	0	0	0
1	В	36	Total	С	Ν	Ο	S	0	0	0
	D	- 50	307	193	57	54	3		U	U

• Molecule 1 is a protein called Beclin 1-associated autophagy-related key regulator.

Chain	Residue	Modelled	Actual	Comment	Reference
А	-1	GLY	-	expression tag	UNP Q8CDJ3
А	0	PRO	-	expression tag	UNP Q8CDJ3
A	1	GLY	-	expression tag	UNP Q8CDJ3
А	2	SER	-	expression tag	UNP Q8CDJ3
A	3	GLY	-	expression tag	UNP Q8CDJ3
A	4	SER	-	expression tag	UNP Q8CDJ3
А	20	LEU	CYS	engineered mutation	UNP Q8CDJ3
A	32	SER	CYS	engineered mutation	UNP Q8CDJ3
А	34	LEU	GLY	engineered mutation	UNP Q8CDJ3
В	-1	GLY	-	expression tag	UNP Q8CDJ3
В	0	PRO	-	expression tag	UNP Q8CDJ3
В	1	GLY	-	expression tag	UNP Q8CDJ3
В	2	SER	-	expression tag	UNP Q8CDJ3
В	3	GLY	-	expression tag	UNP Q8CDJ3
В	4	SER	-	expression tag	UNP Q8CDJ3
В	20	LEU	CYS	engineered mutation	UNP Q8CDJ3
В	32	SER	CYS	engineered mutation	UNP Q8CDJ3
В	34	LEU	GLY	engineered mutation	UNP Q8CDJ3

There are 18 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called Beclin-2.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
2	С	31	Total 251	C 149	N 46	O 56	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
2	D	28	Total 227	C 137	N 41	O 49	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	1	GLY	-	expression tag	UNP A8MW95
С	2	PRO	-	expression tag	UNP A8MW95
С	3	GLY	-	expression tag	UNP A8MW95
С	4	SER	-	expression tag	UNP A8MW95
С	91	LEU	ARG	engineered mutation	UNP A8MW95
D	1	GLY	-	expression tag	UNP A8MW95
D	2	PRO	-	expression tag	UNP A8MW95
D	3	GLY	-	expression tag	UNP A8MW95
D	4	SER	-	expression tag	UNP A8MW95
D	91	LEU	ARG	engineered mutation	UNP A8MW95

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total O 1 1	0	0
3	В	1	Total O 1 1	0	0
3	D	4	Total O 4 4	0	0

SEQUENCE-PLOTS INFOmissingINFO



3 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	72.50Å 72.50Å 68.63Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.11 - 2.60	Depositor
% Data completeness	99.7 (41.11-2.60)	Depositor
(in resolution range)	· · · · · · · · · · · · · · · · · · ·	-
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$7.11 (at 2.61 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.217 , 0.240	Depositor
Wilson B-factor $(Å^2)$	41.2	Xtriage
Anisotropy	0.243	Xtriage
L-test for twinning ²	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1099	wwPDB-VP
Average B, all atoms $(Å^2)$	66.0	wwPDB-VP

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.25% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	1.06	0/309	0.88	0/406	
1	В	0.84	0/308	0.96	0/404	
2	С	1.03	0/250	0.94	0/336	
2	D	0.98	0/226	1.01	0/303	
All	All	0.98	0/1093	0.94	0/1449	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	308	0	334	1	0
1	В	307	0	339	6	0
2	С	251	0	238	4	0
2	D	227	0	222	2	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
3	D	4	0	0	0	0
All	All	1099	0	1133	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 10 close contacts within the same asymmetric unit are listed below, sorted by their



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23:ARG:HD2	2:D:21:GLU:OE2	1.94	0.68
2:D:28:LEU:O	2:D:31:VAL:HG12	1.97	0.63
2:C:35:ASN:OD1	2:C:35:ASN:N	2.32	0.61
1:B:23:ARG:HD3	2:C:21:GLU:OE2	2.11	0.51
1:B:23:ARG:HH21	1:B:26:GLN:NE2	2.09	0.51

clash magnitude.

There are no symmetry-related clashes.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

There are no protein backbone outliers to report in this entry.

4.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

4.3.3 RNA (i)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.



4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

