



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 27, 2023 – 10:13 AM EDT

PDB ID : 3H1I
Title : Stigmatellin and antimycin bound cytochrome bc1 complex from chicken
Authors : Zhang, Z.; Huang, L.; Shulmeister, V.M.; Chi, Y.I.; Kim, K.K.; Hung, L.W.; Crofts, A.R.; Berry, E.A.; Kim, S.H.
Deposited on : 2009-04-12
Resolution : 3.53 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1028 (3.60-3.48)
Clashscore	141614	1109 (3.60-3.48)
Ramachandran outliers	138981	1073 (3.60-3.48)
Sidechain outliers	138945	1074 (3.60-3.48)
RSRZ outliers	127900	1079 (3.62-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	 41% 52% 7%
1	N	446	 39% 52% 7%
2	B	441	 36% 52% 7%
2	O	441	 38% 50% 7%
3	C	380	 39% 53% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	P	380	
4	D	241	
4	Q	241	
5	E	196	
5	R	196	
6	F	110	
6	S	110	
7	G	81	
7	T	81	
8	H	77	
8	U	77	
9	I	47	
9	V	47	
10	J	61	
10	W	61	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	PEE	N	3008	-	X	-	-
11	PEE	P	3005	-	-	-	X
12	UNL	C	3015	-	-	-	X
12	UNL	P	3104	-	-	-	X
15	ANY	C	2002	X	-	-	-
15	ANY	P	3002	X	-	-	-
16	CDL	Q	3003	-	-	-	X
19	FES	E	501	-	-	X	-

2 Entry composition [i](#)

There are 20 unique types of molecules in this entry. The entry contains 32701 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 1, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	443	Total	C	N	O	S	0	0	1
			3440	2155	606	658	21			
1	N	442	Total	C	N	O	S	0	0	0
			3437	2154	605	657	21			

- Molecule 2 is a protein called UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 2, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	421	Total	C	N	O	S	0	0	0
			3141	1974	545	613	9			
2	O	422	Total	C	N	O	S	0	0	0
			3147	1977	546	614	10			

- Molecule 3 is a protein called Cytochrome b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	380	Total	C	N	O	S	0	0	0
			3020	2024	478	505	13			
3	P	379	Total	C	N	O	S	0	0	0
			3012	2019	477	504	12			

- Molecule 4 is a protein called CYTOCHROME C1, HEME PROTEIN, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	241	Total	C	N	O	S	0	0	0
			1898	1212	327	347	12			
4	Q	241	Total	C	N	O	S	0	0	0
			1898	1212	327	347	12			

- Molecule 5 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	196	Total	C	N	O	S	0	0	0
			1513	952	263	292	6			
5	R	196	Total	C	N	O	S	0	0	0
			1513	952	263	292	6			

- Molecule 6 is a protein called UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 14 KDA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	101	Total	C	N	O	S	0	0	0
			891	570	159	159	3			
6	S	101	Total	C	N	O	S	0	0	0
			891	570	159	159	3			

- Molecule 7 is a protein called UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX UBIQUINONE-BINDING PROTEIN QP-C.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
7	G	81	Total	C	N	O	0	0	0
			676	439	120	117			
7	T	79	Total	C	N	O	0	0	0
			658	430	117	111			

- Molecule 8 is a protein called UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 11 KDA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	70	Total	C	N	O	S	0	0	0
			574	350	105	114	5			
8	U	67	Total	C	N	O	S	0	0	0
			553	338	103	107	5			


- Molecule 9 is a protein called Cytochrome b-c1 complex subunit Rieske, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	46	Total	C	N	O	S	0	0	0
			285	169	58	56	2			
9	V	44	Total	C	N	O	S	0	0	1
			275	164	56	53	2			

- Molecule 10 is a protein called UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 7.2 KDA PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
10	J	61	Total	C	N	O	0	0	0
			497	321	87	89			
10	W	59	Total	C	N	O	0	0	0
			478	311	85	82			

- Molecule 11 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (three-letter code: PEE) (formula: C₄₁H₇₈NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
11	A	1	Total	C	O	P	0	0	
			21	12	8	1			
11	C	1	Total	C	N	O	P	0	0
			49	39	1	8	1		
11	E	1	Total	C	N	O	P	0	0
			50	40	1	8	1		
11	N	1	Total	O	P		0	0	
			5	4	1				
11	P	1	Total	C	N	O	P	0	0
			50	40	1	8	1		
11	P	1	Total	C	N	O	P	0	0
			49	39	1	8	1		

- Molecule 12 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	1	Total O 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	C	3	Total O 3 3	0	0
12	E	2	Total O 2 2	0	0
12	P	3	Total O 3 3	0	0
12	R	1	Total O 1 1	0	0

- Molecule 13 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	C	1	Total C Fe N O 43 34 1 4 4	0	0
13	C	1	Total C Fe N O 43 34 1 4 4	0	0
13	P	1	Total C Fe N O 43 34 1 4 4	0	0
13	P	1	Total C Fe N O 43 34 1 4 4	0	0

- Molecule 14 is STIGMATELLIN A (three-letter code: SMA) (formula: $C_{30}H_{42}O_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
14	C	1	Total	C	O			
			37	30	7	0	0	
14	P	1	Total	C	O			
			37	30	7	0	0	

- Molecule 15 is 2-METHYL-BUTYRIC ACID 3-(3-FORMYLAMINO-2-HYDROXY-BENZ OYLAMINO)-8-HEPTYL-2,6-DIMETHYL-4,9-DIOXO-[1,5]DIOXONAN-7-YL ESTER (three-letter code: ANY) (formula: C₂₉H₄₂N₂O₉).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
15	C	1	Total	C	N	O		
			37	26	2	9	0	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
15	P	1	37	26	2	9	0	0

- Molecule 16 is CARDIOLIPIN (three-letter code: CDL) (formula: $C_{81}H_{156}O_{17}P_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
16	C	1	40	21	17	2	0	0
16	D	1	50	31	17	2	0	0
16	P	1	40	21	17	2	0	0
16	Q	1	50	31	17	2	0	0

- Molecule 17 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
17	C	1	Total	C	O	0	0
			6	3	3		
17	P	1	Total	C	O	0	0
			6	3	3		

- Molecule 18 is HEME C (three-letter code: HEC) (formula: $C_{34}H_{34}FeN_4O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
18	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
18	Q	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 19 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
19	E	1	Total	Fe	S	0	0
			4	2	2		
19	R	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 20 is DIUNDECYL PHOSPHATIDYL CHOLINE (three-letter code: PLC) (formula: C₃₂H₆₅NO₈P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
20	E	1	Total 32	C 22	N 1	O 8	P 1	0	0
20	R	1	Total 32	C 22	N 1	O 8	P 1	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN I, MITOCHONDRIAL




• Molecule 1: UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN I, MITOCHONDRIAL






• Molecule 2: UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 2, MITOCHONDRIAL




• Molecule 2: UBIQUINOL-CYTOCHROME-C REDUCTASE COMPLEX CORE PROTEIN 2, MITOCHONDRIAL






• Molecule 3: Cytochrome b




• Molecule 3: Cytochrome b






• Molecule 4: CYTOCHROME C1, HEME PROTEIN, MITOCHONDRIAL




• Molecule 4: CYTOCHROME C1, HEME PROTEIN, MITOCHONDRIAL



Y237
R238
P239
P240
K241


• Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial

Chain E: 20%
33% 59% 8%




• Molecule 5: Cytochrome b-c1 complex subunit Rieske, mitochondrial

Chain R: 8%
37% 53% 10%




• Molecule 6: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 14 KDA PROTEIN


Chain F: 41% 45% 5% 8%



• Molecule 6: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 14 KDA PROTEIN

Chain S: 38% 45% 7% 8%






- Molecule 7: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX UBIQUINONE-BINDING PROTEIN QP-C




- Molecule 7: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX UBIQUINONE-BINDING PROTEIN QP-C




- Molecule 8: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 11 KDA PROTEIN




- Molecule 8: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 11 KDA PROTEIN




- Molecule 9: Cytochrome b-c1 complex subunit Rieske, mitochondrial




- Molecule 9: Cytochrome b-c1 complex subunit Rieske, mitochondrial



- Molecule 10: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 7.2 KDA PROTEIN



- Molecule 10: UBIQUINOL-CYTOCHROME C REDUCTASE COMPLEX 7.2 KDA PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	174.69Å 181.67Å 240.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 3.53 49.53 – 3.53	Depositor EDS
% Data completeness (in resolution range)	90.6 (19.99-3.53) 90.6 (49.53-3.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.23	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 3.57Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.263 , 0.306 0.250 , 0.296	Depositor DCC
R_{free} test set	2560 reflections (2.99%)	wwPDB-VP
Wilson B-factor (Å ²)	89.2	Xtrriage
Anisotropy	0.513	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 70.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.026 for k,h,-l	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	32701	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEC, PEE, GOL, UNL, ANY, PLC, FES, HEM, SMA, CDL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/3511	0.69	0/4757
1	N	0.49	0/3508	0.69	0/4753
2	B	0.43	0/3196	0.67	0/4334
2	O	0.44	0/3202	0.67	0/4343
3	C	0.59	0/3122	0.76	0/4273
3	P	0.53	0/3114	0.72	0/4263
4	D	0.52	0/1956	0.69	0/2658
4	Q	0.43	0/1956	0.67	0/2658
5	E	0.43	0/1547	0.70	3/2103 (0.1%)
5	R	0.46	0/1547	0.71	1/2103 (0.0%)
6	F	0.55	0/911	0.70	0/1219
6	S	0.49	0/911	0.65	0/1219
7	G	0.56	0/698	0.68	0/946
7	T	0.49	0/680	0.64	0/923
8	H	0.48	0/582	0.61	0/779
8	U	0.39	0/561	0.58	0/751
9	I	0.45	0/218	0.69	0/293
9	V	0.44	0/218	0.66	0/293
10	J	0.48	0/508	0.62	0/682
10	W	0.46	0/489	0.63	0/658
All	All	0.49	0/32435	0.69	4/44008 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	R	71	LEU	N-CA-C	5.94	127.03	111.00
5	E	143	GLY	N-CA-C	5.88	127.80	113.10
5	E	71	LEU	N-CA-C	5.87	126.84	111.00
5	E	70	ALA	N-CA-C	-5.04	97.40	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	104	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3440	0	3353	266	0
1	N	3437	0	3349	273	0
2	B	3141	0	3142	276	0
2	O	3147	0	3146	300	0
3	C	3020	0	3070	259	0
3	P	3012	0	3058	284	0
4	D	1898	0	1846	154	0
4	Q	1898	0	1846	161	0
5	E	1513	0	1478	136	0
5	R	1513	0	1478	120	0
6	F	891	0	893	65	0
6	S	891	0	893	73	0
7	G	676	0	659	58	0
7	T	658	0	647	63	0
8	H	574	0	548	27	0
8	U	553	0	535	38	0
9	I	285	0	239	30	0
9	V	275	0	238	30	0
10	J	497	0	490	25	0
10	W	478	0	478	33	0
11	A	21	0	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	C	49	0	72	2	0
11	E	50	0	77	1	0
11	N	5	0	0	0	0
11	P	99	0	149	6	0
12	A	1	0	0	0	0
12	C	3	0	0	0	0
12	E	2	0	0	0	0
12	P	3	0	0	0	0
12	R	1	0	0	0	0
13	C	86	0	60	20	0
13	P	86	0	60	22	0
14	C	37	0	42	3	0
14	P	37	0	42	4	0
15	C	37	0	28	3	0
15	P	37	0	29	1	0
16	C	40	0	24	2	0
16	D	50	0	44	1	0
16	P	40	0	24	2	0
16	Q	50	0	44	5	0
17	C	6	0	8	1	0
17	P	6	0	8	0	0
18	D	43	0	30	6	0
18	Q	43	0	30	4	0
19	E	4	0	0	2	0
19	R	4	0	0	1	0
20	E	32	0	38	2	0
20	R	32	0	38	3	0
All	All	32701	0	32246	2460	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 2460 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:157:ILE:HG13	3:P:158:GLY:H	1.04	1.15
2:O:157:VAL:HG23	9:V:64:LEU:HD21	1.19	1.12
5:E:119:ASP:HB3	5:E:179:ASN:ND2	1.67	1.07
2:B:157:VAL:HG23	9:I:64:LEU:HD21	1.36	1.04
1:N:231:LEU:HD23	1:N:232:PRO:HD2	1.38	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/446 (99%)	323 (73%)	94 (21%)	24 (5%)	2	19
1	N	440/446 (99%)	320 (73%)	95 (22%)	25 (6%)	1	18
2	B	419/441 (95%)	302 (72%)	77 (18%)	40 (10%)	0	8
2	O	420/441 (95%)	302 (72%)	85 (20%)	33 (8%)	1	11
3	C	378/380 (100%)	284 (75%)	72 (19%)	22 (6%)	1	18
3	P	377/380 (99%)	284 (75%)	73 (19%)	20 (5%)	2	19
4	D	239/241 (99%)	188 (79%)	37 (16%)	14 (6%)	1	17
4	Q	239/241 (99%)	186 (78%)	40 (17%)	13 (5%)	2	19
5	E	194/196 (99%)	136 (70%)	42 (22%)	16 (8%)	1	10
5	R	194/196 (99%)	136 (70%)	36 (19%)	22 (11%)	0	6
6	F	99/110 (90%)	73 (74%)	24 (24%)	2 (2%)	7	42
6	S	99/110 (90%)	73 (74%)	23 (23%)	3 (3%)	4	33
7	G	79/81 (98%)	55 (70%)	18 (23%)	6 (8%)	1	12
7	T	77/81 (95%)	52 (68%)	20 (26%)	5 (6%)	1	16
8	H	68/77 (88%)	46 (68%)	18 (26%)	4 (6%)	1	17
8	U	65/77 (84%)	41 (63%)	20 (31%)	4 (6%)	1	17
9	I	29/47 (62%)	20 (69%)	5 (17%)	4 (14%)	0	4
9	V	29/47 (62%)	21 (72%)	4 (14%)	4 (14%)	0	4
10	J	59/61 (97%)	41 (70%)	16 (27%)	2 (3%)	3	31
10	W	57/61 (93%)	36 (63%)	17 (30%)	4 (7%)	1	14
All	All	4002/4160 (96%)	2919 (73%)	816 (20%)	267 (7%)	1	15

5 of 267 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	55	ALA
1	A	72	CYS
1	A	94	GLN
1	A	159	GLN
1	A	282	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/368 (99%)	342 (94%)	23 (6%)	18	52
1	N	365/368 (99%)	343 (94%)	22 (6%)	19	53
2	B	332/347 (96%)	316 (95%)	16 (5%)	25	60
2	O	333/347 (96%)	315 (95%)	18 (5%)	22	56
3	C	329/329 (100%)	306 (93%)	23 (7%)	15	47
3	P	328/329 (100%)	304 (93%)	24 (7%)	14	46
4	D	200/200 (100%)	192 (96%)	8 (4%)	31	65
4	Q	200/200 (100%)	190 (95%)	10 (5%)	24	59
5	E	166/166 (100%)	156 (94%)	10 (6%)	19	53
5	R	166/166 (100%)	157 (95%)	9 (5%)	22	56
6	F	93/96 (97%)	85 (91%)	8 (9%)	10	40
6	S	93/96 (97%)	83 (89%)	10 (11%)	6	31
7	G	71/71 (100%)	65 (92%)	6 (8%)	10	40
7	T	69/71 (97%)	65 (94%)	4 (6%)	20	54
8	H	65/71 (92%)	63 (97%)	2 (3%)	40	71
8	U	63/71 (89%)	59 (94%)	4 (6%)	18	52
9	I	23/26 (88%)	21 (91%)	2 (9%)	10	40
9	V	23/26 (88%)	23 (100%)	0	100	100
10	J	49/49 (100%)	46 (94%)	3 (6%)	18	53
10	W	47/49 (96%)	45 (96%)	2 (4%)	29	63

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	3380/3446 (98%)	3176 (94%)	204 (6%)	19 53

5 of 204 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	N	307	PHE
3	P	47	LEU
8	U	26	GLN
1	N	420	PRO
2	O	146	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 90 such sidechains are listed below:

Mol	Chain	Res	Type
2	O	153	GLN
3	P	82	ASN
2	O	222	GLN
2	O	362	ASN
3	P	332	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 10 are unknown - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
16	CDL	C	2004	-	39,39,99	1.27	5 (12%)	45,51,111	1.13	2 (4%)
17	GOL	C	2011	-	5,5,5	1.25	0	5,5,5	0.61	0
20	PLC	E	2009	-	31,31,41	1.71	6 (19%)	37,39,49	0.71	0
18	HEC	D	501	4	32,50,50	2.19	2 (6%)	24,82,82	1.78	5 (20%)
11	PEE	P	3005	-	49,49,50	1.53	10 (20%)	52,54,55	0.87	3 (5%)
15	ANY	P	3002	-	38,38,41	1.83	9 (23%)	34,52,55	2.22	7 (20%)
20	PLC	R	3009	-	31,31,41	1.63	9 (29%)	37,39,49	0.59	0
16	CDL	P	3004	-	39,39,99	1.26	4 (10%)	45,51,111	1.13	2 (4%)
16	CDL	D	2003	-	49,49,99	1.18	4 (8%)	55,61,111	0.91	1 (1%)
13	HEM	P	502	3	41,50,50	1.78	8 (19%)	45,82,82	1.77	11 (24%)
16	CDL	Q	3003	-	49,49,99	1.15	3 (6%)	55,61,111	0.93	1 (1%)
19	FES	R	501	5	0,4,4	-	-	-	-	-
14	SMA	C	2001	-	38,38,38	1.35	4 (10%)	48,52,52	0.85	3 (6%)
13	HEM	C	502	3	41,50,50	1.95	10 (24%)	45,82,82	2.27	12 (26%)
13	HEM	P	501	3	41,50,50	1.91	9 (21%)	45,82,82	1.85	13 (28%)
13	HEM	C	501	3	41,50,50	1.84	9 (21%)	45,82,82	2.13	11 (24%)
15	ANY	C	2002	-	38,38,41	1.56	4 (10%)	34,52,55	2.22	9 (26%)
11	PEE	E	2005	-	49,49,50	1.43	9 (18%)	52,54,55	0.87	3 (5%)
11	PEE	P	3007	-	48,48,50	1.42	6 (12%)	51,53,55	0.84	3 (5%)
11	PEE	A	2008	-	20,20,50	1.81	5 (25%)	23,25,55	0.71	1 (4%)
17	GOL	P	3011	-	5,5,5	1.21	0	5,5,5	0.59	0
19	FES	E	501	5	0,4,4	-	-	-	-	-
11	PEE	N	3008	-	4,4,50	3.60	4 (100%)	6,6,55	0.66	0
11	PEE	C	2007	-	48,48,50	1.44	8 (16%)	51,53,55	0.85	3 (5%)
18	HEC	Q	501	4	32,50,50	2.34	5 (15%)	24,82,82	1.76	5 (20%)
14	SMA	P	3001	-	38,38,38	1.54	7 (18%)	48,52,52	0.82	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	CDL	C	2004	-	-	24/49/49/110	-
17	GOL	C	2011	-	-	1/4/4/4	-
20	PLC	E	2009	-	-	14/35/35/45	-
18	HEC	D	501	4	-	4/10/54/54	-
11	PEE	P	3005	-	-	27/53/53/54	-
15	ANY	P	3002	-	1/1/10/13	1/37/52/56	0/1/2/2
20	PLC	R	3009	-	-	14/35/35/45	-
16	CDL	P	3004	-	-	22/49/49/110	-
16	CDL	D	2003	-	-	29/59/59/110	-
13	HEM	P	502	3	-	4/12/54/54	-
16	CDL	Q	3003	-	-	28/59/59/110	-
19	FES	R	501	5	-	-	0/1/1/1
14	SMA	C	2001	-	-	15/34/34/34	0/2/2/2
13	HEM	C	502	3	-	5/12/54/54	-
13	HEM	P	501	3	-	6/12/54/54	-
13	HEM	C	501	3	-	6/12/54/54	-
15	ANY	C	2002	-	1/1/10/13	1/37/52/56	0/1/2/2
11	PEE	E	2005	-	-	28/53/53/54	-
11	PEE	P	3007	-	-	28/52/52/54	-
11	PEE	A	2008	-	-	15/24/24/54	-
17	GOL	P	3011	-	-	2/4/4/4	-
19	FES	E	501	5	-	-	0/1/1/1
11	PEE	C	2007	-	-	29/52/52/54	-
18	HEC	Q	501	4	-	6/10/54/54	-
14	SMA	P	3001	-	-	15/34/34/34	0/2/2/2

The worst 5 of 140 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	Q	501	HEC	C2B-C3B	-8.63	1.31	1.40
18	D	501	HEC	C2B-C3B	-8.01	1.32	1.40
18	Q	501	HEC	C3C-C2C	-7.38	1.33	1.40
18	D	501	HEC	C3C-C2C	-7.32	1.33	1.40
13	C	502	HEM	C3C-CAC	-5.89	1.35	1.47

The worst 5 of 96 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	P	3002	ANY	C25-C22-C21	8.03	135.38	111.02
15	C	2002	ANY	C25-C22-C21	7.83	134.77	111.02
13	C	502	HEM	C3B-C2B-C1B	-7.09	101.22	106.49
13	C	502	HEM	C4A-C3A-C2A	6.51	111.52	107.00
13	C	502	HEM	CMB-C2B-C1B	6.05	134.25	125.04

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
15	C	2002	ANY	C22
15	P	3002	ANY	C22

5 of 324 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	C	2007	PEE	O3P-C1-C2-O2
11	C	2007	PEE	C1-O3P-P-O2P
11	C	2007	PEE	C4-O4P-P-O3P
11	C	2007	PEE	C4-O4P-P-O2P
11	C	2007	PEE	C4-O4P-P-O1P

There are no ring outliers.

23 monomers are involved in 91 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
16	C	2004	CDL	2	0
17	C	2011	GOL	1	0
20	E	2009	PLC	2	0
18	D	501	HEC	6	0
11	P	3005	PEE	3	0
15	P	3002	ANY	1	0
20	R	3009	PLC	3	0
16	P	3004	CDL	2	0
16	D	2003	CDL	1	0
13	P	502	HEM	11	0
16	Q	3003	CDL	5	0
19	R	501	FES	1	0
14	C	2001	SMA	3	0
13	C	502	HEM	9	0
13	P	501	HEM	11	0
13	C	501	HEM	11	0
15	C	2002	ANY	3	0
11	E	2005	PEE	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	P	3007	PEE	3	0
19	E	501	FES	2	0
11	C	2007	PEE	2	0
18	Q	501	HEC	4	0
14	P	3001	SMA	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













































5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/446 (99%)	-0.23	4 (0%) 84 73	40, 85, 131, 143	0
1	N	442/446 (99%)	-0.17	8 (1%) 68 54	42, 91, 131, 143	0
2	B	421/441 (95%)	-0.12	3 (0%) 87 79	63, 107, 145, 173	0
2	O	422/441 (95%)	-0.20	6 (1%) 75 62	50, 97, 140, 150	0
3	C	380/380 (100%)	-0.75	0 100 100	17, 40, 73, 122	0
3	P	379/380 (99%)	-0.53	3 (0%) 86 75	20, 71, 106, 136	0
4	D	241/241 (100%)	-0.50	0 100 100	28, 55, 105, 118	0
4	Q	241/241 (100%)	-0.29	2 (0%) 86 75	51, 93, 136, 148	0
5	E	196/196 (100%)	0.50	39 (19%) 1 0	36, 123, 176, 180	0
5	R	196/196 (100%)	0.13	15 (7%) 13 10	50, 97, 152, 159	0
6	F	101/110 (91%)	-0.67	0 100 100	32, 57, 80, 112	0
6	S	101/110 (91%)	-0.24	1 (0%) 82 71	67, 103, 138, 161	0
7	G	81/81 (100%)	-0.43	0 100 100	38, 66, 112, 123	0
7	T	79/81 (97%)	-0.25	3 (3%) 40 29	62, 112, 176, 184	0
8	H	70/77 (90%)	-0.21	0 100 100	50, 90, 109, 132	0
8	U	67/77 (87%)	0.26	3 (4%) 33 24	129, 151, 171, 173	0
9	I	31/47 (65%)	0.64	2 (6%) 18 13	100, 143, 177, 178	0
9	V	31/47 (65%)	0.65	4 (12%) 3 3	107, 152, 194, 197	0
10	J	61/61 (100%)	-0.27	1 (1%) 72 59	57, 72, 124, 165	0
10	W	59/61 (96%)	0.04	1 (1%) 70 57	62, 89, 125, 148	0
All	All	4042/4160 (97%)	-0.24	95 (2%) 59 45	17, 87, 148, 197	0

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	E	132	TRP	5.4
5	E	134	ILE	4.5
9	I	77	ARG	4.3
5	E	121	GLN	3.9
5	E	133	VAL	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
12	UNL	E	2105	1/-	0.60	0.34	35,35,35,35	0
12	UNL	C	3015	1/-	0.63	0.60	32,32,32,32	0
12	UNL	P	3104	1/-	0.69	0.63	42,42,42,42	0
16	CDL	Q	3003	50/100	0.75	0.40	116,149,158,158	0
11	PEE	P	3005	50/51	0.77	0.46	75,97,113,115	0
11	PEE	A	2008	21/51	0.79	0.35	121,125,130,130	0
12	UNL	C	2104	1/-	0.81	0.42	36,36,36,36	0
16	CDL	D	2003	50/100	0.83	0.34	63,103,110,111	0
20	PLC	R	3009	32/42	0.83	0.48	74,93,118,118	0
11	PEE	E	2005	50/51	0.85	0.46	66,86,100,101	0
20	PLC	E	2009	32/42	0.86	0.38	37,60,94,95	0
12	UNL	A	3016	1/-	0.86	0.36	10,10,10,10	0
16	CDL	P	3004	40/100	0.87	0.27	101,108,110,110	0
11	PEE	P	3007	49/51	0.88	0.43	63,96,134,135	0
12	UNL	E	3103	1/-	0.90	0.52	32,32,32,32	0
17	GOL	C	2011	6/6	0.91	0.29	67,70,71,73	0
11	PEE	N	3008	5/51	0.91	0.36	105,105,106,106	0
11	PEE	C	2007	49/51	0.91	0.30	44,56,90,91	0
12	UNL	P	2015	1/-	0.92	0.54	25,25,25,25	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
12	UNL	C	2010	1/-	0.92	0.58	24,24,24,24	0
16	CDL	C	2004	40/100	0.92	0.24	50,72,90,92	0
12	UNL	R	2103	1/-	0.93	0.57	12,12,12,12	0
12	UNL	P	3010	1/-	0.93	0.35	24,24,24,24	0
14	SMA	P	3001	37/37	0.94	0.34	64,69,74,75	0
17	GOL	P	3011	6/6	0.94	0.43	46,48,50,50	0
15	ANY	P	3002	37/40	0.95	0.24	65,69,82,82	0
15	ANY	C	2002	37/40	0.96	0.21	27,36,48,50	0
13	HEM	P	502	43/43	0.96	0.22	49,52,70,77	0
14	SMA	C	2001	37/37	0.97	0.22	20,30,37,39	0
18	HEC	Q	501	43/43	0.97	0.20	64,68,72,76	0
18	HEC	D	501	43/43	0.98	0.16	34,39,48,55	0
13	HEM	P	501	43/43	0.98	0.21	44,49,55,56	0
13	HEM	C	501	43/43	0.98	0.20	24,32,36,39	0
13	HEM	C	502	43/43	0.98	0.20	21,28,43,49	0
19	FES	E	501	4/4	0.99	0.11	87,88,89,90	0
19	FES	R	501	4/4	0.99	0.12	55,56,58,58	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around PEE A 2008:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CDL D 2003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLC R 3009:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PEE E 2005:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLC E 2009:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CDL P 3004:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PEE N 3008:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around CDL C 2004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SMA P 3001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ANY P 3002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ANY C 2002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM P 502:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around SMA C 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)













Electron density around HEM P 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.