



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 05:57 AM BST

PDB ID : 6H1J
Title : TarP native
Authors : Guo, Y.; Stehle, T.
Deposited on : 2018-07-11
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	
1	B	345	
1	C	345	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable ss-1,3-N-acetylglucosaminyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	312	Total 2494	C 1602	N 416	O 468	S 8	0	0	0
1	B	313	Total 2519	C 1620	N 417	O 474	S 8	0	0	0
1	C	313	Total 2490	C 1600	N 416	O 466	S 8	0	1	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP A0A0H3JNB0
A	-16	ARG	-	expression tag	UNP A0A0H3JNB0
A	-15	GLY	-	expression tag	UNP A0A0H3JNB0
A	-14	SER	-	expression tag	UNP A0A0H3JNB0
A	-13	HIS	-	expression tag	UNP A0A0H3JNB0
A	-12	HIS	-	expression tag	UNP A0A0H3JNB0
A	-11	HIS	-	expression tag	UNP A0A0H3JNB0
A	-10	HIS	-	expression tag	UNP A0A0H3JNB0
A	-9	HIS	-	expression tag	UNP A0A0H3JNB0
A	-8	HIS	-	expression tag	UNP A0A0H3JNB0
A	-7	GLY	-	expression tag	UNP A0A0H3JNB0
A	-6	SER	-	expression tag	UNP A0A0H3JNB0
A	-5	LEU	-	expression tag	UNP A0A0H3JNB0
A	-4	VAL	-	expression tag	UNP A0A0H3JNB0
A	-3	PRO	-	expression tag	UNP A0A0H3JNB0
A	-2	ARG	-	expression tag	UNP A0A0H3JNB0
A	-1	GLY	-	expression tag	UNP A0A0H3JNB0
A	0	SER	-	expression tag	UNP A0A0H3JNB0
B	-17	MET	-	initiating methionine	UNP A0A0H3JNB0
B	-16	ARG	-	expression tag	UNP A0A0H3JNB0
B	-15	GLY	-	expression tag	UNP A0A0H3JNB0
B	-14	SER	-	expression tag	UNP A0A0H3JNB0
B	-13	HIS	-	expression tag	UNP A0A0H3JNB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-12	HIS	-	expression tag	UNP A0A0H3JNB0
B	-11	HIS	-	expression tag	UNP A0A0H3JNB0
B	-10	HIS	-	expression tag	UNP A0A0H3JNB0
B	-9	HIS	-	expression tag	UNP A0A0H3JNB0
B	-8	HIS	-	expression tag	UNP A0A0H3JNB0
B	-7	GLY	-	expression tag	UNP A0A0H3JNB0
B	-6	SER	-	expression tag	UNP A0A0H3JNB0
B	-5	LEU	-	expression tag	UNP A0A0H3JNB0
B	-4	VAL	-	expression tag	UNP A0A0H3JNB0
B	-3	PRO	-	expression tag	UNP A0A0H3JNB0
B	-2	ARG	-	expression tag	UNP A0A0H3JNB0
B	-1	GLY	-	expression tag	UNP A0A0H3JNB0
B	0	SER	-	expression tag	UNP A0A0H3JNB0
C	-17	MET	-	initiating methionine	UNP A0A0H3JNB0
C	-16	ARG	-	expression tag	UNP A0A0H3JNB0
C	-15	GLY	-	expression tag	UNP A0A0H3JNB0
C	-14	SER	-	expression tag	UNP A0A0H3JNB0
C	-13	HIS	-	expression tag	UNP A0A0H3JNB0
C	-12	HIS	-	expression tag	UNP A0A0H3JNB0
C	-11	HIS	-	expression tag	UNP A0A0H3JNB0
C	-10	HIS	-	expression tag	UNP A0A0H3JNB0
C	-9	HIS	-	expression tag	UNP A0A0H3JNB0
C	-8	HIS	-	expression tag	UNP A0A0H3JNB0
C	-7	GLY	-	expression tag	UNP A0A0H3JNB0
C	-6	SER	-	expression tag	UNP A0A0H3JNB0
C	-5	LEU	-	expression tag	UNP A0A0H3JNB0
C	-4	VAL	-	expression tag	UNP A0A0H3JNB0
C	-3	PRO	-	expression tag	UNP A0A0H3JNB0
C	-2	ARG	-	expression tag	UNP A0A0H3JNB0
C	-1	GLY	-	expression tag	UNP A0A0H3JNB0
C	0	SER	-	expression tag	UNP A0A0H3JNB0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Cl 2 2	0	0
2	A	5	Total Cl 5 5	0	0
2	C	3	Total Cl 3 3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	209	Total 209	O 209	0	0
4	B	236	Total 236	O 236	0	0
4	C	232	Total 232	O 232	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.37Å 95.25Å 125.47Å 90.00° 96.57° 90.00°	Depositor
Resolution (Å)	44.49 – 1.86 44.49 – 1.86	Depositor EDS
% Data completeness (in resolution range)	98.5 (44.49-1.86) 98.5 (44.49-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.170 , 0.216 0.180 , 0.226	Depositor DCC
R_{free} test set	4187 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtrriage
Anisotropy	0.448	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8192	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/2535	0.64	0/3408
1	B	0.55	0/2562	0.66	0/3445
1	C	0.54	0/2535	0.64	0/3414
All	All	0.55	0/7632	0.65	0/10267

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	100	ARG	Sidechain
1	B	259	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2494	0	2478	4	0
1	B	2519	0	2498	4	0
1	C	2490	0	2460	7	0
2	A	5	0	0	0	0
2	B	2	0	0	0	0
2	C	3	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	209	0	0	1	0
4	B	236	0	0	1	0
4	C	232	0	0	0	0
All	All	8192	0	7436	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:222:THR:HG23	1:C:224:ASN:H	1.60	0.64
1:C:9:PRO:HB3	1:C:75:PRO:HB2	1.91	0.52
1:C:19:ARG:NE	1:C:96:LEU:HD21	2.26	0.50
1:C:176:SER:HB3	1:C:225:GLN:HB3	1.98	0.45
1:A:181:ASP:N	1:A:181:ASP:OD1	2.48	0.44
1:C:173:LYS:O	1:C:180:GLU:HG3	2.18	0.43
1:B:280:ILE:HA	1:B:280:ILE:HD12	1.93	0.43
1:C:50:LEU:HD13	1:C:64:GLN:NE2	2.34	0.43
1:C:19:ARG:HG2	1:C:96:LEU:HD21	2.00	0.43
1:A:269:PHE:O	1:A:279:LYS:HE2	2.19	0.42
1:B:63:LYS:NZ	4:B:512:HOH:O	2.52	0.42
1:B:290:ILE:HD12	1:B:290:ILE:HA	1.93	0.42
1:B:74:VAL:HB	1:B:75:PRO:HD3	2.02	0.42
1:A:146:ILE:CD1	1:A:251:GLN:OE1	2.68	0.41
1:A:126:LYS:NZ	4:A:514:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/345 (89%)	299 (98%)	7 (2%)	0	100	100
1	B	307/345 (89%)	298 (97%)	9 (3%)	0	100	100
1	C	310/345 (90%)	302 (97%)	8 (3%)	0	100	100
All	All	923/1035 (89%)	899 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/313 (85%)	260 (98%)	5 (2%)	57	43
1	B	268/313 (86%)	267 (100%)	1 (0%)	91	89
1	C	260/313 (83%)	257 (99%)	3 (1%)	71	62
All	All	793/939 (84%)	784 (99%)	9 (1%)	73	65

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	LYS
1	A	120	LYS
1	A	195	TYR
1	A	259	ARG
1	A	286	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	286	PHE
1	C	204	TYR
1	C	259	ARG
1	C	286	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/345 (90%)	-0.24	4 (1%) 77 78	22, 31, 54, 83	0
1	B	313/345 (90%)	-0.20	3 (0%) 82 82	18, 28, 56, 76	0
1	C	313/345 (90%)	-0.27	2 (0%) 89 89	22, 32, 51, 80	0
All	All	938/1035 (90%)	-0.24	9 (0%) 82 82	18, 31, 55, 83	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	175	PHE	6.7
1	C	222	THR	3.1
1	A	222	THR	3.0
1	B	209	ASP	2.8
1	A	175	PHE	2.7
1	B	126	LYS	2.4
1	A	221	SER	2.1
1	A	65	LEU	2.1
1	B	18	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	C	401	1/1	0.74	0.15	55,55,55,55	0
2	CL	A	402	1/1	0.96	0.08	42,42,42,42	0
3	MG	B	401	1/1	0.97	0.17	44,44,44,44	0
2	CL	A	404	1/1	0.97	0.06	43,43,43,43	0
2	CL	C	402	1/1	0.97	0.04	41,41,41,41	0
2	CL	B	402	1/1	0.98	0.05	33,33,33,33	0
2	CL	B	403	1/1	0.98	0.07	23,23,23,23	0
2	CL	C	403	1/1	0.98	0.09	45,45,45,45	0
2	CL	C	404	1/1	0.98	0.17	54,54,54,54	0
2	CL	A	401	1/1	0.99	0.06	29,29,29,29	0
2	CL	A	405	1/1	0.99	0.06	37,37,37,37	0
2	CL	A	403	1/1	0.99	0.08	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.