



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 2, 2023 – 02:26 PM JST

PDB ID : 8H1S
Title : Crystal structure of apo-LptDE complex
Authors : Luo, Q.; Huang, Y.
Deposited on : 2022-10-03
Resolution : 3.28 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1177 (3.32-3.24)
Clashscore	141614	1044 (3.30-3.26)
Ramachandran outliers	138981	1026 (3.30-3.26)
Sidechain outliers	138945	1025 (3.30-3.26)
RSRZ outliers	127900	1141 (3.32-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	924	 3% (Poor fit), 63% (0 outliers), 25% (1 outlier), 11% (2+ outliers)
1	C	924	 4% (Poor fit), 63% (0 outliers), 26% (1 outlier), 11% (2+ outliers)
2	B	207	 8% (Poor fit), 65% (0 outliers), 15% (1 outlier), 20% (2+ outliers)
2	D	207	 13% (Poor fit), 58% (0 outliers), 22% (1 outlier), 20% (2+ outliers)

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15855 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LPS-assembly protein LptD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	824	6634	4179	1155	1283	17	0	0	0
1	A	825	6641	4181	1157	1286	17	0	0	0

- Molecule 2 is a protein called LPS-assembly lipoprotein LptE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	166	1290	790	237	259	4	0	0	0
2	B	166	1290	790	237	259	4	0	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.03Å 160.49Å 218.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.09 – 3.28 43.09 – 3.28	Depositor EDS
% Data completeness (in resolution range)	96.4 (43.09-3.28) 96.4 (43.09-3.28)	Depositor EDS
R_{merge}	0.67	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.00 (at 3.25Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.242 , 0.279 0.242 , 0.275	Depositor DCC
R_{free} test set	2952 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	80.8	Xtrriage
Anisotropy	0.315	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	15855	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.65 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.1179e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/6808	0.73	1/9234 (0.0%)
1	C	0.53	0/6799	0.73	2/9218 (0.0%)
2	B	0.47	1/1302 (0.1%)	0.65	0/1758
2	D	0.44	0/1302	0.67	0/1758
All	All	0.53	1/16211 (0.0%)	0.72	3/21968 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	93	GLU	CG-CD	5.54	1.60	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	729	PRO	N-CA-CB	6.13	110.66	103.30
1	C	486	LEU	CA-CB-CG	-5.87	101.80	115.30
1	A	729	PRO	N-CA-CB	5.56	109.98	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6641	0	6295	174	0
1	C	6634	0	6291	173	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1290	0	1277	25	0
2	D	1290	0	1277	36	0
All	All	15855	0	15140	390	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 390 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:444:VAL:HG23	1:C:869:TRP:HB2	1.45	0.94
1:A:628:GLN:OE1	1:A:688:PHE:CZ	2.21	0.93
1:C:150:ALA:HB2	1:C:185:SER:HB3	1.56	0.88
1:C:727:LEU:HD13	1:C:732:LEU:HD22	1.55	0.87
1:C:191:ASP:HB2	1:C:206:GLY:HA3	1.60	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	823/924 (89%)	756 (92%)	53 (6%)	14 (2%)	9	37
1	C	818/924 (88%)	751 (92%)	58 (7%)	9 (1%)	14	46
2	B	164/207 (79%)	156 (95%)	8 (5%)	0	100	100
2	D	164/207 (79%)	157 (96%)	7 (4%)	0	100	100
All	All	1969/2262 (87%)	1820 (92%)	126 (6%)	23 (1%)	13	44

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	96	VAL
1	C	275	ASN
1	C	829	PRO
1	A	487	PRO
1	A	829	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	703/784 (90%)	693 (99%)	10 (1%)	67 82
1	C	702/784 (90%)	694 (99%)	8 (1%)	73 85
2	B	136/169 (80%)	136 (100%)	0	100 100
2	D	136/169 (80%)	135 (99%)	1 (1%)	84 90
All	All	1677/1906 (88%)	1658 (99%)	19 (1%)	73 85

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	535	ASP
1	A	916	TYR
1	A	918	GLN
1	A	747	TYR
2	D	146	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	628	GLN
1	A	819	HIS
1	A	918	GLN
1	A	891	GLN
2	D	120	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	825/924 (89%)	0.27	31 (3%) 40 38	52, 73, 107, 151	0
1	C	824/924 (89%)	0.31	41 (4%) 28 27	58, 79, 114, 140	0
2	B	166/207 (80%)	0.78	16 (9%) 8 8	57, 83, 124, 135	0
2	D	166/207 (80%)	0.95	27 (16%) 1 2	61, 89, 129, 143	0
All	All	1981/2262 (87%)	0.39	115 (5%) 23 23	52, 77, 116, 151	0

The worst 5 of 115 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	179	ALA	6.4
2	B	185	ALA	6.2
2	D	182	LEU	5.5
1	A	174	THR	5.4
1	A	148	ASP	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.