



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 26, 2023 – 06:32 AM EDT

PDB ID : 3H3F
Title : Rabbit muscle L-lactate dehydrogenase in complex with NADH and oxamate
Authors : Bujacz, A.; Bujacz, G.; Swiderek, K.; Paneth, P.
Deposited on : 2009-04-16
Resolution : 2.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	 4% 58% 34% 8%
1	B	331	 2% 66% 30% 5%
1	C	331	 2% 56% 36% 8%
1	D	331	 2% 57% 37% 6%
1	E	331	 6% 50% 42% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	331	<p>2% 61% 32% 7%</p>
1	G	331	<p>% 62% 31% 7%</p>
1	H	331	<p>6% 52% 40% 8%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OXM	F	333	-	-	X	-
4	ACT	C	334	-	-	X	-
4	ACT	D	336	-	-	X	-

2 Entry composition


There are 5 unique types of molecules in this entry. The entry contains 22218 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lactate dehydrogenase A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	331	2562	1635	442	471	14	0	1	0
1	B	331	2568	1641	442	471	14	0	2	0
1	C	331	2559	1633	441	471	14	0	0	0
1	D	331	2575	1644	442	475	14	0	4	0
1	E	331	2564	1636	441	473	14	0	1	0
1	F	331	2563	1636	441	471	15	0	1	0
1	G	331	2571	1639	442	476	14	0	3	0
1	H	331	2568	1640	441	473	14	0	2	0

- Molecule 2 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: C₂₁H₂₉N₇O₁₄P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			6	2	1	3		
3	B	1	Total	C	N	O	0	0
			6	2	1	3		
3	C	1	Total	C	N	O	0	0
			6	2	1	3		
3	D	1	Total	C	N	O	0	0
			6	2	1	3		
3	E	1	Total	C	N	O	0	0
			6	2	1	3		
3	F	1	Total	C	N	O	0	0
			6	2	1	3		
3	G	1	Total	C	N	O	0	0
			6	2	1	3		
3	H	1	Total	C	N	O	0	0
			6	2	1	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	F	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0
4	H	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	146	Total O 146 146	0	0
5	B	202	Total O 202 202	0	0
5	C	143	Total O 143 143	0	0
5	D	139	Total O 139 139	0	0
5	E	116	Total O 116 116	0	0
5	F	153	Total O 153 153	0	0
5	G	158	Total O 158 158	0	0
5	H	139	Total O 139 139	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-lactate dehydrogenase A chain




- Molecule 1: L-lactate dehydrogenase A chain



- Molecule 1: L-lactate dehydrogenase A chain






- Molecule 1: L-lactate dehydrogenase A chain



- Molecule 1: L-lactate dehydrogenase A chain



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.50Å 85.28Å 138.53Å 98.49° 91.67° 111.59°	Depositor
Resolution (Å)	60.00 – 2.38 30.96 – 2.38	Depositor EDS
% Data completeness (in resolution range)	89.7 (60.00-2.38) 89.7 (30.96-2.38)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.160 , 0.207 0.164 , 0.207	Depositor DCC
R_{free} test set	4887 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtrriage
Anisotropy	0.092	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.034 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	22218	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAI, OXM, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	0/2614	0.96	5/3534 (0.1%)
1	B	0.92	0/2622	1.00	3/3545 (0.1%)
1	C	0.88	0/2605	0.96	3/3523 (0.1%)
1	D	0.87	1/2639 (0.0%)	0.97	5/3568 (0.1%)
1	E	0.86	0/2613	0.96	2/3534 (0.1%)
1	F	0.91	1/2613 (0.0%)	0.96	0/3533
1	G	0.88	0/2632	0.97	3/3559 (0.1%)
1	H	0.83	1/2622 (0.0%)	0.95	4/3546 (0.1%)
All	All	0.88	3/20960 (0.0%)	0.97	25/28342 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	184	CYS	CB-SG	-5.16	1.73	1.81
1	F	191	GLU	CG-CD	5.16	1.59	1.51
1	D	126	TYR	CB-CG	-5.04	1.44	1.51

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	268	ARG	NE-CZ-NH2	-8.94	115.83	120.30
1	B	235	ASP	CB-CG-OD1	6.82	124.44	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	42	ASP	CB-CG-OD1	6.26	123.94	118.30
1	H	170	ARG	NE-CZ-NH2	-6.17	117.21	120.30
1	E	156	ARG	NE-CZ-NH1	5.87	123.23	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	13	LYS	Peptide
1	D	282	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2562	0	2644	199	0
1	B	2568	0	2655	124	0
1	C	2559	0	2639	180	0
1	D	2575	0	2653	175	0
1	E	2564	0	2643	204	0
1	F	2563	0	2644	161	0
1	G	2571	0	2647	146	0
1	H	2568	0	2648	200	0
2	A	44	0	27	4	0
2	B	44	0	27	2	0
2	C	44	0	27	4	0
2	D	44	0	27	13	0
2	E	44	0	27	10	0
2	F	44	0	27	4	0
2	G	44	0	27	2	0
2	H	44	0	27	1	0
3	A	6	0	2	1	0
3	B	6	0	2	0	0
3	C	6	0	2	1	0
3	D	6	0	2	0	0
3	E	6	0	2	1	0
3	F	6	0	2	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	6	0	2	1	0
3	H	6	0	2	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
4	C	8	0	6	2	0
4	D	20	0	15	2	0
4	E	20	0	15	1	0
4	F	8	0	6	1	0
4	G	12	0	9	0	0
4	H	16	0	12	0	0
5	A	146	0	0	24	0
5	B	202	0	0	18	0
5	C	143	0	0	13	0
5	D	139	0	0	21	0
5	E	116	0	0	30	0
5	F	153	0	0	16	0
5	G	158	0	0	23	0
5	H	139	0	0	21	0
All	All	22218	0	21474	1332	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 1332 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:223:LYS:H	1:C:223:LYS:CD	1.25	1.45
1:E:317:LYS:HZ2	1:E:317:LYS:CB	0.88	1.41
1:A:106:LEU:O	1:A:106:LEU:HD23	1.26	1.28
1:H:308:THR:CG2	1:H:310:GLU:HG2	1.62	1.28
1:E:100:GLN:CG	1:E:101:GLU:H	1.42	1.27

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/331 (100%)	306 (93%)	22 (7%)	2 (1%)	25	34
1	B	331/331 (100%)	314 (95%)	17 (5%)	0	100	100
1	C	329/331 (99%)	307 (93%)	22 (7%)	0	100	100
1	D	333/331 (101%)	307 (92%)	25 (8%)	1 (0%)	41	53
1	E	330/331 (100%)	307 (93%)	21 (6%)	2 (1%)	25	34
1	F	330/331 (100%)	303 (92%)	25 (8%)	2 (1%)	25	34
1	G	332/331 (100%)	311 (94%)	20 (6%)	1 (0%)	41	53
1	H	331/331 (100%)	306 (92%)	23 (7%)	2 (1%)	25	34
All	All	2646/2648 (100%)	2461 (93%)	175 (7%)	10 (0%)	34	46

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	ALA
1	E	192	HIS
1	D	221	ALA
1	E	312	GLU
1	A	104	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/284 (100%)	252 (88%)	33 (12%)	5	6
1	B	286/284 (101%)	261 (91%)	25 (9%)	10	14
1	C	284/284 (100%)	252 (89%)	32 (11%)	6	7
1	D	288/284 (101%)	254 (88%)	34 (12%)	5	6
1	E	285/284 (100%)	249 (87%)	36 (13%)	4	5
1	F	285/284 (100%)	255 (90%)	30 (10%)	7	8

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	287/284 (101%)	254 (88%)	33 (12%)	5	6
1	H	286/284 (101%)	246 (86%)	40 (14%)	3	3
All	All	2286/2272 (101%)	2023 (88%)	263 (12%)	5	6

5 of 263 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	103	GLU
1	H	131	LYS
1	H	322	LEU
1	D	103	GLU
1	D	75	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	110	GLN
1	F	107	ASN
1	H	297	ASN
1	G	232	GLN
1	E	163	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

39 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	G	334	-	3,3,3	0.66	0	3,3,3	1.87	1 (33%)
2	NAI	D	332	-	42,48,48	1.80	8 (19%)	47,73,73	1.67	10 (21%)
4	ACT	H	335	-	3,3,3	0.84	0	3,3,3	1.50	0
3	OXM	B	333	-	5,5,5	3.54	1 (20%)	4,6,6	1.32	1 (25%)
4	ACT	G	336	-	3,3,3	0.94	0	3,3,3	1.71	2 (66%)
3	OXM	D	333	-	5,5,5	2.82	1 (20%)	4,6,6	1.81	2 (50%)
3	OXM	E	333	-	5,5,5	2.24	1 (20%)	4,6,6	1.75	2 (50%)
3	OXM	F	333	-	5,5,5	2.35	1 (20%)	4,6,6	1.89	1 (25%)
4	ACT	F	334	-	3,3,3	0.88	0	3,3,3	1.07	0
4	ACT	E	336	-	3,3,3	0.90	0	3,3,3	1.65	1 (33%)
2	NAI	B	332	-	42,48,48	1.72	7 (16%)	47,73,73	1.92	13 (27%)
4	ACT	C	334	-	3,3,3	0.92	0	3,3,3	2.26	2 (66%)
4	ACT	F	335	-	3,3,3	0.87	0	3,3,3	1.90	1 (33%)
4	ACT	D	338	-	3,3,3	1.03	0	3,3,3	0.88	0
4	ACT	H	336	-	3,3,3	1.00	0	3,3,3	1.43	0
4	ACT	H	334	-	3,3,3	0.77	0	3,3,3	1.54	0
4	ACT	H	337	-	3,3,3	0.78	0	3,3,3	1.73	2 (66%)
4	ACT	A	334	-	3,3,3	1.17	0	3,3,3	1.60	1 (33%)
4	ACT	D	337	-	3,3,3	0.96	0	3,3,3	0.86	0
4	ACT	D	336	-	3,3,3	0.95	0	3,3,3	1.14	0
2	NAI	G	332	-	42,48,48	1.36	5 (11%)	47,73,73	1.90	15 (31%)
4	ACT	B	334	-	3,3,3	1.12	0	3,3,3	1.90	2 (66%)
2	NAI	E	332	-	42,48,48	1.64	6 (14%)	47,73,73	1.43	8 (17%)
2	NAI	F	332	-	42,48,48	1.55	7 (16%)	47,73,73	1.79	10 (21%)
2	NAI	H	332	-	42,48,48	1.63	6 (14%)	47,73,73	1.73	11 (23%)
4	ACT	E	334	-	3,3,3	1.03	0	3,3,3	0.91	0
3	OXM	G	333	-	5,5,5	3.53	1 (20%)	4,6,6	1.74	2 (50%)
2	NAI	A	332	-	42,48,48	1.48	6 (14%)	47,73,73	1.59	9 (19%)
3	OXM	H	333	-	5,5,5	2.35	1 (20%)	4,6,6	1.67	1 (25%)
3	OXM	C	333	-	5,5,5	2.63	1 (20%)	4,6,6	1.20	1 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	D	335	-	3,3,3	0.88	0	3,3,3	1.62	1 (33%)
4	ACT	E	337	-	3,3,3	1.27	0	3,3,3	0.93	0
4	ACT	G	335	-	3,3,3	0.71	0	3,3,3	1.75	2 (66%)
3	OXM	A	333	-	5,5,5	3.42	1 (20%)	4,6,6	1.83	2 (50%)
4	ACT	E	335	-	3,3,3	1.01	0	3,3,3	2.05	2 (66%)
4	ACT	D	334	-	3,3,3	0.98	0	3,3,3	1.12	0
2	NAI	C	332	-	42,48,48	1.46	5 (11%)	47,73,73	1.73	9 (19%)
4	ACT	E	338	-	3,3,3	0.88	0	3,3,3	1.88	1 (33%)
4	ACT	C	335	-	3,3,3	1.04	0	3,3,3	1.68	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAI	C	332	-	-	5/25/72/72	0/5/5/5
2	NAI	E	332	-	-	10/25/72/72	0/5/5/5
2	NAI	B	332	-	-	5/25/72/72	0/5/5/5
2	NAI	D	332	-	-	9/25/72/72	0/5/5/5
2	NAI	F	332	-	-	8/25/72/72	0/5/5/5
3	OXM	B	333	-	-	0/3/4/4	-
2	NAI	H	332	-	-	8/25/72/72	0/5/5/5
3	OXM	A	333	-	-	0/3/4/4	-
3	OXM	G	333	-	-	0/3/4/4	-
3	OXM	D	333	-	-	0/3/4/4	-
3	OXM	E	333	-	-	2/3/4/4	-
3	OXM	F	333	-	-	0/3/4/4	-
2	NAI	A	332	-	-	3/25/72/72	0/5/5/5
3	OXM	H	333	-	-	3/3/4/4	-
2	NAI	G	332	-	-	11/25/72/72	0/5/5/5
3	OXM	C	333	-	-	0/3/4/4	-

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	333	OXM	C1-C2	-7.71	1.45	1.55
3	B	333	OXM	C1-C2	-7.45	1.46	1.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	333	OXM	C1-C2	-7.35	1.46	1.55
3	D	333	OXM	C1-C2	-5.98	1.47	1.55
2	B	332	NAI	C4N-C3N	-5.79	1.38	1.49

The worst 5 of 116 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	332	NAI	N3A-C2A-N1A	-6.54	118.45	128.68
2	F	332	NAI	N3A-C2A-N1A	-6.29	118.85	128.68
2	C	332	NAI	N3A-C2A-N1A	-5.59	119.94	128.68
2	G	332	NAI	N3A-C2A-N1A	-5.50	120.08	128.68
2	H	332	NAI	C1D-N1N-C2N	-5.25	112.36	121.11

There are no chirality outliers.

5 of 64 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	332	NAI	O4B-C4B-C5B-O5B
2	D	332	NAI	C5D-O5D-PN-O2N
2	E	332	NAI	C5D-O5D-PN-O1N
2	E	332	NAI	C5D-O5D-PN-O2N
2	E	332	NAI	O4D-C4D-C5D-O5D

There are no ring outliers.

17 monomers are involved in 48 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	332	NAI	13	0
3	E	333	OXM	1	0
3	F	333	OXM	3	0
2	B	332	NAI	2	0
4	C	334	ACT	2	0
4	F	335	ACT	1	0
4	D	336	ACT	2	0
2	G	332	NAI	2	0
2	E	332	NAI	10	0
2	F	332	NAI	4	0
2	H	332	NAI	1	0
4	E	334	ACT	1	0
3	G	333	OXM	1	0
2	A	332	NAI	4	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	333	OXM	1	0
3	A	333	OXM	1	0
2	C	332	NAI	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/331 (100%)	-0.16	13 (3%) 39 42	8, 18, 34, 52	0
1	B	331/331 (100%)	-0.31	8 (2%) 59 60	7, 16, 29, 43	0
1	C	331/331 (100%)	-0.19	6 (1%) 68 70	9, 18, 33, 46	0
1	D	331/331 (100%)	-0.23	7 (2%) 63 65	8, 19, 33, 46	0
1	E	331/331 (100%)	-0.05	20 (6%) 21 24	7, 20, 38, 53	0
1	F	331/331 (100%)	-0.28	6 (1%) 68 70	8, 17, 33, 43	0
1	G	331/331 (100%)	-0.30	3 (0%) 84 84	8, 19, 29, 43	0
1	H	331/331 (100%)	-0.00	20 (6%) 21 24	10, 20, 35, 53	0
All	All	2648/2648 (100%)	-0.19	83 (3%) 49 51	7, 19, 33, 53	0

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	100	GLN	6.0
1	A	102	GLY	5.6
1	H	106	LEU	4.8
1	E	99	GLN	4.8
1	H	102	GLY	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OXM	D	333	6/6	0.69	0.38	60,62,63,64	0
4	ACT	H	336	4/4	0.77	0.37	45,46,46,47	0
4	ACT	E	337	4/4	0.83	0.25	37,38,38,38	0
2	NAI	D	332	44/44	0.85	0.27	25,32,44,45	0
4	ACT	B	334	4/4	0.86	0.17	30,30,32,33	0
3	OXM	A	333	6/6	0.86	0.24	34,34,35,36	0
4	ACT	F	334	4/4	0.86	0.12	38,38,40,40	0
4	ACT	A	334	4/4	0.86	0.16	31,32,33,33	0
4	ACT	E	335	4/4	0.87	0.17	32,32,33,33	0
4	ACT	G	336	4/4	0.88	0.21	42,44,44,44	0
3	OXM	H	333	6/6	0.88	0.18	11,16,20,22	0
4	ACT	E	336	4/4	0.89	0.18	41,42,43,43	0
3	OXM	E	333	6/6	0.90	0.26	34,36,37,37	0
4	ACT	C	335	4/4	0.90	0.19	42,43,43,43	0
4	ACT	D	334	4/4	0.90	0.39	40,41,41,41	0
4	ACT	E	338	4/4	0.92	0.25	39,40,40,40	0
4	ACT	D	335	4/4	0.92	0.18	41,41,42,42	0
4	ACT	E	334	4/4	0.93	0.15	37,38,38,38	0
2	NAI	E	332	44/44	0.93	0.15	20,29,33,36	0
4	ACT	D	336	4/4	0.93	0.28	39,40,40,40	0
4	ACT	D	338	4/4	0.93	0.12	32,34,34,34	0
2	NAI	A	332	44/44	0.95	0.12	20,25,29,31	0
4	ACT	F	335	4/4	0.95	0.19	37,38,38,39	0
4	ACT	C	334	4/4	0.95	0.24	27,28,28,28	0
4	ACT	D	337	4/4	0.95	0.32	32,33,33,34	0
4	ACT	H	337	4/4	0.95	0.33	46,46,46,47	0
4	ACT	G	335	4/4	0.96	0.20	39,39,39,40	0
4	ACT	H	334	4/4	0.97	0.07	31,32,32,32	0
4	ACT	H	335	4/4	0.97	0.10	33,33,34,34	0
3	OXM	C	333	6/6	0.97	0.11	12,15,17,17	0
4	ACT	G	334	4/4	0.97	0.10	30,31,31,33	0
2	NAI	B	332	44/44	0.98	0.10	7,14,17,18	0
3	OXM	B	333	6/6	0.98	0.10	9,14,16,17	0
2	NAI	F	332	44/44	0.98	0.09	6,13,17,19	0
2	NAI	G	332	44/44	0.98	0.10	6,13,18,22	0
2	NAI	H	332	44/44	0.98	0.07	10,19,23,27	0
3	OXM	F	333	6/6	0.98	0.07	14,16,17,18	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAI	C	332	44/44	0.99	0.11	6,12,15,17	0
3	OXM	G	333	6/6	0.99	0.09	5,6,7,9	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NAI E 332:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAI A 332:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAI B 332:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAI F 332:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAI G 332:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAI H 332:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.